

Bound Together or Falling Apart? Foraging Association in Red Knots

Pratik R Gupte^{1,2}, Selin Ersoy^{1,2} & Allert I Bijleveld²



1 Groningen Institute for Evolutionary Life Sciences, University of Groningen, 2 Department of Coastal Systems, Netherlands Institute for Sea Research || Contact: p.r.gupte@rug.nl, @pratikunterwegs

Background

1 Waders such as red knots *Calidris canutus* gather in large non-breeding flocks in the Wadden Sea, where they forage on intertidal mudflats

2 Waders are highly social, but they have no 'friends' — no consistent non-random association between individuals (cite, cite)

3 ATLAS allows high frequency tracking of multiple knots, and calculation of pairwise 'coherence' (cite)




^aRed knot — ^bWadden Sea mudflats


^cStudy site — ^dIsland of Griend

^eTracking towers — ^fKnot positions

Data handling




- 1 Obtained the 'lifetime' track
- 2 Identified tidal intervals from water-level data (cite)
- 3 Classified data by tidal interval
- 4 Classified data by hours since high tide



- 1 Distances between time-matched positions
- 2 Count distances < proximity threshold
- 3 Calculate coherence

Good title for results



Acknowledgements Thanks to AIB, Eva Kok, Frank van Marseveen for the tracking data. Thanks to Cpt. Bram Fey, *RV Navicula* crew, & NIOZ technical staff for installing the ATLAS system.

References Myers 1983, Conklin & Calwell 2007, Ekman 1967