


# The ecology and evolution of movement in fluctuating landscapes

**Pratik R Gupte**

February 2019



**Supervised by**

Prof. Franz J Weissing  
Dr. Allert I Bijleveld  
Prof. Ton Groothuis  
Prof. Theunis Piersma

© Pratik R Gupte

**Modelling Adaptive Response Mechanisms Group,**  
Groningen Institute for Evolutionary Life Sciences,  
University of Groningen.

**Bijleveld Lab,**  
Department of Coastal Systems,  
Royal Netherlands Institute for Sea Research.

#### **Contact information**

Groningen Institute for Evolutionary Life Sciences  
University of Groningen  
Nijenborgh 7-5172.0583, 9747 AG Groningen  
Netherlands  
p.r.gupte@rug.nl

#### **Supervisors**

Prof. FJ Weissing  
Dr. AI Bijleveld  
Prof. AGG Groothuis  
Prof. T Piersma

#### **Promotors**

FJ Weissing  
AGG Groothuis

#### **Funding**

Adaptive Life Programme – Groningen Institute for Evolutionary Life Sciences  
Short term causes and long term consequences of adaptations to environmental change

**Cover image** Red knot Y4NGNG (laterally inverted) courtesy Benjamin Gnep; movement track of knot with transmitter 471 courtesy Allert Bijleveld; background generated in R.



**university of  
 groningen**



# Contents

<b>Summary</b>	<b>4</b>
<b>1 Introduction</b>	<b>5</b>
1.1 Animal movement . . . . .	5
1.2 Movement as personality . . . . .	7
1.3 Modelling movement . . . . .	9
<b>2 About this project</b>	<b>10</b>
2.1 Abstract models . . . . .	10
2.2 Wader models and data . . . . .	12
<b>3 Towards mechanistic movement models</b>	<b>12</b>
3.1 Modelling landscapes . . . . .	13
3.1.1 Concept: Spatial predictability . . . . .	13
3.1.2 Concept: Temporal predictability and change . . . . .	14
3.1.3 Practice: Modelling spatio-temporal change in landscapes . . . . .	14
3.2 Modelling agents . . . . .	15
3.2.1 Concept: Agent Based Models . . . . .	15
3.2.2 Concept: Agent-resource and agent-agent interactions . . . . .	17
3.2.3 Concept: Modelling interference competition . . . . .	17
3.2.4 Practice: Agent Based Models using Artificial Neural Networks . . . . .	18
3.3 Confronting models with data . . . . .	19
<b>Timeframe</b>	<b>20</b>
<b>References</b>	<b>21</b>

## Summary

The **movement of animals** has always fascinated humans, but modern biology has only now come to actively study movement as the linchpin of many processes. As movement studies seek to unify phenomena across scales, they must also contend with the idea that there are **consistent individual differences** within many animal populations. These differences, whose evolution, development, and maintenance are still under debate, are estimated to have population level consequences. One system in which consistent individual differences have been identified is that of **red knots** *Calidris canutus*, which forage on intertidal mudflats and vary along an axis of exploratory behaviour. Knots' and other foragers' movement is often strongly influenced by which **spatio-temporal change** in the often unpredictable resource landscape. This system inspires questions about how movement strategies evolve in such landscapes, how they affect the landscapes on which they evolve, as well as how the interaction of the two gives rise to emergent phenomena such as structured communities.

This essay, an introduction to my work over the next two years, is organised over the following parts: first, the **Introduction** contains key concepts in the fields of *Animal movement*, *Movement as personality*, and *Modelling movement*. The second section, **About this project**, lays out the questions I want to investigate for my PhD. The third section, **Towards a model of wader movement**, explains the conceptual and practical aspects of the models and empirical methods I propose to implement towards the goals I set myself. The **Timeframe** outlines the activities I will undertake over the period 2018 – 2022 in terms of research, personal academic training, co-supervision of students, and academic output, culminating in a thesis submission in May 2022, and a thesis defence provisionally in September 2022.

For my PhD, I will attempt to break new ground for the field of animal movement by examining the evolution of movement strategies, with a special focus on the evolution of consistency in fluctuating environments. First, using simple, abstract models, I will examine under which conditions of environmental spatial predictability and temporal change one can expect the evolution of distinct movement types. Second, I will examine the ecological consequences of consistency in movement behaviour in heterogeneous landscapes. Third, I will tailor the former models to investigate consistency in exploratory behaviour in my model system, the shorebirds known as red knots (*Calidris canutus*). Fourth, and finally, I will examine whether red knots tracked in the wild show movement types, and thus test the predictions of theory.

# 1 Introduction

## 1.1 Animal movement

Animal movement across spatio-temporal scales – from seasonal migration, to breeding aggregations, to diurnal cycles – have been an important field of human knowledge throughout recorded history, and likely long before. Even in more modern times, the two most successful paradigms of the natural sciences’ century of insight, evolution by natural selection (Darwin & Wallace 1858) and island biogeography (MacArthur & Wilson 2001), rely at least in part on the movement of animals to make their point. Explicit treatment of the subject began with theoretical developments on the movement of freely moving and ideally rational agents. This led to nearly simultaneous advances in the understanding of the distribution of foragers at the landscape scale (Fretwell & Lucas 1970), and the topology of groups of selfish agents (Hamilton 1971). Theory on population spatial distributions soon incorporated fine-scale phenomena such as individual interactions and population structure, and was already a mature field with strong empirical support when reviewed by Sutherland (1996). Theoretical advances in the study of fine-scale movement itself, on the other hand, have lagged behind the measurement of the movement of individual animals (Holyoak *et al.* 2008).

The recently emerged movement ecology paradigm (MEP; Nathan *et al.* 2008 see Fig. 1) has sought to fill this gap, understanding “movement itself as the focal theme...by providing a unifying framework and common tools” (Nathan 2008). This paradigm urges practitioners to attempt to answer three questions: 1. *Why* do organisms move? 2. *Where* do organisms decide to move? 3. *How* do organisms move? This links (1) external stimuli, such as fear of predators (Laundré *et al.* 2001), (2) sensory capacity, which translates into navigational ability and search strategies (e.g. Bartumeus & Levin 2008), and (3) life-history dependent physiology and internal state (e.g. Fryxell *et al.* 2008). Movement ecology’s rise has coincided with the beginning of “a golden age of animal tracking science” in the form of large-scale and high-resolution animal tracking (Hussey *et al.* 2015; Kays *et al.* 2015). Biologging, i.e., the remote measurement of instantaneous organism state (Cooke *et al.* 2004), and high-resolution animal tracking in particular, have led to remarkable insights into the detailed sub-units of movement behaviour, such as the energetics of predation and locomotion (see e.g. Williams *et al.* 2004; Scantlebury *et al.* 2014).

Despite advances in empirical methods, animal movement studies are still subject to limitations of scale. Among these are emergent group-size effects (Tunstrøm *et al.* 2013), which are difficult to quantify empirically notwithstanding recent advances (Handegard *et al.* 2012; Kays *et al.* 2015; Strandburg-Peshkin *et al.* 2015; Dhanjal-Adams *et al.* 2018); a distinct problem when animal grouping is near-universal (Krause & Ruxton 2002). The impracticality of research across scales of organisation (eg. physiology to population) has similarly confined most movement stud-




Figure 1.1: The movement ecology paradigm places observed movement (*path*:  $U$ ) as both a consequence of prior factors, and a contributor to future ones. From Nathan et al. (2008).

ies to answering at most one MEP question at a time (but see Fryxell *et al.* 2008; Strandburg-Peshkin *et al.* 2015; Curtin *et al.* 2018). This is linked to the choice of study system, which among vertebrates (birds, especially) has been restricted by the technical challenge of hewing to the 3% body-mass rule (Naef-Daenzer *et al.* 2001). Finally, vertebrate movement has been captured by the movement *ecology* paradigm because of the difficulty in studying movement at evolutionary timescales in these systems, compared to the relative tractability of invertebrate communities (see recent eco-evo ex. Venkateswaran *et al.* 2017). Here, simulation models can help to shine “the light of ecology” on evolutionary biology (Grant & Grant 2008), by uncovering the importance of individual interactions for emergent phenomena (e.g. Hildenbrandt *et al.* 2010; Tunstrøm *et al.* 2013), the surprising extrinsic drivers of movement mode (Guttal *et al.* 2012), the evolutionary effects of higher trophic levels (Ioannou *et al.* 2012), and the evolution of movement rules (de Jager *et al.* 2011; Netz 2017).

## 1.2 Movement as personality

The work cited above may be safely claimed to make at least one of two assumptions; the first is variation in traits around a ‘golden mean’ is unimportant noise. The second is the assumption of optimality (Fretwell & Lucas 1970) entailing infinite behavioural plasticity. Neither of these assumptions is very well founded. While the first is methodologically convenient, theory from Darwin (1859) onwards has recognised the pivotal role of variation; models find that differences among individuals can have population level consequences (e.g. Pruitt & Riechert 2011). The second assumption is squarely challenged by the study of consistent individual differences in behaviour. The field of animal personality and behavioural syndromes (Sih *et al.* 2004a, b) and its consequences (Dingemanse & Réale 2005) seeks to quantify between and within individual consistency in behavioural responses “through time or across situations”. Variation along two or more axes of personality (Réale *et al.* 2007) may be correlated into a ‘behavioural type’ (examples in Sih & Bell 2008; Bell *et al.* 2009); for example, ‘boldness’ and ‘aggression’ form two of the major axes of personality, and are often quantified together as a syndrome (Carter *et al.* 2013).

Behavioural ecology has only slowly come to terms with the idea that such individual differences within populations could be advantageous (Wilson 1998). Individual differences have been explained in terms of mechanisms engaged during development (Wolf & Weissing 2010; Groothuis & Trillmich 2011), trade-offs in life-history (Wolf *et al.* 2007a, b), and the feedback loops created between individual state and behaviour (Wolf & Weissing 2010) as prominent proximate causes. Evolutionary explanations of these differences and especially their consistency have remained elusive because the default expectation has been that plasticity is more adaptive (Wilson 1998). Wolf & Weissing (2010) have since shown that behavioural types may yet achieve comparable fitness outcomes if selection is frequency-dependent, and rarer types have a selective advantage (Maynard Smith 1982). Spatio-temporal variation, a ubiquitous feature of environments (Levin 1992), may have a major role to play when it hosts a population where infinite adaptability is costly or habitat choice limited (Wolf & Weissing 2010). In such cases, individuals

may specialise for particular spatial or temporal conditions, maintaining global variability, while local variability may be maintained through imperfect estimation of current habitat state (Wolf & Weissing 2010; Wolf & McNamara 2012). However, there is little doubt that behavioural types have significant eco-evolutionary consequences for populations (Sih *et al.* 2012; Wolf & Weissing 2012).

The movement ecology paradigm places animal movement at the end of a pipeline fed into by physiology, stimuli, and cognition (*state*, Wolf & Weissing 2010, 2012), and it is unsurprising that populations might have evolved behavioural polymorphisms for movement, or *movement types* (*sensu* Wolf & Weissing 2010; Getz *et al.* 2015; Netz 2017). Indeed, movement types are inherent predictions of previous theoretical work that did not necessarily consider movement as its central theme (e.g. Watson & Miller 1971; Lima & Zollner 1996). Animal tracking has since confirmed such individual consistency in movement across scales, from local movements (Austin *et al.* 2004; Leclerc *et al.* 2016) to dispersal (Duckworth & Badyaev 2007; Clobert *et al.* 2009; Cote J. *et al.* 2010; Duckworth & Sockman 2012). While Wolf & Weissing (2010) suggested that some spatio-temporal environmental variability might be causal to consistency evolution, later work showed that such structure can be created from uniformity by individuals themselves, and thus may be co-incidental rather than causal in relation to movement strategy evolution (de Jager *et al.* 2011; Getz *et al.* 2015; Netz 2017). In general, this points to the interplay between individuals' movement and the landscapes – and begs the question of how the rate and predictability of landscape change (*sensu lato* Botero *et al.* 2015) might affect the evolution of movement types. Spiegel *et al.* (2017) have demonstrated that this link between movement types (*fast* and *slow*; see parallels with earlier Wolf *et al.* 2007a) and fitness outcomes in landscapes that vary in spatial *predictability* can have consequences at the population level – from social networks to community composition (see Fig. 2).




Figure 1.2: Spiegel *et al.* (2017) lay out a framework linking personality, movement, and resulting interactions. Personality may be initially measured in terms of movement, but the underlying causes can impact larger-scale processes such as population spatial structure.



### 1.3 Modelling movement

Many aspects of personality and movement can be explored using the waders *Charadrii*, and especially the sandpipers Scolopacidae as a model system. Waders are a ubiquitous group of long-lived birds characterised by their littoral foraging on buried macrozoobenthic prey, and their often extreme flyway-channelled migration between generally disjointed and strongly seasonal global distributions (Boere & Stroud 2006; Gill *et al.* 2009; Piersma & Bonan 2019). Sandpipers are well known for their huge wintering fission-fusion flocks (Myers 1983; Conklin & Colwell 2008), and as symbols of traditional landscapes (Colwell 2018). Waders (hereafter referring only to scolopacids) eking out their precarious existence at the waterline are thus poised to be the new canary in the coalmine: harbingers of global environmental change (Piersma & Lindström 2004; Wikelski & Tertitski 2016; Fitzpatrick & Senner 2018) which has swiftly advanced from ‘near’ to ‘here’ (IPCC 2018).

Wader movement fulfils many of the requirements one would expect from a system that integrates animal personality and movement studies. First, wader movement has been extensively investigated using a variety of approaches at nearly all spatio-temporal scales: small spatio-temporal scales such as foraging (van Gils *et al.* 2003; van Gils 2010), large spatio-temporal scales such as migration (Buehler *et al.* 2006; Piersma 2011), proximate causes for large-scale phenomena (Lank *et al.* 2003; Ydenberg *et al.* 2004; Ruthrauff *et al.* 2013, 2018), and ultimate causes for small-scale phenomena (van Gils *et al.* 2006; Kraan *et al.* 2009a). Of the waders, red knots *Calidris canutus* are among the best studied, with a wealth of biological information, from an understanding of physiology and its interaction with exploratory personality (Bijleveld *et al.* 2014; Mathot *et al.* 2017), to knowledge of group dynamics across scales (Bijleveld *et al.* 2010, 2012a, 2015b), to the effect of foragers on their resource landscapes (Bijleveld *et al.* 2015a).

Many theoretical studies increasingly employ agent-based approaches, which were developed to overcome two flaws in mathematical models (Huston *et al.* 1988). First, in order to prevent an unreasonably large number of parameters (a disadvantage of ABMs), earlier models had abstracted populations to “golden mean” values that elided individual differences (see previous section) essential to evolutionary and ecological processes. The second issue was the assumption of global interactions, i.e., that all individuals affect all others equally, ignoring central problems of scale (Levin 1992) and local interactions (see Legendre 1993) in ecology. ABMs have since been standardised in terminology (Grimm *et al.* 2006, 2010), and are widespread in ecology (DeAngelis 2018). In such models, individual organisms are each modelled separately and allowed to make decisions based upon various inputs, akin to real decision making processes (DeAngelis & Diaz 2019).

In animal ecology, ABMs are especially relevant in modelling movement (reviewed in DeAngelis & Mooij 2005), and have been used extensively (Spiegel *et al.* 2013, 2017; Getz *et al.* 2015, 2016), including to model red knots (van Gils 2010). Simple evolutionary algorithms (see Bäck 1996) as implemented in Getz *et al.* (2015) and Netz (2017) that predicate the reproductive output (replicates) of each agent on some measure of fitness, such as net intake, can add eco-evolutionary

dynamics to otherwise simple ABMs. This allows the modelling of movement across scales and contexts, from small-scale within-patch dynamics, such as competition (see Keddy 2001 for a primer) and the investigation of emergent large-scale evolutionary dynamics, such as community composition (*movement guilds*; Getz *et al.* 2015), and alternative strategies (Netz 2017). A major advance may be made by having the decisions of agents be the output of a ‘black-box’ process in the form of artificial neural networks (ANNs; see Lek & Guégan 1999; Enquist & Ghirlanda 2013; Netz 2017 for example implementation). Neural network approaches allow for surprising results – such as the evolution of rich movement dynamics and alternative strategies, even when beginning from largely similar populations (Netz 2017).

## 2 About this project

In this section, I outline my approaches to examine the *evolution of movement types* (*sensu* Wolf & Weissing 2010; Getz *et al.* 2015) in the context of *environmental predictability and rate of change* (as in Botero *et al.* 2015), using the *red knot system* as a reference (Bijleveld *et al.* 2012a, 2014, 2015a, b; Oudman *et al.* 2016, 2018). Wolf & Weissing (2010) lay out the theoretical basis for the expectation that consistent behavioural differences could arise in spatially structured landscapes, while Botero *et al.* (2015) describe how environments in certain regimes of temporal variability and predictability can give rise to similar polymorphisms. Getz *et al.* (2015) and Netz (2017) show how virtual agents can evolve movement types while also structuring initially featureless landscapes. Bijleveld and colleagues (above) provide empirical data on the foraging and movement ecology of red knots that can inform modeling approaches and which may be used to test model predictions.

### 2.1 Abstract models

I will begin with an abstract approach and some basic questions. Modelling agent movement as the output of an ANN which takes environmental cues as inputs, I will first investigate how different regimes of landscape patchiness and temporal variability influence the evolution of movement types. Specifically, I will be interested in the following:

1. ***How many movement types are evolved under different regimes of spatial predictability and variability?*** This question is aimed at unifying the Wolf & Weissing (2010) and Botero *et al.* (2015) approaches at broad evolutionary scales. Recent mechanistic models based on ANNs suggest that genotype-phenotype mapping is not as clear-cut as in Botero and colleagues’ work, and that polymorphisms (there identified as diversified hedged bets) may arise under a wider set of conditions in populations of fixed size. I will then examine the behaviour of agents (movement) in response to environment quality (resource landscape), with the idea being to identify the conditions under which different strategies evolve.

2. ***What is the link between behaviour and labile physiological state?*** Drawing inspiration from recent work in the red knot system that suggests a strong role for behaviour-sociality-physiology feedbacks (discussed in Wolf & Weissing 2010; Bijleveld *et al.* 2014; Oudman *et al.* 2016; Mathot *et al.* 2017), this question is aimed at investigating whether physiology constrains agents in behavioural parameter space. For example, when movement is made dependent on instantaneous state, it may be constrained by energy reserves, leading to a state-behaviour feedback. Further, intake may face digestive constraints that can strongly affect behaviour (van Gils & Piersma 2004). I will investigate whether these feedbacks can produce behaviour-physiology clustering previously proposed in this system (Bijleveld *et al.* 2015b).
  
3. ***How do movement type frequencies develop over ecological and evolutionary time-scales?*** Foragers moving about a resource landscape can modify or introduce spatial patterns over time (de Jager *et al.* 2011; Getz *et al.* 2015; Netz 2017), yet effects such as facilitation remain understudied. Empirical results suggest top-down landscape structuring, such as by depletion, can be rapid (Bijleveld *et al.* 2015a). Starting with initially unstructured landscapes, I will investigate how structure is modified using a relaxed version of **(1)** in which agents deplete their landscape at different rates. A natural consequences might be that foragers specialised to particular regimes of landscape structure could see a change in their profitability multiple times within their lives. When **(1)** includes the depletion element of **(3)**, agents might be expected to evolve a median (bet-hedged) phenotype that ensures profitability across landscape structures. This question is then aimed at investigating both the landscapes and agents:
  - a) ***How does landscape structure change over ecological and evolutionary time?*** In this scenario, I fall back upon the red knot system as a guide, that foragers are not always present on the landscape and the resource has time to regenerate. Such seasonal systems are widespread making this a reasonable scenario. I will investigate the development of landscape parameters over ecological and evolutionary time (such as the autocorrelation range, or patch size; see Legendre 1993) when agents are themselves evolving upon such landscapes.
  - b) ***How does the number and phenotype of movement types change over evolutionary time?*** Agents' consumption rates (set to be identical) could lead to different rates of landscape change in the scenario outlined above. Foragers would then evolve to maximise fitness in both space and time. As landscapes traverse the axis of patch size (autocorrelation range, or predictability) due to agent foraging, the spatial conditions that evolved types in **(1)** might recur multiple times. This should allow the investigation of whether there is a change in selection pressures on movement types, the time-scale at which it occurs, and whether it is mediated by agents' effect on the landscape.
  - c) ***Do critical transitions occur in the number or profitability of movement types?*** As agent populations evolve movement types rather than bet-hedged phenotypes, these types

may yet show sufficient flexibility to behaviourally buffer against decreasing profitability as the landscape shifts to a regime to which they are poorly adapted. This change may occur within individuals at ecological scales (as resources deplete or change over a season), and at the population level (as landscapes change over evolutionary time). Such systems, where the response (here, movement, or fitness) is initially poorly responsive to the driver (landscape structure, such as patch size), may then undergo abrupt transitions (Scheffer 2009) to an alternative stable state. Behavioural changes relating to movement strategy occur in wader systems (Oudman *et al.* 2018), but the process is not fully understood. I will look into whether changes in such systems can then be characterised as critical transitions. The same approach can be applied to the landscape, to study whether landscapes catastrophically shift to another state (as in van de Koppel *et al.* 1997; Jefferies *et al.* 2006).

## 2.2 Wader models and data

Following the steps above, I will modify the models to simulate the red knot system. Important conceptual changes include the addition of regular environmental change in the form of the tidal cycle. This feature first limits resource landscape availability to agents. Second, it forces agent movement, further constraining the range within which agent movement types can specialise. For example, a very efficient movement type in a model elaborated above (**1**) would be to remain largely sedentary. This is not an option in tidal systems as most waders cannot swim. Important practical changes will be the parameterisation of the model following empirical data or best estimates on resource landscape structure and red knot physiology (body size, energy requirements, perception range).

Finally, I will turn to the empirical red knot tracking data and the benthic sampling data to investigate whether the models sketched above correspond to reality. The following questions suggest themselves.

1. ***Do red knots show movement types that can be identified from a combination of data sources?*** This question entails cooperation with my co-PhD student. Using scores from behavioural assays, state variables such as gizzard mass or body mass, and movement variables such as patch-switching or residence time, I will aim to determine whether knots evolved in models occupy comparable parameter spaces to real birds.
2. ***Do red knots show assortative association based on movement type?*** Here, I investigate expectations from Spiegel *et al.* (2017), that association and the strength of social networks is strongly influenced by personality and the ‘clumped-ness’ of the landscape. This question is suitable to both an experimental and field tracking approach.

## 3 Towards mechanistic movement models

In each of these sub-sections, *Modelling landscapes* and *Modelling agents*, I will lay out how I propose to go about modelling the systems I have described above. I first describe the concept with some justification (usually drawn from red knot, or other avian systems), and then delve into the practical considerations of implementation, again seeking to justify my choices. In each case, I begin with the more abstract model, and then elaborate on a model based on red knots. It is important to note that while initial, abstract models are based on previous work, later, more complex models rely on these abstract models. Should the abstract models produce different dynamics from the ones expected from theory, the implementation and predictions of later models may have to be adjusted accordingly.

Another important point is that these are intended to be ‘mechanistic models of intermediate complexity’. For example, they will explicitly model a mechanism of cue sensing, which was only implicitly assumed in Botero *et al.* (2015). Complexity, though significantly greater than Botero and colleagues’ work, will have an upper limit set near that of Netz (2017) or van den Berg & Weissing (2015).

### 3.1 Modelling landscapes

#### 3.1.1 Concept: Spatial predictability

Spatial pattern and scale matter in ecology (Levin 1992), and foragers benefit from responding to spatial structure in resource landscapes in empirical and theoretical studies (Benhamou 1992; Walsh 1996; Klaassen *et al.* 2006; van Gils *et al.* 2006; van Gils 2010; Bijleveld *et al.* 2016; Oudman *et al.* 2018). Botero *et al.* (2015) model a very abstract predictability as well as variation in time – however, space-time substitution Blois *et al.* (2013) has found wide use in testing landscape-scale predictions in ecological systems (e.g. Hirota *et al.* 2011; Staver *et al.* 2011), and allows the leveraging of extensive empirical sampling of resource landscapes at small (e.g. Bijleveld *et al.* 2012b) and large scales (e.g. Huete *et al.* 2002). In a spatially structured landscape, autocorrelation is an important measure (Legendre 1993), and has been widely interpreted as corresponding to patch size (Kraan *et al.* 2009a, b; van Gils 2010; Bijleveld *et al.* 2016; Oudman *et al.* 2018). This idea is easily illustrated by creating neutral landscapes with varying autocorrelation ranges – landscapes with a high autocorrelation range have larger patches, and are more predictable (see Figure 3). In the wader-mudflat system, the mechanism underlying patch size does not need to be explicitly modelled to implement patch sizes. It suffices to understand that macrobenthic abundance is controlled by bottom-up processes over which agents have little influence.

### 3.1.2 Concept: Temporal predictability and change

The assumption of resources being independent of foraging agents is plainly ridiculous (van de Koppel *et al.* 1997; Jefferies *et al.* 2006; Bijleveld *et al.* 2015a). However, the proposed system is also far from agent-influenced resource generation (see e.g. le Roux *et al.* 2018). This leaves depletion as the main effect of agents on their landscape. Depletion then constitutes one important mechanism of temporal change in model landscapes. Agent-independent mechanisms such as seasonal growth and decline, or within-resource interactions (such as competition and facilitation) are the other major mechanism of temporal change in landscapes.

In initial models, agents will *not* deplete the landscape. However, this would result in the trivial solution of agents remaining stationary at the first point where resources were above some threshold value. Models without depletion must then include some variation in the resource landscape such that agents are forced to move. This can be achieved by having the landscape change, either deterministically or stochastically (see parallel with Botero *et al.* 2015). The first approximates changes such as might be driven by seasonality, while the second is more akin to resource redistribution. The first approach allows near perfect prediction of resource values between times  $t$  and  $t+1$ . In the second case, if the ‘redistribution’ is highly variable, i.e., values at a coordinate at a time  $t$  have a wide range of correlation with values at time  $t+1$ , temporal predictability is reduced. The two extreme cases allow the examination of the interacting effects of spatial and temporal predictability, and identification of the more interesting regime for further investigation.

In a model tailored to waders, the situation is compounded by another source of temporal variation: the tidal cycle. This highly periodic phenomenon both gives and takes away: water restricts exploitation of parts of the landscape (most waders cannot swim; but see interesting exception of phalaropes *Phalaropus*). However, the tide also allows waders to use their pressure-sensitive bills (Piersma *et al.* 1998) to find prey in the waterlogged substrate. This results in gradients of information on prey availability at different scales. At the landscape scale, the optimal prey search area is at the waterline where the substrate is exposed yet sufficiently wet to allow knots to probe for food. At smaller scales, this prey-sensing mechanism severely localises information on potential intake, i.e., a foraging wader can only directly assess the quality of its immediate neighbourhood. The dynamics of coastal systems cause further complications – but also create opportunities – when modelling agents, and these are discussed later.

### 3.1.3 Practice: Modelling spatio-temporal change in landscapes

Both resource and tidal landscapes may be implemented in continuous space. While implemented in some models (e.g. Spiegel *et al.* 2013, 2017), continuous space creates a mismatch of resolution between simulations and empirical data, which is an important consideration in this project. Since data from benthic sampling can be converted into predicted intake rate rasters

with a maximum resolution of 10 m (see method in Bijleveld *et al.* 2012b; examples in Oudman *et al.* 2018), the resource landscape is best modelled as a two-dimensional square grid of side  $n$  cells, a common approach (Nolet & Mooij 2002; van Gils 2010; Getz *et al.* 2015; Netz 2017). Grid values can be modelled using flexible tools (Sciaini *et al.* 2018) as Gaussian random fields (Turner & Gardner 2015; Kery & Royle 2019). Working on the logic of infinite extent (Nolet *et al.* 2006), the grid boundaries are periodic. Landscapes can thus easily be assigned an autocorrelation regime. Grid cells in this landscape are initialised with certain values of resource between 0.0 and 1.0. This allows the subsequent mapping of empirical values using a variety of functions, such as the sigmoidal (Gershenfeld 1999).

The tidal landscape's elevation is easily modelled as a distance or edge gradient (Etherington *et al.* 2015). This creates a region of high elevation which is always exposed, approximating islands such as Griend where waders roost. This model will use a circular distance gradient that avoid issues arising from landscape edges. Temporal change in both water level and resource landscapes can initially be considered to be the output of a sine function with a wavelength  $R$  (as in Botero *et al.* 2015): the “relative timescale of environmental change” of each. For water level, the wavelength would be the number of discrete model timesteps contained in one unit of ecological time. Since the tidal cycle (approx. 13 hours off Griend in 2017; *unpublished data*) is a prominent feature, it is intuitive to also consider this a unit of ecological time, with a single non-breeding season comprising some 6 – 8 months (approx. 450 tidal cycles). Each season would then be a unit of evolutionary time. In the case of the resource landscape, the wavelength can be set to vary across a range to mimic either seasonal (renewal each season) or long-term (climatic cycles) dynamics, corresponding very well to Botero *et al.* (2015).

## 3.2 Modelling agents

### 3.2.1 Concept: Agent Based Models

I will implement ABMs for abstract foragers, and then for foragers similar to waders. While waders are an ideal system in which to study foraging, they are specialised upon a small subset of resources and ecological conditions that will constrain more realistic models. For a general forager, it is important to be able to gain some key information: its position, the profitability of its position, and its potential future positions (*why* and *where* to move; Nathan *et al.* 2008). An agent will move when its current location cannot sustain it, possibly due to some metabolic requirement (Barraquand & Benhamou 2008). There is a strong interplay between *where* to move and *how* to move when agents are capable of more than one movement mode, such as in birds which can switch from cursorial to volant. While the exact mechanism may be abstracted, movement carries a cost, and this cost must be taken into account when deciding when and where to move (Charnov 1976). Movement decisions are made as the result of explicitly written functions (e.g. Getz *et al.* 2015), but this entails and often elides implicit assumptions about the functional response of agents to their environment. This environment may include other agents, allowing




Figure 3.1: Examples of Gaussian random field neutral landscapes generated in R following methods from Sciaini et al. (2018). **(a)** Landscapes of side 100 cells, with increasing autocorrelation range (numbers above panels). Larger autocorrelation ranges result in smoother transitions between areas of high (blue) and low (red) resources, and thus larger patches. **(b)** Spatial autocorrelation in simulated landscapes; each panel corresponds to the one directly above in **(a)**. Compare with figures in Oudman et al. (2018).



for facilitation by local enhancement (Beauchamp 2013).

### **3.2.2 Concept: Agent-resource and agent-agent interactions**

In the models sketched above I began with the assumption that while especially large or numerous foragers (ecosystem engineers) are capable of transforming their resource landscapes (Laundré *et al.* 2001; Jefferies *et al.* 2006; le Roux *et al.* 2018), this cannot be expected for all consumer-resource systems. However, there is evidence from observational studies at small spatial scales (approx. 0.067 km<sup>2</sup>) that medium-sized foragers can significantly deplete their resources (Guillemette *et al.* 1996; Jefferies *et al.* 2006). Markedly smaller waders too can deplete resources over small spatio-temporal scales (Székely & Bamberger 1992; van Gils *et al.* 2003; Bijleveld *et al.* 2015a). Thus, in more realistic models, I will allow foragers to deplete their landscapes. This depletion will replace the sinusoidal temporal variation I proposed for initial abstract models, and will more realistically approximate seasonal dynamics where resource replenishment occurs in a single growing period, and further declines are largely due to harvesting.

Depletion introduces the prospect of exploitative competition (Keddy 2001), where agents affect each other by consuming shared resources. Its inclusion in the simplest models has stark eco-evolutionary consequences – for example, Getz *et al.* (2015) showed that larger population sizes (and thus higher competition) resulted in more movement types and a more rapid rate of evolution. I propose to take this into account by on a grid cell proportional to the number of agents on the cell. Interference competition is widely seen in waders (Goss-Custard 1980), yet is also more challenging to include in discrete time models (Vahl 2006). I will leave this aspect of wader ecology out of the model (but see next sub-section).

### **3.2.3 Concept: Modelling interference competition**

I intend to co-supervise a master's student with an interest in evolutionary game theory, and modelling skills. This master's project will extend the work of Vahl (2006), and take into account the following: first, that mechanisms can strongly influence the dynamics of games (van den Berg & Weissing 2015), second, that competition is often state dependent (van Gils & Piersma 2004), and third, that both direct and indirect competition (Vahl *et al.* 2005; Bijleveld *et al.* 2012a) can have consequences for agent space-use (Vahl & Kingma 2007). I envision the following modules for this project: individual based models following Vahl's ideas, with pairwise interactions determining some contest outcome that translates to fitness. These will be implemented in an evolutionary system, where successful agents replicate, either with fixed or flexible population size. Second, some state dependence of behaviour to examine behaviour – physiology trade-offs. Third, to have the agent decisions be the output of neural networks. Fourth, implementation in an initially limited spatial system (with the option of expansion), such as a two dimensional 3

$\times 3$  grid, or a single dimensional vector. In this final, more realistic system, movement would be an option, allowing interesting dynamics between direct competition, state-dependence of behaviour, and spatial distribution in a game theoretical framework.

### 3.2.4 Practice: Agent Based Models using Artificial Neural Networks

Agents in ABMs possess attributes, including knowledge of their coordinate position on the grid  $(x, y)$  and some proxy of internal state (energy reserves; e.g. van Gils 2010). Other attributes may comprise the overall phenotype of the agent. For example, Getz *et al.* (2015); Getz *et al.* (2016) draw three parameters that correspond broadly to giving-up value  $\rho$ , competitiveness  $\delta$ , and sociability  $\alpha$  from suitable distributions, and assign them to agents. These values are implastic through the agent's life, and while a useful starting point, assumes behavioural consistency. The ANN approach allows for far richer dynamics, where agents assess their environment (described above) and make a single decision about what move to make in the next time-step. This environment may include landscape values and/or other agents (Netz 2017). These cues are the activations of the agent's ANN nodes, while state variables can be mapped to node weights and biases. Agent movement is often limited to the Moore neighbourhood of size 1, i.e., from agent grid position to the eight cells around it in each timestep (Getz *et al.* 2015; Netz 2017). Movements of unit distance in unit time assume constant speed; clearly unfounded since real animals such as knots can achieve at least two broad movement speeds by flying or walking (Bijleveld *et al.* *unpublished data*). The ANN output sketched above – next grid position – allows agents to choose their movement distance at each time step. At an ecological scale, this first allows the investigation of individual consistency in step length, and then an examination of whether movement types have characteristic step lengths, e.g. that are some whole number multiple of the autocorrelation range. Movement carries a travel cost, which is subtracted from energy reserves at each time-step.

In more complex models, cues of where best to forage need not be equally available to foraging agents. Knots can sense only very local macrobenthos availability using their pressure sensitive bills (Piersma *et al.* 1998). At intermediate and local scales, knots may use public information from other foragers (Bijleveld *et al.* 2015b), while it is to be assumed that across scales, knots are able to avoid deep water. This creates a cue gradient: at very small range, agents have much more reliable information than at larger ones, where local enhancement due to conspecific presence may play a greater role (Beauchamp 2013). Thus agents in such models may evolve movement types that trade the cost of poor information for the cost of exploitative competition, and vice-versa, all while balancing the cost of travel with the intake from the landscape. Evolutionary models require that traits are inherited, implying reproduction, birth and death. Agent fitness is best modelled as some function, possibly sigmoidal, of net intake, i.e., agents do have an upper limit to the offspring produced; for example, many sandpipers produce a clutch of four eggs (Piersma & Bonan 2019), and this is a realistic upper limit. Agents must therefore also have some mechanism approximating death; this can be modelled as energy reserves reaching zero, after

which the agent dies. Agents must reproduce at some time — in systems worldwide, most species have a fixed breeding season, and this can be modelled as agents that survive until the next ecological time-step producing offspring. The value of energy reserves at which agents decide to reproduce may also be allowed to evolve, possibly evolving interesting life-history strategies (Wolf *et al.* 2007a; as in Wolf & McNamara 2012).

### 3.3 Confronting models with data

Models are always wrong, but they do yield insights into real systems. Empirical data on the red knot system is collected both from the agents (MacCurdy *et al.* 2015 in ; examples in Bijleveld *et al.* 2016; Oudman *et al.* 2018) and their resource landscape (Bijleveld *et al.* 2012b). These data include morphometric measures, as well as movement measures which may be obtained using a number of tools now available for the processing of animal tracking data (most recent review by Joo *et al.* 2019). Coupled with behavioural scores from aviary experiments (currently ongoing at NIOZ; see e.g. Bijleveld *et al.* 2012a, 2015b), these can be used to cluster individuals. These clusters can then be compared with movement types evolved in simulations. I aim to given the resource landscape structure experienced in the Wadden Sea by *islandica* red knots, and then to examine how well these predictions hold up in the face of data.

Individual associations of red knots are not expected to be non-random, consistent with the pattern for other waders (Myers 1983; Conklin & Colwell 2008). However, that expectation might yet hold for movement types rather than specific individuals, with assortative association within or between types in different environmental regimes (Spiegel *et al.* 2017). The strength and nature of associations can be quantified from simulated movement tracks, and then tested using extensive tracking data.

## Timeframe

This is an approximate timeframe. Events farther in the future are more likely to change. Note that field visits and review meetings are not included, and that flexible plans are marked by asterisks.

Id	Year	Research	Training/conferences	Co-supervision	Output
<b>2018</b>					
1	May – Jun	Visit NIOZ			
2	Jul – Sep	Field visits; knot associations	Computing cluster crs.; <i>Biomove symp.</i>		Poster Biomove symp.
3	Oct – Dec	Lit. rev. for intro. essay	<i>C++ programming; PhD course CRI</i>	B.Sc. essays (2)	Crs. proj. (2); Intro. Essay
<b>2019</b>					
<b>Building abstract models</b>					
4	Jan – Mar	Move. + competition model descriptions	GRS/GRC; <i>Sci. Integrity Course</i>	M.Sc. student 1, 2	Final Intro. Essay
5	Apr – Jun	Proj. 1. Movement mods. variable envmnt.; Proj. 2. Interference mods.	Visit NIOZ	M.Sc. student 1, 2	Proj. 1+2. methods
6	Jul – Sep	Proj. 3. Aviary scores ~ field movement	Collaborators visit*	M.Sc. student 2	Prep. Proj. 3. m/s.
7	Oct – Dec	Analyse data Proj. 1+2.	Machine learning course*		Proj. 3 m/s.
<b>2020</b>					
<b>Building shorebird models</b>					
8	Jan – Mar	Proj. 4. Develop shorebird models	<i>Modelling in R</i> course;	M.Sc. student 3, 4*	Proj. 1+2 m/s.
9	Apr – Jun	Finalise shorebird models	Collaborators visit*	M.Sc. student 3, 4*	Shorebird mods. methods
10	Jul – Sep	Proj. 5. M.Sc. students' project topics	ISBE/ESEB	M.Sc. student 3, 4*	
11	Oct – Dec	Compare models with data			
<b>2021</b>					
<b>Drawing down</b>					
12	Jan – Mar	Compare models and data			Proj. 4+5 m/s prep.
13	Apr – Jun	Paper writing	Visit collaborator/future lab*		Proj. 4+5 m/s
14	Jul – Sep	Paper writing			
15	Oct – Dec	Thesis and paper writing	GSSE <i>Career Perspectives Courses</i>		
<b>2022</b>					
<b>Preparing to leave</b>					
16	Jan – Mar	Thesis and paper writing	Post-doc grant writing		Post-doc grant appl.
17	Apr – May	PhD ends; thesis submission			PhD thesis
18	Sep	Preparation & PhD defence			

# References

- Austin, D., Bowen, W.D. & McMillan, J.I. (2004). Intraspecific variation in movement patterns: Modeling individual behaviour in a large marine predator. *Oikos*, 105, 15–30.
- Bäck, T. (1996). *Evolutionary Algorithms in Theory and Practice: Evolution Strategies, Evolutionary Programming, Genetic Algorithms*. Oxford University Press.
- Barraquand, F. & Benhamou, S. (2008). Animal movements in heterogeneous landscapes: Identifying profitable places and homogeneous movement bouts. *Ecology*, 89, 3336–3348.
- Bartumeus, F. & Levin, S.A. (2008). Fractal reorientation clocks: Linking animal behavior to statistical patterns of search. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 19072–7.
- Beauchamp, G. (2013). *Social Predation: How Group Living Benefits Predators and Prey*. Elsevier.
- Bell, A.M., Hankison, S.J. & Laskowski, K.L. (2009). The repeatability of behaviour: A meta-analysis. *Animal Behaviour*, 77, 771–783.
- Benhamou, S. (1992). Efficiency of area-concentrated searching behaviour in a continuous patchy environment. *Journal of Theoretical Biology*, 159, 67–81.
- Bijleveld, A.I., Egas, M., Gils, J.A.V. & Piersma, T. (2010). Beyond the information centre hypothesis: Communal roosting for information on food, predators, travel companions and mates? *Oikos*, 119, 277–285.
- Bijleveld, A.I., Folmer, E.O. & Piersma, T. (2012a). Experimental evidence for cryptic interference among socially foraging shorebirds. *Behavioral Ecology*, 23, 806–814.
- Bijleveld, A.I., Gils, J.A. van, Meer, J. van der, Dekinga, A., Kraan, C. & Veer, H.W. van der *et al.* (2012b). Designing a benthic monitoring programme with multiple conflicting objectives. *Methods in Ecology and Evolution*, 3, 526–536.
- Bijleveld, A.I., MacCurdy, R.B., Chan, Y.-C., Penning, E., Gabrielson, R.M. & Cluderay, J. *et al.* (2016). Understanding spatial distributions: Negative density-dependence in prey causes predators to trade-off prey quantity with quality. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20151557.
- Bijleveld, A.I., Massourakis, G., van der Marel, A., Dekinga, A., Spaans, B. & van Gils, J.A. *et al.* (2014). Personality drives physiological adjustments and is not related to survival. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20133135.
- Bijleveld, A.I., Twietmeyer, S., Piechocki, J., Gils, J.A. van & Piersma, T. (2015a). Natural selection by pulsed predation: Survival of the thickest. *Ecology*, 96, 1943–1956.
- Bijleveld, A.I., van Gils, J.A., Jouta, J. & Piersma, T. (2015b). Benefits of foraging in small groups: An experimental study on public information use in red knots *Calidris Canutus*. *Behavioural Processes, Cause and Function in Behavioral Biology: A tribute to Jerry Hogan*, 117, 74–81.
- Blois, J.L., Williams, J.W., Fitzpatrick, M.C., Jackson, S.T. & Ferrier, S. (2013). Space can substitute for time in predicting climate-change effects on biodiversity. *Proceedings of the National Academy of Sciences*, 110, 9374–9379.
- Boere, G.C. & Stroud, D. (2006). The flyway concept: What it is and what it isn't. In: *Waterbirds Around the World: A Global Overview of the Conservation, Management and Research of the World's Waterbird Flyways*. Scottish Natural Heritage, Edinburgh, UK, pp. 40–47.
- Botero, C.A., Weissing, F.J., Wright, J. & Rubenstein, D.R. (2015). Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences*, 112, 184–189.
- Buehler, D.M., Baker, A.J. & Piersma, T. (2006). Reconstructing palaeoflyways of the late Pleistocene and early Holocene Red Knot *Calidris canutus*, 15.
- Carter, A.J., Feeney, W.E., Marshall, H.H., Cowlshaw, G. & Heinsohn, R. (2013). Animal personality: What are behavioural ecologists measuring? *Biological Reviews*, 88, 465–475.
- Charnov, E.L. (1976). Optimal foraging, the marginal value theorem. *Theoretical Population Biology*, 9, 129–136.
- Clobert, J., Galliard, J.-F.L., Cote, J., Meylan, S. & Massot, M. (2009). Informed dispersal, heterogeneity in animal disper-

- sal syndromes and the dynamics of spatially structured populations. *Ecology Letters*, 12, 197–209.
- Colwell, M.** (2018). *Curlew Moon*. 1st edn. Harper Collins.
- Conklin, J.R. & Colwell, M.A.** (2008). Individual associations in a wintering shorebird population: Do Dunlin have friends? *Journal of Field Ornithology*, 79, 32–40.
- Cooke, S.J., Hinch, S.G., Wikelski, M., Andrews, R.D., Kuchel, L.J. & Wolcott, T.G. et al.** (2004). Biotelemetry: A mechanistic approach to ecology. *Trends in Ecology & Evolution*, 19, 334–343.
- Cote J., Clobert J., Brodin T., Fogarty S. & Sih A.** (2010). Personality-dependent dispersal: Characterization, ontogeny and consequences for spatially structured populations. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 4065–4076.
- Curtin, N.A., Bartlam-Brooks, H.L.A., Hubel, T.Y., Lowe, J.C., Gardner-Medwin, A.R. & Bennitt, E. et al.** (2018). Remarkable muscles, remarkable locomotion in desert-dwelling wildebeest. *Nature*, 563, 393–396.
- Darwin, C.** (1859). *On the Origin of Species by means of natural selection*. John Murray, London.
- Darwin, C. & Wallace, A.R.** (1858). On the Tendency of Species to form Varieties; and on the Perpetuation of Varieties and Species by Natural Means of Selection. *Journal of the Proceedings of the Linnean Society of London. Zoology*, 3, 45–62.
- de Jager, M., Weissing, F.J., Herman, P.M.J., Nolet, B.A. & Koppel, J. van de.** (2011). Lévy Walks Evolve Through Interaction Between Movement and Environmental Complexity. *Science*, 332, 1551–1553.
- DeAngelis, D.L.** (2018). *Individual-based models and approaches in ecology: Populations, communities and ecosystems*. CRC Press.
- DeAngelis, D.L. & Diaz, S.G.** (2019). Decision-Making in Agent-Based Modeling: A Current Review and Future Prospectus. *Frontiers in Ecology and Evolution*, 6.
- DeAngelis, D.L. & Mooij, W.M.** (2005). Individual-based modeling of ecological and evolutionary processes. *Annual Review of Ecology, Evolution, and Systematics*, 36, 147–168.
- Dhanjal-Adams, K.L., Bauer, S., Emmenegger, T., Hahn, S., Lisovski, S. & Liechti, F.** (2018). Spatiotemporal Group Dynamics in a Long-Distance Migratory Bird. *Current Biology*, 28, 2824–2830.e3.
- Dingemanse, N.J. & Réale, D.** (2005). Natural selection and animal personality. *Behaviour*, 142, 1159–1184.
- Duckworth, R.A. & Badyaev, A.V.** (2007). Coupling of dispersal and aggression facilitates the rapid range expansion of a passerine bird. *Proceedings of the National Academy of Sciences*, 104, 15017–15022.
- Duckworth, R.A. & Sockman, K.W.** (2012). Proximate mechanisms of behavioural inflexibility: Implications for the evolution of personality traits. *Functional Ecology*, 26, 559–566.
- Enquist, M. & Ghirlanda, S.** (2013). *Neural networks and animal behavior*. Princeton University Press.
- Etherington, T.R., Holland, E.P. & O'Sullivan, D.** (2015). NLMpy: A python software package for the creation of neutral landscape models within a general numerical framework. *Methods in Ecology and Evolution*, 6, 164–168.
- Fitzpatrick, J.W. & Senner, N.R.** (2018). Shorebirds, the World's Greatest Travelers, Face Extinction. *The New York Times*.
- Fretwell, S.D. & Lucas, H.L.** (1970). On territorial behavior and other factors influencing habitat distribution in birds. *Acta Biotheoretica*, 19, 16–36.
- Fryxell, J.M., Hazell, M., Börger, L., Dalziel, B.D., Haydon, D.T. & Morales, J.M. et al.** (2008). Multiple movement modes by large herbivores at multiple spatiotemporal scales. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 19114–9.
- Gershenfeld, N.A.** (1999). *The nature of mathematical modeling*. Cambridge University Press.
- Getz, W.M., Salter, R., Lyons, A.J. & Sippl-Swezey, N.** (2015). Panmictic and Clonal Evolution on a Single Patchy Resource Produces Polymorphic Foraging Guilds. *PLOS ONE*, 10, e0133732–e0133732.
- Getz, W.M., Salter, R., Seidel, D.P. & van Hooft, P.** (2016). Sympatric speciation in structureless environments. *BMC Evolutionary Biology*, 16, 50–50.
- Gill, R.E., Lee, T.T., David C., D., Handel, Colleen M., Mulcahy, Daniel M. & Gottschalck, Jon C. et al.** (2009). Extreme endurance flights by landbirds crossing the Pacific Ocean: Ecological corridor rather than barrier? *Proceedings of the*

*Royal Society B: Biological Sciences*, 276, 447–457.

- Goss-Custard, J.D.** (1980). Competition for food and interference among waders. *Ardea*, 55, 31–53.
- Grant, P.R. & Grant, B.R.** (2008). *How and why species multiply: The radiation of darwin's finches*. Princeton University Press.
- Grimm, V., Berger, U., Bastiansen, F., Eliassen, S., Ginot, V. & Giske, J. et al.** (2006). A standard protocol for describing individual-based and agent-based models. *Ecological Modelling*, 198, 115–126.
- Grimm, V., Berger, U., DeAngelis, D.L., Polhill, J.G., Giske, J. & Railsback, S.F.** (2010). The ODD protocol: A review and first update. *Ecological Modelling*, 221, 2760–2768.
- Groothuis, T.G.G. & Trillmich, F.** (2011). Unfolding personalities: The importance of studying ontogeny. *Developmental Psychobiology*, 53, 641–655.
- Guillemette, M., Himmelman, J.H. & Reed, A.** (1996). Availability and consumption of food by common eiders wintering in the Gulf of St. Lawrence: Evidence of prey depletion. *Canadian Journal of Zoology*, 74, 32–38.
- Guttal, V., Romanczuk, P., Simpson, S.J., Sword, G.A. & Couzin, I.D.** (2012). Cannibalism can drive the evolution of behavioural phase polyphenism in locusts. *Ecology Letters*, 15, 1158–1166.
- Hamilton, W.D.** (1971). Geometry for the selfish herd. *Journal of theoretical Biology*, 31, 295–311.
- Handegard, N.O., Boswell, K.M., Ioannou, C.C., Leblanc, S.P., Tjøstheim, D.B. & Couzin, I.D.** (2012). The Dynamics of Coordinated Group Hunting and Collective Information Transfer among Schooling Prey. *Current Biology*, 22, 1213–1217.
- Hildenbrandt, H., Carere, C. & Hemelrijk, C.** (2010). Self-organized aerial displays of thousands of starlings: A model. *Behavioral Ecology*, 21, 1349–1359.
- Hirota, M., Holmgren, M., Nes, E.H.V. & Scheffer, M.** (2011). Global Resilience of Tropical Forest and Savanna to Critical Transitions. *Science*, 334, 232–235.
- Holyoak, M., Casagrandi, R., Nathan, R., Revilla, E. & Spiegel, O.** (2008). Trends and missing parts in the study of movement ecology. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 19060–5.
- Huete, A., Didan, K., Miura, H., Rodriguez, E.P., Gao, X. & Ferreira, L.F.** (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens. Environ.*, 83, 195–213.
- Hussey, N.E., Kessel, S.T., Aarestrup, K., Cooke, S.J., Cowley, P.D. & Fisk, A.T. et al.** (2015). Aquatic animal telemetry: A panoramic window into the underwater world. *Science*, 348, 1255642–1255642.
- Huston, M., DeAngelis, D. & Post, W.** (1988). New Computer Models Unify Ecological Theory. *BioScience*, 38, 682–691.
- Ioannou, C.C., Guttal, V. & Couzin, I.D.** (2012). Predatory fish select for coordinated collective motion in virtual prey. *Science*, 337, 1212–1215.
- IPCC.** (2018). *Summary for Policymakers. In: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*. World Meteorological Organisation, Geneva.
- Jefferies, R.L., Jano, A.P. & Abraham, K.F.** (2006). A biotic agent promotes large-scale catastrophic change in the coastal marshes of Hudson Bay. *Journal of Ecology*, 94, 234–242.
- Joo, R., Boone, M.E., Clay, T.A., Patrick, S.C., Clusella-Trullas, S. & Basille, M.** (2019). Navigating through the R packages for movement. *arXiv:1901.05935 [q-bio, stat]*.
- Kays, R., Crofoot, M.C., Jetz, W. & Wikelski, M.** (2015). Terrestrial animal tracking as an eye on life and planet. *Science*, 348, aaa2478.
- Keddy, P.A.** (2001). *Competition*. Population and Community Biology Series. 2nd edn. Springer Netherlands.
- Kery, M. & Royle, J.A.** (2019). *Applied hierarchical modeling in ecology: Analysis of distribution, abundance and species richness in R and BUGS: Dynamic and advanced models*. Academic Press.
- Klaassen, R.H.G., Nolet, B.A. & Bankert, D.** (2006). Movement of Foraging Tundra Swans Explained by Spatial Pattern in Cryptic Food Densities. *Ecology*, 87, 2244–2254.
- Kraan, C., Gils, J.A.V., Spaans, B., Dekinga, A., Bijleveld, A.I. & Roomen, M.V. et al.** (2009a). Landscape-scale experiment demonstrates that Wadden Sea intertidal flats are used to capacity by molluscivore migrant shorebirds. *Journal of*

*Animal Ecology*, 78, 1259–1268.

- Kraan, C., Meer, J. van der, Dekinga, A. & Piersma, T.** (2009b). Patchiness of macrobenthic invertebrates in homogenized intertidal habitats: Hidden spatial structure at a landscape scale. *Marine Ecology Progress Series*, 383, 211–224.
- Krause, J. & Ruxton, G.D.** (2002). *Living in groups*. Oxford University Press.
- Lank, D.B., Butler, R.W., Ireland, J. & Ydenberg, R.C.** (2003). Effects of predation danger on migration strategies of sandpipers. *Oikos*, 103, 303–319.
- Laundré, J.W., Hernández, L. & Altendorf, K.B.** (2001). Wolves, elk, and bison: Reestablishing the “landscape of fear” in Yellowstone National Park, U.S.A. *Canadian Journal of Zoology*, 79, 1401–1409.
- le Roux, E., Kerley, G.I.H. & Cromsigt, J.P.G.M.** (2018). Megaherbivores Modify Trophic Cascades Triggered by Fear of Predation in an African Savanna Ecosystem. *Current Biology*, 28, 2493–2499.e3.
- Leclerc, M., Vander Wal, E., Zedrosser, A., Swenson, J.E., Kindberg, J. & Pelletier, F.** (2016). Quantifying consistent individual differences in habitat selection. *Oecologia*, 180, 697–705.
- Legendre, P.** (1993). Spatial Autocorrelation: Trouble or New Paradigm? *Ecology*, 74, 1659–1673.
- Lek, S. & Guégan, J.F.** (1999). Artificial neural networks as a tool in ecological modelling, an introduction. *Ecological Modelling*, 120, 65–73.
- Levin, S.A.** (1992). The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology*, 73, 1943–1967.
- Lima, S.L. & Zollner, P.A.** (1996). Anti-predatory vigilance and the limits to collective detection: Visual and spatial separation between foragers. *Behavioral Ecology and Sociobiology*, 38, 355–363.
- MacArthur, R.H. & Wilson, E.O.** (2001). *The Theory of Island Biogeography*. Princeton University Press.
- MacCurdy, R., Bijleveld, A., Gabrielson, R., Cluderay, J., Spaulding, E. & Oudman, T. et al.** (2015). Automatic, intensive wildlife radiotracking. In: *Untying the knot*. p. 17.
- Mathot, K.J., Dekinga, A. & Piersma, T.** (2017). An experimental test of stateBehaviour feedbacks: Gizzard mass and foraging behaviour in red knots. *Functional Ecology*, 31, 1111–1121.
- Maynard Smith, J.** (1982). *Evolution and the theory of games*. Cambridge University Press.
- Myers, J.P.** (1983). Space, time and the pattern of individual associations in a group-living species: Sanderlings have no friends. *Behavioral Ecology and Sociobiology*, 12, 129–134.
- Naef-Daenzer, B., Widmer, F. & Nuber, M.** (2001). A test for effects of radio-tagging on survival and movements of small birds. *Avian Science*, 1, 15–23.
- Nathan, R.** (2008). An emerging movement ecology paradigm. *Proc. Natl. Acad. Sci.*, 105, 19050–19051.
- Nathan, R., Getz, W.M., Revilla, E., Holyoak, M., Kadmon, R. & Saltz, D. et al.** (2008). A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*, 105, 19052–19059.
- Netz, C.** (2017). A neural-network model for predator-prey coevolution. Master. Faculty of Science; Engineering, Groningen.
- Nolet, B.A. & Mooij, W.M.** (2002). Search paths of swans foraging on spatially autocorrelated tubers. *Journal of Animal Ecology*, 71, 451–462.
- Nolet, B.A., Klaassen, R.H.G. & Mooij, W.M.** (2006). The use of a flexible patch leaving rule under exploitative competition: A field test with swans. *Oikos*, 112, 342–352.
- Oudman, T., Bijleveld, A.I., Kavelaars, M.M., Dekinga, A., Cluderay, J. & Piersma, T. et al.** (2016). Diet preferences as the cause of individual differences rather than the consequence. *Journal of Animal Ecology*, 85, 1378–1388.
- Oudman, T., Piersma, T., Ahmedou Salem, M.V., Feis, M.E., Dekinga, A. & Holthuijsen, S. et al.** (2018). Resource landscapes explain contrasting patterns of aggregation and site fidelity by red knots at two wintering sites. *Movement Ecology*, 6, 24–24.
- Piersma, T.** (2011). Flyway evolution is too fast to be explained by the modern synthesis: Proposals for an “extended” evolutionary research agenda. *Journal of Ornithology*, 152, 151–159.
- Piersma, T. & Bonan, A.** (2019). Sandpipers, Snipes, Phalaropes (Scolopacidae). In: *Handbook of the Birds of the World Alive*.



Lynx Edicions, Barcelona.

- Piersma, T. & Lindström, Å.** (2004). Migrating shorebirds as integrative sentinels of global environmental change. *Ibis*, 146, 61–69.
- Piersma, T., van Aelst, R., Kurk, K., Berkhoudt, H. & Maas, L.R.M.** (1998). A new pressure sensory mechanism for prey detection in birds: The use of principles of seabed dynamics? *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 265, 1377–1383.
- Pruitt, J.N. & Riechert, S.E.** (2011). How within-group behavioural variation and task efficiency enhance fitness in a social group. *Proceedings of the Royal Society B: Biological Sciences*, 278, 1209–1215.
- Réale, D., Reader, S.M., Sol, D., McDougall, P.T. & Dingemanse, N.J.** (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews*, 82, 291–318.
- Ruthrauff, D.R., Dekinga, A., Gill, R.E. & Piersma, T.** (2018). Energetic solutions of Rock Sandpipers to harsh winter conditions rely on prey quality. *Ibis*, 160, 397–412.
- Ruthrauff, D.R., Gill, R.E. & Tibbitts, T.L.** (2013). Coping with the Cold: An Ecological Context for the Abundance and Distribution of Rock Sandpipers during Winter in Upper Cook Inlet, Alaska. *Arctic*, 66, 269–278.
- Scantlebury, D.M., Mills, M.G.L., Wilson, R.P., Wilson, J.W., Mills, M.E.J. & Durant, S.M. et al.** (2014). Flexible energetics of cheetah hunting strategies provide resistance against kleptoparasitism. *Science*, 346, 79–81.
- Scheffer, M.** (2009). *Critical Transitions in Nature and Society*. Princeton University Press.
- Sciaini, M., Fritsch, M., Scherer, C. & Simpkins, C.E.** (2018). NLMR and landscapetools: An integrated environment for simulating and modifying neutral landscape models in R. *Methods in Ecology and Evolution*, 0.
- Sih, A. & Bell, A.M.** (2008). Insights for behavioral ecology from behavioral syndromes. In: *Advances in the Study of Behavior*. Academic Press, pp. 227–281.
- Sih, A., Bell, A. & Johnson, J.** (2004a). Behavioral syndromes: An ecological and evolutionary overview. *Trends in Ecology & Evolution*, 19, 372–378.
- Sih, A., Bell, A.M., Johnson, J.C. & Ziemba, R.E.** (2004b). Behavioral Syndromes: An Integrative Overview. *The Quarterly Review of Biology*, 79, 241–277.
- Sih, A., Cote, J., Evans, M., Fogarty, S. & Pruitt, J.** (2012). Ecological implications of behavioural syndromes. *Ecology Letters*, 15, 278–289.
- Spiegel, O., Getz, W.M. & Nathan, R.** (2013). Factors influencing foraging search efficiency: Why do scarce lappet-faced vultures outperform ubiquitous white-backed vultures? *The American naturalist*, 181, E102–15.
- Spiegel, O., Leu, S.T., Bull, C.M. & Sih, A.** (2017). What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecology Letters*, 20, 3–18.
- Staver, A.C., Archibald, S. & Levin, S.A.** (2011). The Global Extent and Determinants of Savanna and Forest as Alternative Biome States. *Science*, 334, 230–232.
- Strandburg-Peshkin, A., Farine, D.R., Couzin, I.D. & Crofoot, M.C.** (2015). Shared decision-making drives collective movement in wild baboons. *Science*, 348, 1358–1361.
- Sutherland, W.J.** (1996). *From individual behaviour to population ecology*. Oxford University Press.
- Székely, T. & Bamberger, Z.** (1992). Predation of Waders (Charadrii) on Prey Populations: An Exclosure Experiment. *Journal of Animal Ecology*, 61, 447–456.
- Tunstrøm, K., Katz, Y., Ioannou, C.C., Huepe, C., Lutz, M.J. & Couzin, I.D.** (2013). Collective states, multistability and transitional behavior in schooling fish. *PLoS Comput Biol*, 9, e1002915–e1002915.
- Turner, M.G. & Gardner, R.H.** (2015). Introduction to Models. In: *Landscape Ecology in Theory and Practice: Pattern and Process* (eds. Turner, M.G. & Gardner, R.H.). Springer New York, New York, NY, pp. 63–95.
- Vahl, W.K.** (2006). *Interference competition among foraging waders*. s.n.
- Vahl, W.K. & Kingma, S.A.** (2007). Food divisibility and interference competition among captive ruddy turnstones, *Arremonia interpres*. *Animal Behaviour*, 74, 1391–1401.
- Vahl, W.K., Lok, T., van der Meer, J., Piersma, T. & Weissing, F.J.** (2005). Spatial clumping of food and social dominance

- affect interference competition among ruddy turnstones. *Behavioral Ecology*, 16, 834–844.
- van de Koppel, J., Rietkerk, M. & Weissing, F.J.** (1997). Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology & Evolution*, 12, 352–356.
- van den Berg, P. & Weissing, F.J.** (2015). The importance of mechanisms for the evolution of cooperation. *Proceedings of the Royal Society B: Biological Sciences*, 282, 20151382.
- van Gils, J.A.** (2010). State-dependent Bayesian foraging on spatially autocorrelated food distributions. *Oikos*, 119, 237–244.
- van Gils, J.A. & Piersma, T.** (2004). Digestively constrained predators evade the cost of interference competition. *Journal of Animal Ecology*, 73, 386–398.
- van Gils, J.A., Schenk, I.W., Bos, O. & Piersma, T.** (2003). Incompletely Informed Shorebirds That Face a Digestive Constraint Maximize Net Energy Gain When Exploiting Patches. *The American Naturalist*, 161, 777–793.
- van Gils, J.A., Spaans, B., Dekinga, A. & Piersma, T.** (2006). Foraging in a tidally structured environment by red knots (*Calidris Canutus*): Ideal, but not free. *Ecology*, 87, 1189–1202.
- Venkateswaran, V., Shrivastava, A., Kumble, A.L.K. & Borges, R.M.** (2017). Life-history strategy, resource dispersion and phylogenetic associations shape dispersal of a fig wasp community. *Movement Ecology*, 5, 25–25.
- Walsh, P.D.** (1996). Area-restricted Search and the Scale Dependence of Path Quality Discrimination. *Journal of Theoretical Biology*, 183, 351–361.
- Watson, A. & Miller, G.R.** (1971). Territory Size and Aggression in a Fluctuating Red Grouse Population. *Journal of Animal Ecology*, 40, 367–383.
- Wikelski, M. & Tertitski, G.** (2016). Living sentinels for climate change effects. *Science*, 352, 775–776.
- Williams, T.M., Fuiman, L.A., Horning, M. & Davis, R.W.** (2004). The cost of foraging by a marine predator, the Weddell seal *Leptonychotes weddellii*: Pricing by the stroke. *Journal of Experimental Biology*, 207, 973–982.
- Wilson, D.S.** (1998). Adaptive individual differences within single populations. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 353, 199–205.
- Wolf, M. & McNamara, J.M.** (2012). On the evolution of personalities via frequency-dependent selection. *The American Naturalist*, 179, 679–692.
- Wolf, M. & Weissing, F.J.** (2010). An explanatory framework for adaptive personality differences. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365, 3959–3968.
- Wolf, M. & Weissing, F.J.** (2012). Animal personalities: Consequences for ecology and evolution. *Trends in Ecology & Evolution*, 27, 452–461.
- Wolf, M., van Doorn, G.S., Leimar, O. & Weissing, F.J.** (2007a). Life-history trade-offs favour the evolution of animal personalities. *Nature*, 447, 581–584.
- Wolf, M., van Doorn, G.S., Leimar, O. & Weissing, F.J.** (2007b). *Wolf et al. Reply. Nature*, 450, E5–E6.
- Ydenberg, R.C., Butler, R.W., Lank, D.B., Smith, B.D. & Ireland, J.** (2004). Western sandpipers have altered migration tactics as peregrine falcon populations have recovered. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271, 1263–1269.