



BITS Pilani
Pilani | Dubai | Goa | Hyderabad

Introduction to Statistical Methods(ISM) (AIMLC ZC418)

Dr YVK Ravi Kumar
yvk.ravikumar@pilani.bits-pilani.ac.in

➤ **Lead Faculty:** Dr. G Krishna Kumari(gkrishnakumari@wilp.bits-pilani.ac.in)



Session 1 **(25th April 2026)**

Overview of the course & Basic of Statistics


Overview of the course

- **Module 1** : Basic Probability & Statistics
- **Module 2** : Conditional Probability & Bayes' theorem
- **Module 3** : Probability Distributions
- **Module 4** : Hypothesis Testing
- **Module 5** : Prediction & Forecasting
- **Module 6** : Prediction & Forecasting; Gaussian Mixture model & Expectation Maximization



TEXT BOOKS

- T1 : Statistics for Data Scientists, An introduction to Probability, Statistics and Data Analysis, Maurits Kaptein et al, Springer 2022
- T2 : Probability and Statistics for Engineering and Sciences, 8th Edition, Jay L Devore, Cengage Learning
- T3 : Introduction to Time Series and Forecasting, Second Edition , Peter J Brockwell, Richard A Davis, Springer.



Evaluation Components

	Name	Type	Weightage
EC 1	Quiz 1 & Quiz 2	Online	10%
	Assignment 1 Situating Learning	Online Online	10% 10%
EC 2	Mid semester Exam	Closed Book	30%
EC 3	Comprehensive Exam	Open Book	40%


Module 1: (Basic Probability & Statistics)

Contact Session	List of Topic Title	Reference
CS - 1	Measures of Central Tendency & Measures of Variability, Data – Symmetric & Asymmetric, outlier detection, 5 point summary	T1 & T2



Statistics

- Statistics is a branch of Applied Mathematics
- Statistics is the back-bone of Data Science, AI & ML and all Information Technologies




Statistics


➤ **Statistics may be defined as science that is employed to**

- Collect the data
- Present and organize the data in a systematic manner
- Analyse the data
- Infer about the data
- Take decision from the data.

Statistics is a Data(Information) driven decision making Science.



Data - Types



Levels of Data Measurement

- Nominal — Lowest level of measurement
- Ordinal
- Interval
- Ratio — Highest level of measurement

❑ **Nominal** ○ A **nominal scale** classifies data into distinct categories in which no ranking is implied
○ **Example** : Gender, Marital Status, voting to different party

❑ **Ordinal scale** ○ An **ordinal scale** classifies data into distinct categories in which ranking is implied
○ **Example**:
1. Product satisfaction: Satisfied, Neutral, Unsatisfied
2. Faculty rank: Professor, Associate Professor, Assistant Professor
3. Student Grades: A, B, C, D, F
4. Medals won: Gold, Silver, Bronze

❑ **Interval scale** ○ An **interval scale** is an ordered scale in which the difference between measurements is a meaningful quantity but the measurements do not have a true zero point.
○ **Example**
1. Temperature in Fahrenheit and Celsius
2. Months of the Year: there's no month called zero and we can't say January is twice as much as June.

❑ **Ratio Scale** ○ A **ratio scale** is an ordered scale in which the difference between the measurements is a meaningful quantity and the measurements have a true zero point.
○ **Examples**
Weight , Age , Salary

Variables

Types of Variable

Qualitative (Categorical): express a qualitative attribute such as hair color, eye color, religion.

Quantitative(Numerical): measured in terms of numbers such as height, weight, number of people.

Nominal:
No ordering is possible such as hair color, eye color, religion.

Ordinal:
Ordering is possible such as health, which can take values such as poor, reasonable, good, or excellent.

Discrete: countable and have a **finite number of possibilities** such as number of people

Continuous: not countable and have an **infinite number of possibilities** such as height

INTERVAL: ratio of values of variable do not have any meaning and it does not have an inherently defined zero value such as temperature

RATIO: ratio of values of variable have meaning and it have an inherently defined zero value such as length

Example

Question	Type of Variable
Types of vehicle owned Two wheeler, four wheeler	Categorical : Nominal
Product satisfaction Unsatisfied, neutral, fairly satisfied, satisfied	Categorical : Ordinal
To how many magazines do you currently subscribed Zero, One, Two, Three, Four	Discrete
How tall are you (in inches)	Continuous: Ratio
Weight (in Kilograms)	Continuous: Ratio
Temperature (in degrees Celsius or degrees Fahrenheit)	Continuous: Interval

Measures of Central Tendency

- Measure of central tendency provides a very convenient way of representing (reducing/summarizing) a data set with a single number that describes the **significance** of the whole data set.
- Also defined as a single value that is used to describe the “**center**” of the data.
- Three commonly used measures of central tendency:
 1. Mean
 2. Median
 3. Mode

Mean

- Also referred as the “**arithmetic average**”
- The most commonly used measure of the center of data
- Numbers that describe what is average or typical of the distribution
- Computation of Sample Mean:

$$\bar{Y} = \frac{\sum Y}{n}$$

 -
- Computation of the Mean for grouped Data
- $\bar{Y} = \frac{\sum f Y}{N}$ where $f Y$ = a score multiplied by its frequency

Mean: Ungrouped Data

- Suppose you define the time to get ready as the time (rounded to the nearest minute) from when you get out of bed to when you leave your home. You collect the times shown below for 10 consecutive work days:

Day	1	2	3	4	5	6	7	8	9	10
Time (min)	39	29	43	52	39	44	40	31	44	35

$$\bar{X} = \frac{\text{Sum of the values}}{\text{Number of values}}$$

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$$

$$\bar{X} = \frac{39 + 29 + 43 + 52 + 39 + 44 + 40 + 31 + 44 + 35}{10}$$


$$\bar{X} = \frac{396}{10} = 39.6$$

Mean: Frequency distribution

Data of Children watching TV in Bengaluru

Hours Spent Watching TV	Frequency (f)	fY
1	104	104
2	130	260
3	98	294
Total	332	658

$$\bar{Y} = \frac{\sum fY}{N} = \frac{658}{332} = 1.98$$



Mean

➤ PROPERTIES

- Mean is the most stable among other measures of central tendency because every score contributes to the value of the mean.
- It may easily be affected by the extreme scores.
- The sum of each score's distance from the mean is zero.
- It may not be an actual score in the distribution
- It is very easy to compute.

➤ When to Use the Mean

- Data is quantitative in nature without extreme/outlier observations.
- It can be applied to interval and ratio levels of measurements
- Sampling stability is desired.
- Other measures are to be computed such as standard deviation, coefficient of variation and skewness

The Mode

- The category or score with the largest frequency (or percentage) in the distribution.
- The mode can be calculated for variables with levels of measurement that are: nominal, ordinal, or interval-ratio.

Example:

- A systems manager in charge of a company's network keeps track of the number of server failures that occur in a day. Compute the mode for the following data, which represents the number of server failures in a day for the past two weeks:.

1 3 0 3 26 2 7 4 2 3 3 6 3

- Because 3 appears five times, more times than any other value, the mode is 3. Thus, the systems manager can say that the most common occurrence is having three server failures in a day.


Mode

Properties

- It can be used when the data are qualitative as well as quantitative.
- It may not be unique.
- It is not affected by extreme values.

When to Use the Mode

- When the data set is measured on a nominal scale
- A decision is desirably taken on the basis maximum frequent occurrences.



The Median

- The score that **divides the distribution into two equal parts**, so that half the cases are above it and half below it.
- You compute the median value by following one of two rules:
 - **Rule 1** If there are an *odd* number of values in the data set, the median is the middle-ranked value.
 - **Rule 2** If there are an *even* number of values in the data set, then the median is the *average* of the two middle ranked values.

Median

Properties

- The median is the **middle positional score**, or average of middle scores in a distribution.
 - Fifty percent (50%) lies below the median value and 50% lies above the median value.
 - It is also known as the middle score or the 50th percentile/middle quartile.
 - Median is never affected by extreme observations/Outliers

Uses

- Even though it is applicable for all types of measurement scales, it is more useful for ordinal measurement scales.
- It is a best measure where the influences of extreme observations/Outliers are not required.

The Median

- **Example-1:** The three-year annualized returns for the seven small-cap growth funds with low risk are ranked from the smallest to the largest:

The Median is the 4th position value i.e. 22.4.

- **Example-2:** Consider the time taken for reaching office from home in minutes arranged from lowest to highest

29 31 35 39 39 40 43 44 44 52

- The number of observations are 10. Hence the median is the average of values at the 5th and 6th Position, i.e. average of 39 and 40 = **39.5**

Data distribution-Skew (Asymmetry) & Symmetry

➤ Skewed

○ Negatively (Left): $\text{mean} < \text{median} < \text{Mode}$

- Distributions with a left skew have long left tails;


○ Positively (Right) : $\text{mean} > \text{median} > \text{Mode}$

- Distributions with a right skew have long right tails.

➤ Symmetry: $\text{Mean} = \text{Median} = \text{Mode}$

➤ Bimodal : has two distinct modes

➤ Multi-modal : has more than 2 distinct modes



Note: In science, an empirical relationship is a relationship that is supported by experiment and observation but not necessarily supported by theory. For moderately skewed/unskew data, an empirical relationship is observed as $\text{Mode} = 3\text{Median} - 2\text{Mean}$. Further Mean is applied for unskew/moderately skewed data whereas median is a preferred measure for heavily skewed data.

Why mean(measure of central tendency) is not enough?

Sl. No.	X_1
1	2
2	8
3	5
4	3
5	7
6	8
7	5
8	2
9	5
Total	45

Statistical measures	Group 1
Mean	5
Median	5
Mode	5

Sl. No.	X_2
1	1
2	15
3	5
4	5
5	6
6	3
7	5
8	2
9	3
Total	45

Statistical measures	Group 2
Mean	5
Median	5
Mode	5

Sl. No.	X_1
1	2
2	8
3	5
4	3
5	7
6	8
7	5
8	2
9	5
Total	45

Statistical measures	Group 1 & 2
Mean	5
Median	5
Mode	5

Do we need any other measure?

Answer: Yes

Measures of variability/dispersion/spread



➤ Measures of Variability:

- The Range
- The Variance
- The Standard Deviations
- Quartile deviation



Measure of Variability

- Variability can be defined several ways:
 - A quantitative distance measure based on the differences between scores
 - Describes distance of the spread of scores or distance of a score from the mean
- Purposes of Measure of Variability:
 - Describe the distribution
 - Measure how well a measure of central tendency represents the distribution



The Ranges

- The distance covered by the scores in a distribution – From smallest value to highest value
- For continuous data, real limits are used

$$\text{Range} = \text{URL for } X_{\max} - \text{LRL for } X_{\min}$$

- Based on two scores, not all the data – An imprecise, unreliable measure of variability


Example: For a set of scores: 7, 2, 7, 6, 5, 6, 2

$$\text{Range} = \text{Highest Score minus Lowest score} = 7 - 2 = 5$$

The Standard Deviation

- Most common and most important measure of variability is the standard deviation
 - A measure of the standard, or average, distance from the mean
 - Describes whether the scores are clustered closely around the mean or are widely scattered
- Calculation differs for population and samples
- Variance is a necessary *companion concept* to standard deviation but *not the same* concept

The Standard Deviation



Exercise : Find out the deviations of all the data points with the mean....and then find the 'mean deviation'.

The Standard Deviation

- Sum of mean deviations is always 'zero' !
(because Mean is a balance point)
- Then, how do you find 'Standard Deviation' ?



Need a new strategy

The Standard Deviation

➤ New Strategy :

- a) First square each deviation score
- b) Then sum the Squared Deviations (SS)
- c) Average the squared deviations

- Mean Squared Deviation is known as “**Variance**”
- Variability is now measured in squared units

$$\textit{Standard Deviation} = \sqrt{\textit{Variance}}$$

The Variance

Variance equals mean (average) squared deviation (distance) of the scores from the mean

$$\text{Variance} = \frac{\text{sum of squared deviations}}{\text{number of scores}}$$

where $SS = \sum (X - \mu)^2$

The Population Variance

- ❖ Population variance equals mean (average) squared deviation (distance) of the scores from the population mean
- ❖ Variance is the average of squared deviations, so we identify population variance with a lowercase Greek letter sigma squared: σ^2
- ❖ Standard deviation is the square root of the variance, so we identify it with a lowercase Greek letter sigma: σ

Sl. No.	X ₁
1	2
2	8
3	5
4	3
5	7
6	8
7	5
8	2
9	5
Total	45

$$\bar{X} = \frac{\sum_{i=1}^9 x_i}{N} = \frac{45}{9} = 5$$

$$S = \sqrt{\frac{\sum (x - \bar{X})^2}{n - 1}}$$

$$S = \sqrt{\frac{44}{8}} = 2.345$$

Sl. No.	X ₂
1	1
2	15
3	5
4	5
5	6
6	3
7	5
8	2
9	3
Total	45

$$\bar{X} = \frac{\sum_{i=1}^9 x_i}{N} = \frac{45}{9} = 5$$

$$S = \sqrt{\frac{\sum (x - \bar{X})^2}{n - 1}}$$

$$S = \sqrt{\frac{134}{8}} = 4.093$$

Standard Deviation and Variance for a Sample

➤ Goal of inferential statistics:

- Draw general conclusions about population based on limited information from a sample

➤ Samples differ from the population


- Samples have less variability
- Computing the Variance and Standard Deviation in the same way as for a population would give a biased estimate of the population values

Sample Standard Deviation and Variance

- Sum of Squares (SS) is computed as before
- Formula for Variance has $n-1$ rather than N in the denominator
- Notation uses s instead of σ

$$\text{variance of sample} = s^2 = \frac{SS}{n-1}$$

$$\text{standard deviation of sample} = s = \sqrt{\frac{SS}{n-1}}$$



Learning Check

Select the correct option

- a) A sample of four scores has $SS = 24$. What is the variance?
- (1) The variance is 6
 - (2) The variance is 7
 - (3) The variance is 8
 - (4) The variance is 12
- b) A sample systematically has less variability than a population
- b) The standard deviation is the distance from the Mean to the farthest point on the distribution curve

True / False ?

True / False ?

Learning Check

Select the correct option



- a) A sample of four scores has $SS = 24$. What is the variance?
- (1) The variance is 6
 - (2) The variance is 7
 - (3) The variance is 8**
 - (4) The variance is 12
- b) A sample systematically has less variability than a population
- b) The standard deviation is the distance from the Mean to the farthest point on the distribution curve

True

False

Descriptive Statistics


- A standard deviation describes scores in terms of distance from the mean
- Describe an entire distribution with just two numbers (M and s)
- Reference to both allows reconstruction of the measurement scale from just these two numbers
- Means and standard deviations together provide extremely useful descriptive statistics for characterizing distributions



Five point summary of Data

The five number summary of data includes 5 items:

- **Minimum.**
- **Q1** (the first quartile, or the 25% mark).
- **Median.**
- **Q3** (the third quartile, or the 75% mark).
- **Maximum.**



Interquartile range (IQR) & Quartile Deviation (QD)

- It is measure of Variation
- Also Known as Mid-spread : Spread in the Middle 50%
- Difference Between Third & First Quartiles:
- Not Affected by Extreme Values
- Interquartile Range = $IQR = Q_3 - Q_1$

Data in Ordered Array: 11 12 13 16 16 17 17 18 21


Position of $Q_1 = \frac{1 \cdot (9 + 1)}{4} = 2.50$, $Q_1 = 12.5$

Position of $Q_3 = \frac{3 \cdot (9 + 1)}{4} = 7.50$, $Q_3 = 17.5$


Interquartile Range = $IQR = Q_3 - Q_1 = 17.5 - 12.5 = 5$

$QD = IQR/2 = 5/2 = 2.5$ (variation among the data in the central 50% of a data set)

Box and Whisker plot




Box and Whisker plot



Detection of outliers

- Data in Ordered Array of the previous data: 11 12 13 16 16 17 17 18
- $Q_1 - 1.5 \text{ IQR} = 5$, 11 12 13 16 16 17 17 18, $Q_3 + 1.5 \text{ IQR} = 25$
- As all the above data points lie in between $Q_1 - 1.5 \text{ IQR}$ and $Q_3 + 1.5 \text{ IQR}$, the above data is free from Outliers/Extreme Values



Potential outliers



- ❖ The lower limit and upper limit of a data set are given by:

$$\text{Lower limit} = Q_1 - 1.5 \times \text{IQR}$$

$$\text{Upper limit} = Q_3 + 1.5 \times \text{IQR}$$

- ❖ Data points that lie below the lower limit or above the upper limit are **potential outliers**.

Outlier Detection



Statistical Summary

Cost	Weight	Weight1	Length	Height	Width	Width 1
count	159.000000	159.000000	159.000000	159.000000	159.000000	159.000000
mean	398.326415	26.247170	28.415723	31.227044	8.970994	4.417486
std	357.978317	9.996441	10.716328	11.610246	4.286208	1.685804
min	0.000000	7.500000	8.400000	8.800000	1.728400	1.047600
25%	120.000000	19.050000	21.000000	23.150000	5.944800	3.385650
50%	273.000000	25.200000	27.300000	29.400000	7.786000	4.248500
75%	650.000000	32.700000	35.500000	39.650000	12.365900	5.584500
max	1650.000000	59.000000	63.400000	68.000000	18.957000	8.142000

Summary

1. **Measures of Central Tendency:** Mean, Median, Mode
2. **Measures of Variability:** Range, Standard Deviation , Variance
3. Symmetric and Asymmetric distribution
4. Five point summary
5. Outliers



Practice Problem:


- Q.1 A sample of 77 individuals working at a particular office was selected and the noise level (dBA) experienced by everyone is the following data:

55.3, 55.3, 55.3, 55.9, 55.9, 55.9, 55.9, 56.1, 56.1, 56.1, 56.1, 56.1, 56.1,
56.8, 56.8, 57.0, 57.0, 57.0, 57.8, 57.8, 57.8, 57.9, 57.9, 57.9, 58.8, 58.8,
58.8, 59.8, 59.8, 59.8, 62.2, 62.2, 63.8, 63.8, 63.8, 63.9, 63.9, 63.9, 64.7,
64.7, 64.7, 65.1, 65.1, 65.1, 65.3, 65.3, 65.3, 65.3, 67.4, 67.4, 67.4, 67.4,
68.7, 68.7, 68.7, 68.7, 69.0, 70.4, 70.4, 71.2, 71.2, 71.2, 73.0, 73.0, 73.1,
73.1, 74.6, 74.6, 74.6, 74.6, 79.3, 79.3, 79.3, 79.3, 83.0, 83.0, 83.0.

- Find a) Arithmetic Mean, SD, variance, and IQR
b) Draw box and whisker plot
c) Comment on the outliers, if any.

Discussion

- Arithmetic Mean: 64.89 dBA
- Standard Deviation (SD): 7.80 dBA
- Variance: 60.88 (dBA)^2
- Interquartile Range (IQR): 12.60 dBA



- Lower Bound: $Q1 - 1.5 \times IQR = 59.8 - (1.5 \times 12.6) = 40.9$
- Upper Bound: $Q3 + 1.5 \times IQR = 72.4 + (1.5 \times 12.6) = 91.3$

No data points fall below 40.9, but values such as 79.3 and 83.0 are potential upper outliers, as they exceed 91.3.

Practice Problems:

Q.2 The data given below is the total fat, in grams per serving, for a sample of 20 chicken sandwiches from fast-food chains.

7 8 4 5 16 20 20 24 19 30 23 30 25 19 29 29 30 30 40 56

- Compute the mean, median, first quartile, and third quartile.
- Compute the variance, standard deviation, range, interquartile range, Are there any outliers? Explain.
- Are the data skewed? If so, how?
- Based on the results of (a) through (c), what conclusions can you reach concerning the total fat of chicken sandwiches?


Practice Problem:

Q.3 The following data represent the battery life (in shots) for three pixel digital cameras:

300	180	85	170	380	460	260	35	380	120
110	240								

➤ List the Five-point summary.

Q.4 For the data set below:



82	45	64	80	82	74	79	80	80	78	80	80	48	73	80	79	81	70	78	73
----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----

- Obtain and interpret the quartiles.
- Determine and interpret the interquartile range.
- Find and interpret the five-number(point) summary.
- Identify potential outliers, if any.
- Construct and interpret a boxplot.

Practice Problem:

Q.5 A bank branch located in a commercial place of a city has developed an improved process for serving customers during the noon-to-1:00 p.m. lunch period. The waiting time, in minutes (defined as the time the customer enters the line to when he or she reaches the teller window), of a sample of 15 customers during this hour is recorded over a period of one week. The results are: 4.21, 5.55, 3.02, 5.13, 4.77, 2.34, 3.54, 3.20, 4.50, 6.10, 0.38, 5.12, 6.46, 6.19, 3.79.

Another branch, located in a residential area, is also concerned with the noon-to-1 p.m. lunch hour. The waiting time, in of a sample of 15 customers during this hour is recorded over a period of one week. The results are listed below: 9.66, 5.90, 8.02, 5.79, 8.73, 3.82, 8.01, 8.35, 10.49, 6.68, 5.64, 4.08, 6.17, 9.91, 5.47.

- a. List the five-number summaries of the waiting times at the two bank branches.
- b. Construct box-and-whisker plots and describe the shape of the distribution of each for the two bank branches.
- c. What similarities and differences are there in the distributions of the waiting time at the two bank branches?



Thank You!