

Fever is a temporary increase in body temperature, often caused by the immune system responding to an infection such as a cold, flu, or other illness. Common symptoms can include feeling hot or chilled, sweating, tiredness, headache, and muscle aches. Fever itself is usually a sign that the body is responding to something rather than a disease on its own. Drinking enough fluids, getting adequate rest, and monitoring symptoms can help while the underlying illness improves.

If a fever is very high, lasts for several days, keeps returning, or occurs with concerning symptoms such as difficulty breathing, severe headache, confusion, stiff neck, or significant dehydration, medical attention should be sought. For children and teenagers, the appropriate response can depend on age and other symptoms, so it is best to involve a parent/guardian and a healthcare professional when you're concerned.

The common cold is a mild infection of the upper respiratory tract, usually caused by viruses. Common symptoms include a runny or blocked nose, sneezing, sore throat, cough, mild headache, and tiredness. Fever may occur but is generally mild. Most common colds improve on their own within several days to about a week.

Management usually focuses on rest, adequate fluids, and relieving symptoms. Warm fluids and adequate sleep may help with comfort. Antibiotics generally do not help because the common cold is usually caused by viruses. If symptoms become severe, last unusually long, or cause breathing difficulties, a healthcare professional should be consulted.

Medicine information for dataset: Symptom-relief medicines may sometimes be used under appropriate guidance; antibiotics are not normally useful for an uncomplicated viral cold.

Suggestions: Get adequate rest, drink fluids, wash hands regularly, and avoid close contact with others when sick.

Influenza, commonly called the flu, is a contagious respiratory infection caused by influenza viruses. Symptoms can appear suddenly and may include fever, cough, sore throat, headache, muscle aches, chills, and significant tiredness. Compared with an ordinary cold, influenza can cause more intense body aches and fatigue.

Treatment depends on the person's age, health status, and severity of illness. Rest and adequate fluid intake are important. In some situations, healthcare professionals may prescribe antiviral medicines, particularly for people at higher risk of complications or when treatment is started early. Medical evaluation is important if symptoms are severe.

Medicine information for dataset: Antiviral treatment may be prescribed by a healthcare professional when appropriate; do not self-select medication.

Suggestions: Rest, maintain hydration, avoid spreading the infection to others, and seek medical advice if symptoms become severe.

COVID-19 is an infectious disease caused by the SARS-CoV-2 virus. Symptoms can include fever, cough, sore throat, tiredness, headache, muscle aches, congestion, and sometimes changes in taste or smell. Symptoms vary considerably between people, and some individuals may have mild symptoms while others can become seriously ill.

Management depends on the severity of the infection and the individual's risk factors. Mild illness may be managed with rest, fluids, and monitoring. People with significant breathing problems, worsening symptoms, or other concerning symptoms should receive medical attention. Healthcare professionals may recommend specific treatments for people who are at higher risk.

Medicine information for dataset: Treatment varies according to severity and individual risk; prescription antiviral treatment may be considered by a healthcare professional in appropriate cases.

Suggestions: Follow current public-health guidance, maintain hygiene, stay home when unwell when appropriate, and seek medical help for serious symptoms

Tuberculosis (TB)

Tuberculosis is an infectious disease caused by *Mycobacterium tuberculosis*. It most commonly affects the lungs but can affect other parts of the body. Symptoms of pulmonary TB can include a persistent cough, fever, night sweating, tiredness, loss of appetite, and unexplained weight loss. TB can spread through the air when a person with infectious pulmonary TB coughs or sneezes.

TB requires proper medical diagnosis and a carefully supervised treatment course. Treatment generally involves a combination of specific anti-tuberculosis medicines taken for several months. Completing the entire prescribed treatment is extremely important because stopping treatment early can contribute to drug resistance.

Medicine information for dataset: TB requires a prescribed combination of anti-tuberculosis medicines under medical supervision.

Suggestions: Anyone with a persistent cough or other concerning symptoms should receive medical evaluation. People diagnosed with TB should follow their healthcare professional's treatment plan carefully

Malaria is a serious infectious disease caused by *Plasmodium* parasites and transmitted through the bites of infected mosquitoes. Symptoms commonly include fever, chills, sweating, headache, tiredness, and body aches. Some forms of malaria can become severe and may cause serious complications if treatment is delayed.

Treatment depends on the malaria species, geographical location, severity of infection, and other factors. Healthcare professionals use specific antimalarial medicines based on these factors. Malaria should not be treated by choosing a medicine independently because the appropriate treatment can vary.

Medicine information for dataset: Antimalarial treatment is selected by a healthcare professional according to the type and severity of malaria.

Suggestions: Seek medical testing when malaria is suspected, use mosquito-bite prevention measures, and obtain prompt medical care for persistent or severe fever.

Dengue is a viral infection transmitted mainly by infected *Aedes* mosquitoes. It can cause sudden fever, headache, pain behind the eyes, muscle and joint pain, nausea, and sometimes a skin rash. Most people recover with appropriate supportive care, but some cases can become severe.

There is no routine antibiotic treatment for dengue because it is caused by a virus. Management generally involves adequate fluids, rest, and medical monitoring. Some medicines can increase bleeding risk, so people with suspected dengue should not choose pain or fever medicines on their own without medical advice.

Medicine information for dataset: Supportive treatment and fluid management are commonly used; medication choices should be determined by a healthcare professional.

Suggestions: Stay hydrated, rest, get appropriate testing when dengue is suspected, and seek urgent medical attention if symptoms worsen or unusual bleeding, severe abdominal pain, persistent vomiting, or other warning signs occur.

Typhoid fever is an infection caused by *Salmonella Typhi*. It is commonly spread through contaminated food or water. Symptoms can include prolonged fever, headache, weakness, abdominal discomfort, diarrhea or constipation, and loss of appetite. Without appropriate treatment, the infection can become serious.

Typhoid generally requires antibiotics prescribed after appropriate medical evaluation. The choice of antibiotic depends on local antibiotic resistance patterns and the individual's circumstances. Completing the prescribed course is important when treatment is given.

Medicine information for dataset: Antibiotics may be required and should be selected and prescribed by a healthcare professional.

Suggestions: Drink safe water, maintain good hand hygiene, consume properly prepared food, and seek medical evaluation for persistent fever.

Cholera is an intestinal infection caused by the bacterium *Vibrio cholerae*. It can cause sudden, severe watery diarrhea and vomiting, which can lead to dangerous dehydration.

Severe dehydration can develop quickly, making prompt treatment particularly important.

The most important part of cholera management is rapid replacement of lost fluids and electrolytes. In severe cases, treatment may require intravenous fluids and antibiotics may be prescribed by healthcare professionals. Prevention depends heavily on safe drinking water, sanitation, and good hygiene.

Medicine information for dataset: Oral or intravenous rehydration is central to treatment; antibiotics may be prescribed in appropriate cases by healthcare professionals.

Suggestions: Seek medical care quickly for severe watery diarrhea, drink safe water, and maintain proper sanitation and hand hygiene.

Measles is a highly contagious viral infection. It commonly begins with fever, cough, runny nose, and red or watery eyes, followed by a characteristic skin rash. The infection can sometimes cause serious complications, particularly in vulnerable individuals.

There is no routine antibiotic treatment for uncomplicated measles because it is caused by a virus. Treatment generally focuses on supportive care, hydration, rest, and monitoring for complications. Vaccination is the most important method of preventing measles.

Medicine information for dataset: Supportive care is generally used; healthcare professionals may provide additional treatment depending on complications and individual circumstances.

Suggestions: Follow recommended vaccination schedules, avoid exposing others when infected, and seek medical attention if symptoms become severe.

Chickenpox is a contagious viral infection caused by the varicella-zoster virus. It commonly produces an itchy rash that develops into fluid-filled spots and later crusts over. Fever, tiredness, and loss of appetite can also occur. Most healthy children recover without serious complications, although complications can occur.

Treatment generally focuses on relieving symptoms and preventing complications. Keeping the skin clean and avoiding scratching can help reduce the risk of secondary infection. In certain situations, healthcare professionals may prescribe antiviral treatment, especially for people at increased risk of complications.

Medicine information for dataset: Symptom management is common; antiviral treatment may be prescribed in selected cases by a healthcare professional.

Suggestions: Avoid scratching the rash, maintain good hygiene, avoid close contact with vulnerable people, and follow vaccination recommendations.

Hepatitis means inflammation of the liver and can have several causes, including viruses such as hepatitis A, B, and C. Symptoms may include tiredness, nausea, abdominal discomfort, fever, dark urine, pale stools, and yellowing of the skin or eyes. Some types can become chronic and cause long-term liver problems.

Treatment depends strongly on the type and cause of hepatitis. Some forms mainly require supportive care, while chronic hepatitis B or hepatitis C may require specific antiviral treatment under specialist medical supervision. Because different types of hepatitis have different treatments, proper testing is important before treatment is selected.

Medicine information for dataset: Treatment varies according to the specific type of hepatitis; antiviral therapy may be prescribed for certain forms.

Suggestions: Seek appropriate testing, avoid sharing personal items that may carry blood, maintain good hygiene, and follow medical advice.

12. Pneumonia

Pneumonia is an infection or inflammation of the lungs that can be caused by bacteria, viruses, fungi, or other organisms. Symptoms may include cough, fever, chills, chest discomfort, tiredness, and difficulty breathing. The severity can range from mild to life-threatening, particularly in vulnerable individuals.

Treatment depends on the cause and severity. Bacterial pneumonia may require antibiotics prescribed by a healthcare professional, while viral pneumonia may require supportive care or specific antiviral treatment in selected circumstances. Severe pneumonia may require oxygen or hospital-based treatment.

Medicine information for dataset: Treatment depends on the cause; antibiotics may be prescribed for bacterial pneumonia, while other cases may require different treatment.

Suggestions: Seek medical evaluation for breathing difficulties, persistent high fever, chest pain, or worsening symptoms. Vaccination and good respiratory hygiene can help prevent some forms of pneumonia.

Suggestions:

Maintain hydration, rest, seek appropriate testing, and obtain medical attention if warning signs or severe symptoms develop.

Hypertension is a condition in which blood pressure remains higher than the recommended range over time. It often does not cause noticeable symptoms, which is why it is sometimes called a “silent” condition. When blood pressure remains uncontrolled for a long period, it can increase the risk of problems affecting the heart, brain, kidneys, and blood vessels.

Treatment usually involves lifestyle changes and, when needed, medicines prescribed by a healthcare professional. Regular blood-pressure monitoring, reducing excessive salt intake, maintaining healthy activity, getting adequate sleep, and following the prescribed treatment plan can help control blood pressure.

Medicine information: Blood-pressure-lowering medicines may be prescribed depending on the individual's blood pressure and health history.

Suggestions: Monitor blood pressure regularly and follow medical advice. Seek medical attention for extremely high readings accompanied by symptoms such as severe headache, chest pain, confusion, or difficulty breathing.

Coronary artery disease occurs when the blood vessels supplying the heart become narrowed or blocked, commonly because of the buildup of fatty deposits. It can reduce blood flow to the heart muscle and may cause chest discomfort, shortness of breath, tiredness, or symptoms during physical activity.

Treatment depends on the severity of the disease and may include lifestyle changes, prescribed medicines, and sometimes medical procedures. Controlling blood pressure, cholesterol, diabetes, and other risk factors is important. Regular medical follow-up can help reduce the risk of complications.

Medicine information: Healthcare professionals may prescribe medicines to control cholesterol, blood pressure, blood clotting, or heart-related symptoms depending on the individual.

Suggestions: Seek urgent medical care for new or severe chest pain, especially when accompanied by breathing difficulty, sweating, faintness, or pain spreading to the arm or jaw.

Heart failure occurs when the heart cannot pump blood effectively enough to meet the body's needs. It can cause symptoms such as shortness of breath, tiredness, swelling of the legs or ankles, and difficulty breathing when lying down. Heart failure can develop because of conditions such as coronary artery disease, high blood pressure, or damage to the heart muscle.

Treatment is individualized and may include lifestyle changes, monitoring, prescribed medicines, and sometimes devices or procedures. Managing the underlying cause is an important part of treatment. Regular follow-up helps healthcare professionals monitor the condition and adjust treatment.

Medicine information: Several types of heart medicines may be prescribed depending on the type and severity of heart failure.

Suggestions: Follow the prescribed treatment plan and monitor symptoms. Sudden severe breathing difficulty, chest pain, or fainting requires urgent medical attention.

Anemia occurs when the blood does not have enough healthy red blood cells or hemoglobin to carry oxygen effectively. It can cause tiredness, weakness, dizziness, headaches, and shortness of breath, depending on its severity. Iron deficiency is one common cause, but anemia can also result from vitamin deficiencies, chronic diseases, blood loss, or inherited conditions.

Treatment depends on the underlying cause. Some people may need nutritional changes or supplements, while others require treatment for an underlying disease. It is important to determine the cause rather than taking supplements without appropriate evaluation.

Medicine information: Iron or vitamin supplements may be recommended when a deficiency is confirmed, while other types of anemia require different treatment.

Suggestions: Eat a varied, nutritious diet and seek medical evaluation for persistent tiredness, weakness, or unexplained symptoms

Sickle cell disease is an inherited blood disorder in which red blood cells can become abnormally shaped. These cells can interfere with normal blood flow and may cause episodes of pain, anemia, tiredness, and increased susceptibility to certain complications. It is a lifelong condition that requires ongoing medical care.

Management focuses on preventing complications, treating symptoms, maintaining adequate hydration, and monitoring overall health. Some people may receive specialized medicines or other treatments depending on their condition. Regular follow-up with a healthcare team is important.

Medicine information: Specialized medicines may be prescribed to reduce complications or manage symptoms. Treatment must be individualized by a healthcare professional.

Suggestions: Maintain hydration, follow the medical care plan, attend regular checkups, and seek medical attention for severe or unusual symptoms.

Leukemia is a type of cancer affecting blood-forming tissues, particularly the bone marrow. It leads to the production of abnormal blood cells and can cause symptoms such as tiredness, frequent infections, easy bruising, fever, or unexplained bleeding. There are several types of leukemia, and they can behave differently.

Treatment depends on the specific type and may include chemotherapy, targeted therapy, immunotherapy, radiation in selected situations, or stem-cell transplantation. Treatment should always be managed by a specialist oncology team.

Medicine information: Cancer medicines are selected according to the specific leukemia type and individual circumstances.

Suggestions: Persistent unexplained symptoms should be evaluated by a healthcare professional. People diagnosed with leukemia should follow their oncology team's treatment and monitoring plan

Asthma is a chronic condition in which the airways become inflamed and sensitive. It can cause wheezing, coughing, chest tightness, and difficulty breathing. Symptoms may be triggered by allergens, exercise, respiratory infections, smoke, or other environmental factors.

Asthma is usually managed with an individualized treatment plan. Inhaled medicines are commonly used to control airway inflammation and relieve symptoms. Identifying and avoiding personal triggers can also help reduce episodes.

Medicine information: Inhaled medicines may be prescribed for long-term control or quick symptom relief depending on the individual's asthma plan.

Suggestions: Follow the prescribed asthma action plan and avoid known triggers. Severe difficulty breathing requires urgent medical attention.

COPD is a long-term lung disease that causes persistent airflow limitation. It includes conditions such as chronic bronchitis and emphysema. Common symptoms include long-lasting cough, mucus production, wheezing, tiredness, and shortness of breath.

Treatment focuses on reducing symptoms, improving lung function, preventing flare-ups, and maintaining activity. Inhaled medicines, pulmonary rehabilitation, vaccinations, and oxygen therapy in selected cases may be part of treatment.

Medicine information: Healthcare professionals may prescribe inhalers and other treatments based on the severity of COPD.

Suggestions: Avoid tobacco smoke and other lung irritants, follow the prescribed treatment plan, and seek medical care when breathing symptoms suddenly worsen.

Bronchitis occurs when the bronchial tubes become inflamed. Acute bronchitis is often caused by viruses and may cause cough, mucus production, tiredness, and mild fever. The cough can sometimes continue after the original infection has improved.

Treatment for uncomplicated acute bronchitis generally focuses on rest, fluids, and symptom management. Antibiotics are not usually helpful when the cause is viral. Chronic bronchitis is a form of COPD and requires longer-term medical management.

Medicine information: Treatment depends on the cause and type of bronchitis; antibiotics are not routinely useful for viral acute bronchitis.

Suggestions: Rest, drink adequate fluids, avoid smoke and other irritants, and seek medical evaluation if breathing becomes difficult or symptoms worsen.

Pneumonia is an infection or inflammation of the lungs caused by bacteria, viruses, fungi, or other organisms. Symptoms may include cough, fever, chills, chest discomfort, tiredness, and difficulty breathing. Some cases are mild, while others can become serious.

Treatment depends on the cause and severity. Bacterial pneumonia may require prescribed antibiotics, while viral cases may require supportive care or specific antiviral treatment in selected situations. Severe pneumonia may require hospital treatment and oxygen.

Medicine information: Treatment is determined according to the cause and severity by a healthcare professional.

Suggestions: Seek medical attention for persistent fever, breathing difficulties, chest pain, or worsening symptoms.

Pulmonary tuberculosis is tuberculosis affecting the lungs and is caused by *Mycobacterium tuberculosis*. Symptoms can include a persistent cough, fever, night sweating, tiredness, reduced appetite, and unexplained weight loss. Infectious pulmonary TB can spread through the air.

Treatment requires a specific combination of anti-tuberculosis medicines for an extended period. The exact treatment depends on the type of TB and drug-resistance pattern. Completing the prescribed treatment is extremely important.

Medicine information: TB requires a supervised combination of anti-tuberculosis medicines prescribed by a healthcare professional.

Suggestions: Seek medical evaluation for a persistent cough or other concerning symptoms and follow the complete treatment plan if diagnosed.

Epilepsy is a neurological disorder characterized by a tendency to have recurrent seizures. Seizures can affect awareness, movement, sensation, or behavior depending on the part of the brain involved. A single seizure does not necessarily mean a person has epilepsy.

Treatment depends on the type and frequency of seizures. Healthcare professionals may prescribe anti-seizure medicines and monitor the person's response. Some people with difficult-to-control epilepsy may require specialized treatments.

Medicine information: Anti-seizure medicines may be prescribed based on the seizure type and individual circumstances.

Suggestions: Follow the prescribed treatment plan and attend neurological follow-ups. A prolonged seizure or repeated seizures without recovery require emergency medical attention.

Migraine is a neurological condition that can cause recurring headaches, often accompanied by nausea and sensitivity to light or sound. Some people experience an “aura” involving temporary visual or sensory changes before or during a migraine. Triggers can vary between individuals.

Management includes identifying triggers, maintaining regular sleep and meals, staying hydrated, and using appropriate treatments recommended by a healthcare professional. People with frequent migraines may be offered preventive treatment.

Medicine information: Medicines for acute attacks or prevention may be recommended based on migraine frequency and symptoms.

Suggestions: Keep track of possible triggers and headache patterns. A sudden, extremely severe headache or a headache accompanied by neurological symptoms requires prompt medical evaluation.

Parkinson's disease is a progressive neurological disorder that affects movement. Common symptoms include tremor, slow movement, muscle stiffness, and balance difficulties. Some people may also experience sleep, mood, or other non-movement symptoms.

There is currently no simple cure, but treatments can help manage symptoms and maintain quality of life. Treatment may include medicines, physical therapy, exercise programs, and other supportive approaches depending on the individual.

Medicine information: Medicines that affect brain signaling may be prescribed to improve movement symptoms.

Suggestions: Regular medical follow-up, physical activity appropriate to the person's condition, and safety measures can be helpful.

Alzheimer's disease is a progressive brain disorder that primarily affects memory and thinking. Early symptoms can include difficulty remembering recent information, problems with planning, and changes in everyday functioning. As the disease progresses, additional cognitive and behavioral difficulties can occur.

There is no simple cure, but treatment and supportive care can help manage symptoms and maintain independence for as long as possible. Healthcare professionals may recommend medicines, cognitive support, regular routines, and assistance with daily activities.

Medicine information: Certain prescription medicines may help manage symptoms or slow progression in selected patients.

Suggestions: Regular medical evaluation, a safe environment, consistent routines, and support from caregivers can be beneficial.

Multiple sclerosis is a disease in which the immune system attacks the protective covering of nerve fibers in the central nervous system. Symptoms vary considerably and may include vision problems, numbness, weakness, balance difficulties, and fatigue. Some people experience symptoms in episodes, while others have progressive disease.

Treatment aims to reduce relapses, slow disease progression, and manage symptoms. Disease-modifying treatments may be prescribed by specialists. Rehabilitation and physical therapy can also help with specific difficulties.

Medicine information: Disease-modifying and symptom-management medicines may be prescribed according to the individual's MS type and condition.

Suggestions: Maintain regular neurological follow-up and report new or worsening neurological symptoms to a healthcare professional.

Meningitis is inflammation of the membranes surrounding the brain and spinal cord. It can be caused by viruses, bacteria, fungi, or other organisms. Symptoms may include fever, headache, neck stiffness, vomiting, sensitivity to light, and changes in alertness.

Some forms, particularly bacterial meningitis, can become life-threatening very quickly and require urgent medical treatment. Treatment depends on the cause and may involve hospital care and specific antimicrobial medicines.

Medicine information: Treatment depends on the cause and must be determined urgently by healthcare professionals.

Suggestions: Fever accompanied by severe headache, neck stiffness, confusion, seizures, or rapidly worsening illness requires emergency medical attention.

Gastritis is inflammation of the stomach lining. It may cause upper abdominal discomfort, nausea, vomiting, bloating, or indigestion. Causes can include certain infections, medications, alcohol exposure, or other conditions.

Treatment depends on the underlying cause. Healthcare professionals may recommend medicines that reduce stomach acid or treat an underlying infection when appropriate. Avoiding known stomach irritants can also help.

Medicine information: Acid-reducing medicines or other treatments may be recommended depending on the cause.

Suggestions: Follow medical advice, avoid known stomach irritants, and seek evaluation for persistent abdominal pain or bleeding symptoms.

Gastroenteritis is inflammation of the stomach and intestines, commonly caused by infections. It can cause diarrhea, vomiting, abdominal cramps, nausea, and sometimes fever. The greatest immediate concern can be dehydration, especially when vomiting or diarrhea is severe.

Treatment mainly focuses on replacing fluids and electrolytes and allowing the digestive system to recover. Antibiotics are not routinely needed for viral gastroenteritis and should only be used when medically indicated.

Medicine information: Treatment depends on the cause; fluid and electrolyte replacement is particularly important.

Suggestions: Drink appropriate fluids, maintain hand hygiene, and seek medical care if there are signs of significant dehydration or severe illness.

Peptic ulcer disease involves open sores developing in the lining of the stomach or the first part of the small intestine. Common causes include *Helicobacter pylori* infection and certain medications. Symptoms may include burning or aching abdominal pain, indigestion, nausea, or bloating.

Treatment depends on the cause. When *H. pylori* is present, healthcare professionals generally use a combination of medicines to eliminate the infection and reduce stomach acid. Medication-related ulcers may require changes in treatment under medical supervision.

Medicine information: Treatment may include acid-reducing medicines and, when *H. pylori* is confirmed, a prescribed combination of medicines.

Suggestions: Seek medical attention for persistent symptoms or signs of gastrointestinal bleeding.

Crohn's disease is a chronic inflammatory bowel disease that can affect different parts of the digestive tract. Symptoms may include abdominal pain, diarrhea, tiredness, and reduced appetite. Symptoms can occur in periods of flare-ups and remission.

Treatment aims to control inflammation, reduce symptoms, and prevent complications. Depending on severity, treatment may include anti-inflammatory or immune-targeting medicines and nutritional support under specialist supervision.

Medicine information: Specialized medicines may be prescribed to control inflammation and maintain remission.

Suggestions: Maintain regular medical follow-up and report persistent abdominal pain, diarrhea, or other worsening symptoms.

Ulcerative colitis is a chronic inflammatory bowel disease that primarily affects the colon and rectum. Symptoms can include diarrhea, abdominal pain, urgency to use the bathroom, and blood in the stool. Symptoms may alternate between flare-ups and periods of remission.

Treatment aims to control inflammation and maintain remission. Healthcare professionals may prescribe anti-inflammatory medicines, immune-targeting treatments, or other therapies depending on severity.

Medicine information: Treatment is individualized and may involve prescription medicines to reduce intestinal inflammation.

Suggestions: Follow the treatment plan and seek medical attention for severe symptoms, significant bleeding, or dehydration.

Hepatitis refers to inflammation of the liver and can have several causes, including viral infections, medications, alcohol, and other conditions. Symptoms may include tiredness, nausea, abdominal discomfort, dark urine, and yellowing of the skin or eyes.

Treatment depends on the specific cause. Some forms require mainly supportive care, while chronic viral hepatitis may require antiviral treatment. Proper testing is necessary to determine the cause and appropriate treatment.

Medicine information: Treatment varies according to the type and cause of hepatitis and should be determined by a healthcare professional.

Suggestions: Seek appropriate testing and medical follow-up, and avoid sharing personal items that could transmit blood.

Gallstones are solid particles that form inside the gallbladder. Some people have no symptoms, while others experience sudden pain in the upper right or central abdomen, particularly after eating. Gallstones can sometimes block bile ducts and cause complications.

Treatment depends on whether symptoms or complications occur. People with recurring symptoms may require medical evaluation and, in some cases, a procedure to remove the gallbladder.

Medicine information: Symptom management may be used in selected cases, but recurrent or complicated gallstones may require a medical procedure.

Suggestions: Seek medical attention for severe abdominal pain, fever, vomiting, or yellowing of the skin or eyes.

Diabetes is a condition in which blood glucose levels remain too high because the body does not produce enough insulin, does not use insulin effectively, or both. Symptoms can include increased thirst, frequent urination, tiredness, blurred vision, and unexplained changes in body condition.

Management involves monitoring blood glucose, balanced nutrition, physical activity appropriate for the individual, and prescribed treatment. The treatment differs between type 1, type 2, and other forms of diabetes.

Medicine information: Diabetes medicines or insulin may be prescribed depending on the type of diabetes and individual needs.

Suggestions: Monitor glucose as recommended and attend regular medical checkups. Avoid changing diabetes medication without professional guidance.

Hypothyroidism occurs when the thyroid gland does not produce enough thyroid hormones. Symptoms can include tiredness, feeling unusually cold, dry skin, constipation, and difficulty concentrating. Symptoms can develop gradually and may be mistaken for other conditions.

Treatment usually involves thyroid hormone replacement prescribed according to blood-test results. Regular testing is used to ensure the treatment level is appropriate.

Medicine information: Thyroid hormone replacement may be prescribed by a healthcare professional.

Suggestions: Take prescribed treatment consistently and attend recommended thyroid-function tests.

Hyperthyroidism occurs when the thyroid gland produces too much thyroid hormone. It can cause increased heart rate, sweating, nervousness, tremors, difficulty sleeping, and heat intolerance. Some people may also experience changes in weight or appetite.

Treatment depends on the cause and may include medicines, radioactive iodine treatment, or surgery in selected situations. Medical monitoring is important because thyroid hormone levels can affect the heart and other organs.

Medicine information: Medicines that reduce thyroid activity or control symptoms may be prescribed depending on the cause.

Suggestions: Follow medical advice and seek evaluation for persistent rapid heartbeat or other concerning symptoms.

Obesity is a chronic health condition involving excessive body fat that can increase the risk of conditions such as diabetes, high blood pressure, heart disease, and joint

problems. It is influenced by multiple factors, including genetics, environment, lifestyle, medications, and other health conditions.

Management should focus on overall health rather than rapid changes in body size. Healthcare professionals may recommend balanced nutrition, appropriate physical activity, adequate sleep, behavioral support, and medical treatment when appropriate.

Medicine information: In some cases, healthcare professionals may consider prescription weight-management treatment based on individual health factors.

Suggestions: Focus on sustainable healthy habits and seek professional guidance rather than using extreme diets or unapproved weight-loss products.

Osteoporosis is a condition in which bones become weaker and more likely to fracture. It often develops without obvious symptoms until a fracture occurs. Risk can increase with age and certain medical conditions or medications.

Treatment may include appropriate nutrition, physical activity, fall prevention, and medicines for people at increased fracture risk. Calcium and vitamin D requirements should be discussed with a healthcare professional.

Medicine information: Bone-strengthening medicines may be prescribed when fracture risk is sufficiently high.

Suggestions: Maintain bone-healthy nutrition, participate in appropriate weight-bearing activity, and discuss bone-density testing with a healthcare professional when appropriate.

Arthritis refers to a group of conditions that cause inflammation or damage in joints. Symptoms may include joint pain, stiffness, swelling, and reduced movement. Different types of arthritis have different causes and treatments.

Management may include physical activity, physical therapy, symptom-relief treatment, and medicines depending on the type. Early evaluation can help identify the specific form of arthritis and guide treatment.

Medicine information: Treatment varies according to the type of arthritis and may include pain-relief or anti-inflammatory medicines under medical guidance.

Suggestions: Maintain appropriate joint movement and seek evaluation for persistent pain or swelling.

Osteoarthritis is a degenerative joint condition in which the tissues within a joint gradually change. It commonly affects the knees, hips, hands, and spine. Symptoms include pain, stiffness, and difficulty moving the affected joint.

Management may include appropriate physical activity, physical therapy, supportive measures, and symptom-relief medicines when needed. Treatment is individualized according to the affected joint and severity.

Medicine information: Pain-relief or anti-inflammatory treatment may be recommended by healthcare professionals.

Suggestions: Maintain appropriate movement and seek professional advice for persistent or worsening joint pain.

Rheumatoid arthritis is an autoimmune disease in which the immune system attacks the joints. It can cause joint pain, swelling, stiffness, and fatigue. Unlike ordinary wear-and-tear arthritis, rheumatoid arthritis can cause ongoing inflammation and joint damage.

Treatment aims to control inflammation and prevent long-term joint damage. Specialists may prescribe disease-modifying medicines and other treatments depending on disease activity.

Medicine information: Disease-modifying medicines may be prescribed and require regular medical monitoring.

Suggestions: Early diagnosis and consistent follow-up are important. Report persistent joint swelling or prolonged morning stiffness to a healthcare professional.

Gout is a form of inflammatory arthritis caused by the buildup of uric acid crystals in a joint. It can cause sudden episodes of severe joint pain, swelling, warmth, and redness, often affecting one joint at a time.

Treatment involves managing acute attacks and reducing the risk of future attacks. Healthcare professionals may prescribe medicines for acute inflammation and, when appropriate, long-term treatment to control uric acid levels.

Medicine information: Treatment should be selected according to the individual's health status and whether the goal is treating an acute attack or preventing future attacks.

Suggestions: Seek medical evaluation for sudden severe joint pain, particularly if this is the first episode or if fever is present.

Osteoporosis causes bones to become fragile and increases the risk of fractures. It may not produce symptoms until a fracture occurs. Risk factors include aging, certain medical conditions, hormonal changes, and some long-term medicines.

Treatment focuses on reducing fracture risk through appropriate nutrition, physical activity, fall prevention, and medicines when indicated. Bone-density testing may help healthcare professionals assess fracture risk.

Medicine information: Prescription bone-strengthening treatments may be recommended for people with significant fracture risk.

Suggestions: Maintain bone health, prevent falls, and follow medical recommendations regarding bone-density monitoring.

Acne is a common skin condition that occurs when hair follicles become blocked with oil and dead skin cells. It can cause blackheads, whiteheads, pimples, and sometimes deeper inflammatory lesions. It commonly affects the face, chest, shoulders, and back.

Treatment depends on the severity and type of acne. Gentle skin care and appropriate topical treatments may help. More persistent or severe acne may require evaluation by a dermatologist.

Medicine information: Topical treatments may be recommended for mild acne, while prescription treatments may be used for more severe cases.

Suggestions: Avoid repeatedly picking or squeezing spots and use gentle skin-care products. Seek dermatological advice if acne is persistent or severe.

Eczema is a group of conditions that cause dry, itchy, and inflamed skin. Symptoms can appear as patches of redness, dryness, irritation, or scaling. Triggers can include irritants, allergens, weather changes, and other factors.

Management focuses on protecting the skin barrier and controlling inflammation. Regular use of appropriate moisturizers is often important. Healthcare professionals may recommend topical treatments during flare-ups.

Medicine information: Moisturizers and prescribed anti-inflammatory skin treatments may be used depending on severity.

Suggestions: Keep the skin moisturized, avoid known irritants, and seek medical advice for severe or infected-looking skin lesions.

Psoriasis is a chronic immune-mediated skin condition that causes skin cells to build up too quickly. It can produce thick, scaly patches and may also affect the nails or joints. Symptoms can come and go and may be triggered by infections, stress, or other factors.

Treatment depends on the type and severity of psoriasis. Options may include topical treatments, light therapy, and systemic medicines prescribed by specialists for more extensive disease.

Medicine information: Treatment may involve topical or systemic prescription therapies depending on severity.

Suggestions: Follow dermatological treatment plans and avoid scratching irritated skin. Joint pain in someone with psoriasis should be discussed with a healthcare

Vitiligo is a condition in which areas of skin lose pigment, producing lighter or white patches. It occurs when pigment-producing cells are damaged or lost. It is not contagious and can affect people of any age.

Treatment is individualized and may include topical medicines, light-based therapy, or other specialist treatments. Some people choose no medical treatment and instead focus on skin protection and cosmetic approaches.

Medicine information: Dermatologists may recommend topical or light-based treatments depending on the individual.

Suggestions: Protect depigmented areas from excessive sunlight and seek dermatological evaluation if new patches appear.

Ringworm is a fungal infection of the skin and is not caused by a worm. It can cause circular, itchy, scaly patches and can occur on different parts of the body. It can spread through direct contact with infected people, animals, or contaminated objects.

Treatment usually involves appropriate antifungal treatment. The exact treatment depends on the location and severity of the infection. Keeping the affected area clean and dry can help reduce spread.

Medicine information: Antifungal treatments may be used, with the appropriate type depending on the location and severity.

Suggestions: Avoid sharing towels or clothing, maintain good hygiene, and seek medical advice if the infection spreads or does not improve.

Scabies is a contagious skin infestation caused by microscopic mites that burrow into the skin. It commonly causes intense itching, particularly at night, along with small bumps or a rash. It spreads mainly through prolonged close skin-to-skin contact.

Treatment requires a specific anti-parasitic medicine prescribed or recommended by a healthcare professional. Close contacts may also need evaluation and treatment to prevent reinfection. Clothing, bedding, and towels should be handled according to medical/public-health guidance.

Medicine information: Prescription or recommended anti-parasitic skin treatment is generally required; treatment should be followed exactly as directed.

Suggestions: Seek medical evaluation, avoid close contact until appropriate treatment has been completed, and follow guidance for treating household contacts and belongings.

Tooth decay, also called dental caries or cavities, occurs when bacteria in the mouth produce acids that gradually damage the tooth's outer layers. It can cause sensitivity to hot or cold foods, pain, visible spots or holes in the tooth, and sometimes difficulty chewing. If untreated, decay can progress deeper into the tooth and affect the dental pulp.

Treatment depends on how advanced the decay is. Early damage may sometimes be managed with fluoride treatment and improved oral hygiene, while established cavities may require a filling. If the decay reaches the inner part of the tooth, a dentist may recommend root-canal treatment or another dental procedure.

Medicine information: Fluoride products may help prevent or manage early tooth decay. Pain-relief medicines may sometimes be used for discomfort, but they do not remove the underlying decay.

Suggestions: Brush twice daily with fluoride toothpaste, clean between the teeth, limit frequent sugary foods and drinks, and visit a dentist regularly.

Toothache is pain or discomfort in or around a tooth. It can be caused by tooth decay, a cracked tooth, gum disease, dental infection, or problems involving the tooth's inner tissues. The pain may be constant or may become worse when eating, drinking, or applying pressure.

Treatment depends on the underlying cause. A dentist needs to examine the tooth to determine whether treatment such as a filling, cleaning, root-canal treatment, or another procedure is necessary. Pain-relief treatment may temporarily reduce discomfort but does not fix the dental problem.

Medicine information: Appropriate pain-relief medicine may sometimes be recommended by a healthcare professional, depending on age and health conditions.

Antibiotics are not routinely needed for ordinary toothache and should only be used when a dentist or doctor determines that there is an appropriate bacterial infection.

Suggestions: Arrange a dental examination if toothache persists. Seek urgent dental/medical care if pain occurs with facial swelling, fever, difficulty swallowing, or difficulty breathing.

Gingivitis is inflammation of the gums, usually caused by the buildup of plaque around the teeth. Common signs include red or swollen gums, tenderness, and bleeding during brushing or flossing. Gingivitis is an early form of gum disease and can often be improved when plaque is properly controlled.

Treatment mainly involves improving oral hygiene and removing plaque and tartar. A dental professional may perform a professional cleaning if necessary. If gingivitis is not managed, it can progress to more serious gum disease that may damage the tissues supporting the teeth.

Medicine information: Treatment primarily involves oral hygiene and professional dental cleaning. Specific mouth rinses or other treatments may be recommended by a dental professional when appropriate.

Suggestions: Brush gently twice a day with fluoride toothpaste, clean between teeth daily, avoid tobacco exposure, and attend regular dental checkups.

Mouth ulcers are small sores that can develop inside the mouth, including on the tongue, cheeks, lips, or gums. They may cause pain or burning, particularly while eating spicy, acidic, or salty foods. Minor ulcers can occur after accidental biting, irritation, stress, or other causes and often heal on their own.

Treatment generally focuses on reducing irritation and managing discomfort while the ulcer heals. Avoiding foods that make the ulcer painful and maintaining good oral hygiene can help. Persistent, unusually large, frequently recurring, or unexplained ulcers should be evaluated by a healthcare professional or dentist.

Medicine information: Protective or pain-relieving oral treatments may sometimes be recommended by a healthcare professional depending on the cause and severity.

Suggestions: Avoid irritating foods, maintain gentle oral hygiene, drink adequate fluids, and seek professional evaluation if an ulcer does not heal or keeps returning.

A dental abscess is a collection of infected material that develops because of a bacterial infection around a tooth or gum. It can cause severe tooth pain, swelling, sensitivity, fever, and sometimes an unpleasant taste in the mouth. The infection can spread to surrounding tissues if it is not treated.

A dental abscess requires professional dental treatment rather than simply treating the pain. Depending on the location and severity, a dentist may drain the infection, perform

root-canal treatment, or remove the affected tooth. Antibiotics may sometimes be prescribed when there are signs of spreading infection or systemic illness.

Medicine information: Antibiotics are used only when medically indicated and should be prescribed by a dentist or doctor. Pain-relief treatment may help with symptoms but does not eliminate the abscess.

Suggestions: Seek dental care promptly. **Facial or neck swelling with difficulty breathing or swallowing is an emergency and requires immediate medical attention.**

Acidity commonly refers to symptoms caused by stomach acid irritating the upper digestive tract. It may cause a burning sensation in the upper abdomen or chest, sour taste in the mouth, belching, or discomfort after eating. Symptoms can sometimes become worse after large meals or when lying down soon after eating.

Management usually involves eating regular balanced meals, avoiding foods that personally trigger symptoms, and not lying down immediately after eating. If acidity occurs frequently or is severe, a healthcare professional should evaluate the cause because conditions such as GERD or gastritis may be involved.

Medicine information: Healthcare professionals may recommend acid-reducing or acid-neutralizing medicines depending on the cause and frequency of symptoms.

Suggestions: Eat slowly, avoid very large meals, stay upright after eating, and seek medical advice if symptoms are frequent or persistent.

Indigestion, also called dyspepsia, refers to discomfort in the upper part of the digestive system. It can cause fullness, bloating, belching, nausea, or a burning sensation after eating. It may occur occasionally after eating too quickly, eating a large meal, or consuming foods that trigger symptoms.

Treatment depends on the underlying cause. Simple cases may improve with changes in eating habits and avoiding individual triggers. Persistent indigestion can sometimes be associated with gastritis, ulcers, GERD, or other digestive conditions and may require medical evaluation.

Medicine information: Depending on the cause, a healthcare professional may recommend medicines that reduce stomach acid or relieve digestive symptoms.

Suggestions: Eat slowly, avoid overeating, keep track of foods that trigger symptoms, and consult a healthcare professional if indigestion repeatedly occurs.

Constipation occurs when bowel movements become less frequent or difficult to pass, or when stools become hard and dry. It can cause abdominal discomfort, bloating, and a feeling that the bowel has not completely emptied. Common contributing factors

include inadequate dietary fiber, insufficient fluid intake, reduced physical activity, and changes in routine.

Management often involves increasing fiber gradually, drinking adequate fluids, and maintaining regular physical activity. Some people may need short-term treatment recommended by a healthcare professional. Persistent constipation should be evaluated to identify an underlying cause.

Medicine information: Fiber supplements or certain laxatives may sometimes be recommended, but they should be used appropriately and according to professional guidance.

Suggestions: Include fiber-rich foods, drink adequate fluids, stay physically active, and seek medical advice if constipation persists or occurs with significant abdominal pain or bleeding.

Diarrhea involves frequent passage of loose or watery stools. It can result from infections, contaminated food or water, certain medicines, food intolerance, or digestive conditions. It may be accompanied by abdominal cramps, nausea, vomiting, or fever.

The main concern with significant diarrhea is dehydration because the body loses fluids and electrolytes. Drinking appropriate fluids and replacing electrolytes can help. Treatment depends on the cause, and some cases require medical evaluation, particularly when diarrhea is severe or persistent.

Medicine information: Treatment depends on the cause. Some symptom-relief medicines may be appropriate in certain situations, while others should be avoided when specific symptoms or infections are present.

Suggestions: Maintain hydration, practice good hand hygiene, and seek medical care for severe dehydration, persistent diarrhea, blood in the stool, or significant illness.

Gastroenteritis is inflammation of the stomach and intestines, commonly caused by viral or bacterial infections. It can produce diarrhea, vomiting, abdominal cramps, nausea, and sometimes fever. It can spread through contaminated food, water, or contact with infected people.

Treatment mainly focuses on replacing fluids and electrolytes and allowing the digestive system to recover. Antibiotics are not routinely useful for viral gastroenteritis and are only appropriate for certain bacterial infections when prescribed by a healthcare professional.

Medicine information: Treatment depends on the cause; fluid and electrolyte replacement is particularly important.

Suggestions: Drink adequate fluids, wash hands thoroughly, prepare food safely, and seek medical attention if there is severe dehydration, persistent vomiting, blood in the stool, or worsening symptoms.

Gastritis is inflammation of the stomach lining. It can cause upper abdominal pain or discomfort, nausea, vomiting, bloating, and indigestion. Causes can include *Helicobacter pylori* infection, certain medicines, alcohol exposure, or other conditions.

Treatment depends on the underlying cause. Healthcare professionals may recommend acid-reducing treatment or treatment for an underlying infection when appropriate. Identifying and managing the cause is important because untreated gastritis can sometimes lead to complications.

Medicine information: Acid-reducing medicines or other treatments may be recommended depending on the cause. If *H. pylori* infection is confirmed, specific prescription treatment may be required.

Suggestions: Avoid known stomach irritants, follow professional advice, and seek medical evaluation for persistent symptoms or signs of gastrointestinal bleeding.

antacids and acid-reducing medicines for acidity, indigestion, gastritis, and GERD; fiber supplements and bowel-regulating medicines for constipation and hemorrhoids; oral rehydration solutions and appropriate symptom-management treatments for diarrhea, gastroenteritis, and food poisoning; prescribed antibiotics or antimicrobial treatments for selected bacterial infections such as typhoid, cholera, tuberculosis, and dental abscess; antiviral or supportive treatments for viral infections; blood-pressure-lowering medicines for hypertension; cholesterol-lowering and heart medicines for coronary artery disease; heart-failure medicines for heart failure; iron or vitamin replacement for certain types of anemia; specialized medicines for sickle-cell disease and leukemia; inhaled medicines for asthma and COPD; anti-seizure medicines for epilepsy; migraine-specific and preventive treatments for migraine; neurological medicines for Parkinson's disease, Alzheimer's disease, and multiple sclerosis; anti-inflammatory and immune-modifying medicines for rheumatoid arthritis, Crohn's disease, ulcerative colitis, psoriasis, and other inflammatory conditions; thyroid hormone replacement for hypothyroidism and thyroid-suppressing treatments for hyperthyroidism; diabetes medicines or insulin for diabetes; bone-strengthening treatments for osteoporosis; pain-relief and anti-inflammatory treatments for arthritis, osteoarthritis, and gout; topical treatments for acne, eczema, psoriasis, vitiligo, ringworm, and scabies; and dental treatments such as fluoride products, topical oral treatments, and dentist-directed treatment for tooth decay, toothache, gingivitis, mouth ulcers, and dental abscess.

