

Arsenicosis and malnutrition

Kunal Kanti Majumdar

Professor, Department of Community Medicine, KPC Medical College & Hospital, Jadavpur, Kolkata, India

Submission: 07-05-2016

Revision: 20-05-2016

Publication: 29-06-2016

Arsenic exposure through drinking water is a major health problem affecting many countries in the World, viz., Bangladesh, India, Argentina, Mongolia, China, Chili, Thailand, Taiwan, Mexico, and some parts of USA.¹⁻⁴ Prolonged exposure of arsenic of 5–90 µg/kg body weight/day results in arsenicosis,⁵ characterized by hyper- and hypopigmentation, keratosis, various systemic manifestations like weakness, anemia, chronic lung disease, peripheral neuropathy, liver fibrosis, gangrene of limbs, and cancers of skin, lungs and urinary bladder.⁶⁻⁸ Studies on populations of Taiwan, India, and Argentina exposed to arsenic through drinking water have suggested that malnutrition increases the risk of arsenic-induced diseases.⁹⁻¹³ Several human studies have identified associations between malnutrition and arsenic-induced skin lesions, skin cancer, and cardiovascular effects.^{12,14,15} Literature surveys indicate arsenic resistance and its relation with nutritional status.¹⁶⁻¹⁹ Laboratory experiments have demonstrated that specific micronutrients can modify arsenic metabolism and toxicity.¹³ Inhabitants of Taiwan and the Antofagasta region in northern Chile with severe health effects due to ingestion of high arsenic contaminated drinking water were reported to have a poor nutritional status.²⁰

Inorganic arsenic is metabolized to monomethylarsonic acid (MMA) and dimethylarsinic acid (DMA), and this methylation facilitates urinary arsenic excretion, which is dependent on availability of S-adenosylmethionine (SAM). The methyl group from SAM may be derived from dietary components such as methionine, choline, folate, and other nutrients.^{21,22} Dietary protein, iron, zinc, and niacin are associated with urinary excretion of MMA and DMA.²³ Previous studies suggest that persons with more complete methylation have a lower risk of adverse arsenic-related health outcomes.²⁴ Diet poor in methionine in protein is likely to decrease the ability to methylate arsenic and increase the arsenic toxicity.²⁵ Folate and cobalamin (vitamin B12) have been suggested to play an important role in the detoxification of ingested arsenic.²⁶ Studies carried out in experimental animals have shown that severe protein deficiencies can impair arsenic methylation and excretion.²² Vitamin C reduces the toxicity of arsenic.²⁷

Intake of carbohydrate, protein, animal protein, fat, vitamins, and minerals was estimated from a diet survey by 24-hour recall method in arsenic exposed study population of south 24 parganas and it was found in that study that

deficiencies of some nutrients like animal protein, calcium, fiber, folate, and vitamin C may increase the risk of arsenic-induced skin lesions.²⁸ Some researchers have postulated that deficiencies in some nutrients such as beta-carotene, methionine, and zinc may increase susceptibility to arsenic-induced health effects.¹³ Experimental animals with a low dietary intake of methionine, choline, and protein were found to have lowered methylation of inorganic arsenic.²²

Nutrition surveys in eight states of India conducted by the National Nutrition Bureau of India revealed calorie consumption less than the RDA of energy (male—2875 and female—2225 kcal/day).²⁹ Studies in Murshidabad district of West Bengal revealed that in more than 50% of the household surveyed families with poor nutrition suffer more from arsenic toxicity.²⁷ Other studies also revealed that undernourishment was found to increase the risk of skin lesions and skin cancer in arsenic-exposed populations.^{9,11,15} In Western States like Alaska, studies revealed that populations consuming high concentrations of arsenic from their drinking water often did not show arsenical skin lesions. The reason may be their nutritional status.³⁰ Experimental research in animals has also found that low protein and amino acids in diets increase risks of arsenic related health effects.^{12,22,31} The study conducted among residents of California where arsenic level in the drinking water supplies had been near 100 µg/l suggests that low intakes of dietary protein, iron, zinc and niacin lead to decreased production of DMA and increased level of MMA in arsenic exposed individuals.³² Studies showed that consumption of high levels of niacin (vitamin B3) was associated with arsenic methylation.³³ There are studies indicating that consumption of a diet rich in riboflavin, pyridoxin, vitamin A, C, and E can significantly reduce the harmful effects of developments of skin lesions.²⁶ Other nutrients like niacin, iron, calcium, protein, and thiamin were also reported to be protective against arsenic toxicity.³⁴ Inadequate intake of folate, methionine, cysteine, vitamin B6 and B12, calories and proteins are associated with arsenic related health effects in humans.^{12,22,34-36} It was experimentally proved that low zinc concentrations were found in blood and urine of arsenic effected patients.³⁷

A cross-sectional study was conducted in Chakdah and Haringhata block of Nadia district of West Bengal to determine the dietary intake of nutrients and its correlation with manifestations of arsenicosis.

The study revealed that low socioeconomic status along with less dietary intake of calorie, protein and micronutrients may have a definite role in increasing the risk of development arsenicosis. It was found that there was significant difference in intake of protein along with micronutrients like thiamine, riboflavin, niacin, zinc, and choline in the diet of cases compared to exposed controls in both the sexes, indicating the probable role of above nutrients in development of diseases.³⁸

Based on the above findings all patients living in arsenic endemic area should be advised to consume a variety of green leafy vegetables, locally available pulses and fruits rich in antioxidant and other micronutrients along with low-cost animal protein like egg, fish to decrease the susceptibility from arsenic-related health effects.

REFERENCES

- WHO. Arsenic and Arsenic Compounds (Environmental Health Criteria 224). Geneva: International Programme on Chemical Safety; 2001.
- Thornton I and Farago M. The geochemistry of arsenic. In: Abernathy CO, Calderon RL, Chappell WR, eds. Arsenic Exposure and Health Effects. London, UK: Chapman & Hall; 1997. p. 1–16.
- Alam MGM, Snow ET and Tanaka A. Arsenic and heavy metal contamination of rice, pulses and vegetables grown in Samta village, Bangladesh. In: Chappell WR, Abernathy CO, Calderon RL, Thomas DJ, eds. Arsenic Exposure and Health Effects V, vol. 8. UK: Elsevier BV; 2003. p. 103–114.
- NRC (National Research Council). Arsenic in Drinking Water. Washington, DC: National Academic Press; 1999.
- WHO Technical Publication. In: Caussy D, ed. A Field Guide for Detection, Management and Surveillance of Arsenicosis Cases, vol 30. New Delhi: WHO Regional Office for South East Asia; 2005. p. 19–22.
- Guha Mazumder DN. Criteria for case definition of arsenicosis. In: Chappell WR, Abernathy CO, Calderon RL, Thomas DJ, eds. Arsenic Exposure and Health Effects V, vol. 9. UK: Elsevier BV; 2003. p. 117–133.
- Guha Mazumder DN. Clinical aspects of chronic arsenic toxicity. J Assoc Phys India 2001;49:650–655.
- Saha KC. Saha's grading of arsenicosis progression and treatment. In: Chappell WR, Abernathy CO, Calderon RL, Thomas DJ, eds. Arsenic Exposure and Health Effects V, vol. 30. UK: Elsevier BV; 2003. p. 391–414.
- Guha Mazumder DN, Chakraborty AK, Ghose A Gupta JD, Chakraborty DP, Dey SB, et al. Chronic arsenic toxicity from drinking tube well water in rural West Bengal. Bull. Wild Health Org. 1998;66:499–506.
- Zaldivar R. Arsenic contamination of drinking water and foodstuffs causing endemic chronic poisoning. Beitr Pathol 1978;151:384–400.
- Hsueh YM, Cheng GS, Wu MM, Yu HS, Kuo TL and Chen CJ. Multiple risk factors associated with arsenic induced skin cancer: effects of chronic liver disease and malnutritional status. Br J Cancer 1995;71:109–114.
- Chen C-J, Kuo TL and WU MM. Arsenic and cancers (letter). Lancet 1988;331:414–415.
- Chung JS, Haque R, Guha Mazumder DN, Moore LE, Ghosh N, Samanta S, et al. Blood concentrations of methionine, selenium, beta-carotene and other micronutrients in a case control study of arsenic induced skin lesions in West Bengal, India. Environ Res 2006;101:230–237.
- Pal A, Chowdhury U, Mandal D and Nayak B. Arsenic burden from cooked rice in the population of arsenic affected and nonaffected areas and Kolkata city in West Bengal, India. Environ Sci Technol 2009;43:3349–3355.
- Guha Mazumder DN, Haque R, Ghosh N, De BK, Santra A, Chakraborty D, et al. Arsenic levels in drinking water and the prevalence of skin lesions in West Bengal, India. Int J Epidemiol 1998;27:871–877.
- Johnson JL and Rajagopalan KV. The interaction of arsenic with the molybdenum center of chicken liver xanthine dehydrogenase. Bioinorg Chem 1978;8:439–444.
- Calabrese EJ. Nutrition and Environmental Health. New York: John Wiley; 1980.
- Harding-Barlow I. In: Lederer WH, Fensterheim RJ, eds. Arsenic: Industrial, Biomedical, Environmental Perspective. New York: Van Nostrand Reinhold; 1983. p. 203.
- US EPA. Special Report on Ingested Inorganic Arsenic. Skin Cancer, Nutritional Essentially (EPA/625/3-87/013). Washington, DC: Environmental Protection Agency; 1988.
- Vahter M. Genetic polymorphism in the biotransformation of inorganic arsenic and its role in toxicity. Toxicol Lett 2000;112–113:209–217.
- Maity S and Chatterjee AK. Differential response of cellular antioxidant mechanism of liver and kidney to arsenic exposure and its relation to dietary protein deficiency. Environ Pharmacol Toxicol 2000;8:227–235.
- Vahter M and Marafante E. Effects of low dietary intake of methionine, choline or proteins on the biotransformation of arsenite in the rabbit. Toxicol Lett 1987;37:41–46.
- Steinmaus C, Carrigan K, Kalman D, Atallah R, Yuan Y and Smith AH. Dietary intake and arsenic methylation in a U.S. population. Environ Health Perspect 2005;113:1153–1159.
- Vahter M. Methylation of inorganic arsenic in different mammalian species and population groups. Sci Prog 1999;82(Part 1):69–88.
- Heck JE, Nieves JW, Chen Y, Parvez F, Brandt-Rauf PW, Graziano JH, et al. Dietary intake of methionine, cysteine and protein and urinary arsenic excretion in Bangladesh. Environ Health Perspect 2009;117:99–104.
- Zablotska LB, Chen Y, Graziano JH, Parvez F, Geen AV, Howe GR, et al. Protective effects of B vitamins antioxidants on the risk of arsenic related skin lesions in Bangladesh. Environ Health Perspect 2008;116:1056–1062.
- Roychowdhury T, Uchino T, Tokunaga H and Ando M. Survey of arsenic in food composites from an arsenic-affected area of West Bengal, India. Food Chem Toxicol 2002;40:1611–1621.
- Mitra SR, Guha Mazumder DN, Basu A, Blosh G, Haque R, Samanta S, et al. Nutritional factors and susceptibility to arsenic caused skin lesions in West Bengal, India. Environ Med 2004;112:1104–1109.
- Bamji MS. Vitamin deficiencies in rice-eating populations: effects of B-vitamin supplements. Experientia Suppl 1983;44:245–263.
- Harrington JM, Middaugh JP, Morse DL and Houseworth J. A survey of a population exposed to high concentrations of arsenic in well water in Fairbanks, Alaska. Am J Epidemiol 1978;108:337–385.
- Maity S and Chatterjee AK. Effects on levels on glutathione and some related enzymes in tissues after an acute arsenic exposure in rats and their relationship to dietary protein deficiency. Arch Toxicol 2001;75:531–537.
- Yang YH and Blackwell RQ. Nutritional and environmental conditions in the endemic black foot area. Formos Sci 1961;15:101–129.
- Steinmaus CM, Yuan Y and Smith AH. The temporal stability of arsenic concentrations in well water in western Nevada. Environ Res 2005;99:164–168.
- Heck JE, Gamble MV, Chen Y, Graziano JH, Slavonic V, Parvez F, et al. Consumption of folate-related nutrients and metabolism of arsenic in Bangladesh. Am J Clin Nutr 2007;85:1367–1374.
- Hoffman DJ, Sanderson CJ, LeCaptain LJ, Cromartie E and Pendleton GW. Interactive effects of arsenate, selenium and dietary protein on survival, growth and physiology in mallard ducklings. Arch Environ Contam Toxicol 1992;22:55–62.
- Lammon CA and Hood RD. Effects of protein deficient diets on the developmental toxicity of inorganic arsenic in mice. Birth Defects Res B: Dev Rep Toxicol 2004;71:124–134.
- Kreppel H, Liu J, Liu Y, Reichl FX and Klassen CD. Zinc induced arsenic tolerance in mice. Fundam Appl Toxicol 1994;23:32–37.
- Deb Debasree, Majumdar KK and Guha Mazumder DN. Arsenicosis and dietary intake among men and women. Proc Natl Acad Sci, India Sect B: Biol Sci 2013;83:405–413.

Address for Correspondence:

Dr Kunal Kanti Majumdar,
Professor, Department of Community Medicine,
KPC Medical College & Hospital, Jadavpur, Kolkata, India.
E-mail: kunalmajumdar1@gmail.com, **Phone:** +91 9830401906.

How to cite this article: Majumdar KK. Arsenicosis and malnutrition. IntJMRP 2016;1(1):16-17