Tables

Table 1: Quality-control flow (main analysis).

Step	Trials	Sessions	Athletes
All male football CMJ trials	833	333	215
R1: valid body mass	828	329	212
R2: quiet-stance mass within 5% of body mass	813	328	212
R3: no force on plate during flight	811	327	212
R4: complete metrics and age	811	327	212
R5: end of landing (lowest COM) found	805	327	212

Note. Rules R1–R4 were fixed before any result was examined; R5 applies only to the main landing definition (touch-down to the lowest point of the centre of mass), and the 1-s landing window therefore retained 811 trials. Sessions, test occasions on which an athlete performed one or more trials. COM, centre of mass.

Table 2: Characteristics of the analysed sample (805 trials, 212 athletes).

Variable	*n*	Mean ± SD	Median (IQR)	Range
Age at first test (years)	212	16.01 ± 5.94	14.10 (12.00 to 16.80)	8.80 to 37.9
Body mass at first test (kg)	212	53.31 ± 15.90	51.73 (40.86 to 64.03)	25.03 to 95
Jump height, flight time (cm)	805	28.19 ± 8.43	26.96 (21.62 to 34.69)	8.54 to 51.4
Concentric peak force (N)	805	1269.26 ± 420.74	1227.01 (923.47 to 1582.99)	473.52 to 3
Concentric peak force (× BW)	805	2.35 ± 0.38	2.29 (2.11 to 2.51)	1.65 to 4.70
Eccentric braking impulse (N s)	805	27.80 ± 15.43	24.60 (16.59 to 35.70)	0.00 to 95.1
Concentric impulse, first 100 ms (N s)	805	62.24 ± 26.66	58.70 (41.76 to 79.14)	1.51 to 187
Peak landing force (× BW)	805	5.39 ± 1.97	4.97 (3.88 to 6.67)	2.19 to 14.8
Contraction time (s)	805	0.78 ± 0.19	0.77 (0.66 to 0.90)	0.31 to 1.48
Countermovement depth (cm)	805	−26.55 ± 9.90	−27.13 (−33.55 to −20.71)	−52.39 to 0
Touch-down to lowest COM (s)	805	0.25 ± 0.11	0.24 (0.15 to 0.31)	0.07 to 0.93
Trials per athlete	212	3.80 ± 2.28	3.00 (2.00 to 5.00)	1.00 to 16.0
Trials per session	327	2.46 ± 0.78	2.00 (2.00 to 3.00)	1.00 to 7.00
Sessions per athlete	212	1.54 ± 0.80	1.00 (1.00 to 2.00)	1.00 to 6.00

Note. Age and body mass are per athlete at the first analysed trial; all other rows are per trial unless stated. Jump height, from flight time. Countermovement depth is the vendor estimate of centre-of-mass displacement (negative downwards). Touch-down to lowest COM, duration of the landing phase in the main analysis. BW, body weight; COM, centre of mass; IQR, interquartile range.

Table 3: Retained principal components of the concatenated take-off and landing curve (main analysis).

	Variance	Cumulative	Landing share	Largest	Pearson *r* with				
Component	(%)	(%)	of loading (%)	loading at	LPF	JH	CT	CMD	LD
PC1	37.3	37.3	99.7	Landing 40%	0.39	0.13	−0.04	0.01	−0.75

PC2	16.0	53.3	99.9	Landing 26%	0.22	0.13	0.02	0.00	−0.14
PC3	12.6	65.9	99.5	Landing 16%	0.39	0.34	0.00	−0.14	−0.32
PC4	8.9	74.8	99.3	Landing 9%	0.45	0.24	0.15	−0.18	−0.04

Note. Landing share of loading, percentage of the squared loading vector that falls in the landing block (the remainder falls in the take-off block). Largest loading at, phase and percentage of the phase at which the loading has its largest absolute value. LPF, peak landing force (× BW); JH, jump height; CT, contraction time; CMD, countermovement depth; LD, landing duration (touch-down to lowest COM). Signs are fixed so that the largest absolute loading is positive.

Table 4: Dispersion of component scores in the full sample and among trials matched on jump height or on relative concentric peak force (main analysis).

	SD, full	Jump-height band		Relative peak-force band	
Component	sample	SD	Ratio (95% CI)	SD	Ratio (95% CI)
PC1	7.28	6.45	0.89 (0.70–1.06)	7.33	1.01 (0.77–1.16)
PC2	4.77	4.62	0.97 (0.77–1.16)	5.01	1.05 (0.82–1.24)
PC3	4.22	4.03	0.95 (0.74–1.13)	4.25	1.00 (0.81–1.17)
PC4	3.56	4.16	1.17 (0.81–1.44)	3.75	1.05 (0.87–1.23)
Whole curve[a]	11.92	11.99	1.01	11.95	1.00

Note. Matched bands comprise trials between the 45th and 55th percentiles of jump height (26.0–28.2 cm; 82 trials, 51 athletes) or of concentric peak force relative to body weight (2.25–2.32 BW; 81 trials, 58 athletes). Ratio, SD in the band divided by SD in the full sample; 95% CI from 2,000 athlete-level bootstrap resamples. Component SDs depend on the number of grid points and are comparable only within this table; ratios are scale-free. [a]Square root of the summed point-wise variance of the 202-point curve.

Table 5: Variance of each component score explained by the four vendor metrics, age and body mass, and decomposition of component-score variance (main analysis).

	R^2 of component score on metrics, age and body mass (%)					Variance of component score (%)		
Component	In-sample	Adjusted	Athlete CV	One trial per athlete	Athlete means	Between athletes	Between sessions	Within session
PC1	7.7	7.0	4.0	10.0	11.1	29.6	34.9	35.5
PC2	2.7	2.0	0.3	4.3	5.4	21.4	17.0	61.6
PC3	13.1	12.4	11.8	17.8	21.2	22.9	2.3	74.8
PC4	9.1	8.4	6.3	11.1	13.8	10.0	20.3	69.6

Note. Predictors: jump height, concentric peak force, eccentric braking impulse, concentric impulse over the first 100 ms, age and body mass (ordinary least squares). In-sample and adjusted, all 805 trials. Athlete CV, out-of-sample R^2 from five-fold cross-validation with athletes, not trials, allocated to folds. One trial per athlete, one randomly drawn trial per athlete (212 trials). Athlete means, scores and predictors averaged within athlete. Variance shares from a three-level random-intercept model (athlete, session within athlete, trial within session) fitted by restricted maximum likelihood.

Table 6: Main analysis compared with the sensitivity analysis in which the landing phase ended 1 s after touch-down.

Quantity	Main: landing to lowest COM	Sensitivity: 1-s landing
Trials / athletes after QC	805 / 212	811 / 212
Variance explained, PC1–PC4 (%)	37.3, 16.0, 12.6, 8.9	20.2, 16.6, 15.1, 10.2
Cumulative, PC1–PC4 (%)	74.8	62.2
Landing share of loading, PC1–PC4 (%)	99.7, 99.9, 99.5, 99.3	96.4, 92.5, 3.7, 80.2
PC1 SD ratio, jump-height band (95% CI)	0.89 (0.70–1.06)	0.82 (0.69–0.96)
Whole-curve SD ratio, jump-height band	1.01	1.01
In-sample R^2, PC1–PC4 (%)	7.7, 2.7, 13.1, 9.1	19.1, 5.3, 68.3, 17.8
Athlete-CV R^2, PC1–PC4 (%)	4.0, 0.3, 11.8, 6.3	13.4, 2.8, 66.4, 15.1
Between-athlete variance, PC1–PC4 (%)	29.6, 21.4, 22.9, 10.0	45.3, 18.7, 26.5, 31.6
r(PC1, landing peak force)	0.39	0.71
r(PC3, contraction time)	0.00	–0.79

Note. COM, centre of mass; CI, confidence interval; CV, cross-validation. Between-athlete variance from the three-level random-intercept model. In the 1-s window PC3 is a take-off component (Figure 6), which explains its correlation with contraction time.

Figures




Figure 1: Analysis pipeline. Steps 1–4 build one 202-point, body-weight-normalised force–time curve per trial; steps 5–7 address the three aims; step 8 examines the structure of the repeated measurements and the sensitivity of the conclusions to analytical choices. All decision rules were fixed before the results were examined. CMJ, countermovement jump; PCA, principal component analysis; OLS, ordinary least squares; CV, cross-validation; CI, confidence interval.




Figure 2: (A) Mean body-weight-normalised vertical force (line) ± 1 SD (shaded) of the 805 analysed countermovement jumps. The take-off phase (movement onset to take-off) and the landing phase (touch-down to the lowest point of the centre of mass, COM) were each time-normalised to 0–100% and concatenated; flight is omitted. The dotted line marks body weight. (B) Variance explained by each of the first ten principal components (bars) and cumulatively (line); the dashed line marks the four retained components.




Figure 3: Interpretation of the four retained principal components (main analysis). Each panel shows the mean curve (dashed) and the mean curve plus (vermillion) or minus (blue) two standard deviations of the component score. Values below zero in the reconstructions are an artefact of adding a linear component to the mean and do not correspond to observed trials. The vertical line separates the take-off and landing phases.




Figure 4: Trials matched on jump height do not share a common force–time curve. (A) Curves of the 82 trials between the 45th and 55th percentiles of jump height (27.1 ± 0.65 cm), coloured by their PC1 score. (B) Ratio of the standard deviation of each component score among trials matched on jump height (circles) or on relative concentric peak force (triangles) to that in the full sample, with athlete-level bootstrap 95% confidence intervals. A ratio of 1 indicates that matching on the metric leaves curve-shape variation unchanged.




Figure 5: Proportion of the variance of each component score explained by the four vendor metrics, age and body mass, under four evaluation schemes, for the main landing definition (left) and the 1-s landing window (right). CV, cross-validation with athletes allocated to folds.




Figure 6: Sensitivity analysis with the landing phase ending 1 s after touch-down: mean curve (dashed) ± two standard deviations of each retained component (vermillion, +2 SD; blue, −2 SD). Under this definition PC3 is located almost entirely in the take-off phase (96.3% of its squared loading) and describes the duration and amplitude of the propulsive force.