

# Constructive Algorithms for Discrepancy Minimization

(Nikhil Bansal, FOCS 2010)

Pravesh Kothari

The University of Texas at Austin

`kothari@cs.utexas.edu`

January 30, 2012

# Discrepancy

- “*Approximating a probability space by a smaller probability space*”

# Discrepancy

- “*Approximating a probability space by a smaller probability space*”
- Example: Approximating a Continuous Probability Space by A Discrete Space
- Question: How **uniformly** can you distribute  $n$ -points in a unit square?
- What does **UNIFORM** mean?

# Axis Parallel Rectangles in a Grid

- Question: How **uniformly** can you distribute  $n$ -points in a unit square?
- What does **UNIFORM** mean?




Figure: Distributing Points in a Unit Square

For any axis parallel rectangle  $R$  inside the unit square,

$$|Prob_{x \in [0,1]^2}[x \in R] - Prob_{y \in S}[y \in R]|$$

is small.

- $|(\# \text{ of points from } S \text{ in } R) - n \cdot \text{Area}(R)|$  is small.

## Discrepancy: Example 2

- “Approximating a large discrete space by a smaller discrete space.”
- **Given:** A set  $X$  of  $n$  points in a unit square.
- **Problem:** Color each point in  $X$  red/blue such that each *axis parallel rectangle* has “roughly” the same number of red or blue points.
- **Discrepancy:** max over Rectangle  $R$

$$((\# \text{ of red points in } R) - (\# \text{ of blue points in } R))$$

## Discrepancy: Example 2

- “Approximating a large discrete space by a smaller discrete space.”
- **Given:** A set  $X$  of  $n$  points in a unit square.
- **Problem:** Color each point in  $X$  red/blue such that each *axis parallel rectangle* has “roughly” the same number of red or blue points.
- **Discrepancy:** max over Rectangle  $R$

$$((\# \text{ of red points in } R) - (\# \text{ of blue points in } R))$$

- **Non Trivial:** Random Coloring:  $O(\sqrt{n \log n})$   
**Best:**

## Discrepancy: Example 2

- “Approximating a large discrete space by a smaller discrete space.”
- **Given:** A set  $X$  of  $n$  points in a unit square.
- **Problem:** Color each point in  $X$  **red/blue** such that each *axis parallel rectangle* has “roughly” the same number of **red** or **blue** points.
- **Discrepancy:** max over Rectangle  $R$

$$((\# \text{ of red points in } R) - (\# \text{ of blue points in } R))$$

- **Non Trivial:** Random Coloring:  $O(\sqrt{n \log n})$   
**Best:**  $O(\log^{2.5} n)$

# Combinatorial Discrepancy

- **Given :** Set System  $(X, \mathbb{S})$ .  $X = \{1, 2, \dots, n\} = [n]$  and  $\mathbb{S} = \{S_1, S_2, \dots, S_m\}$ . Each  $S_i \subseteq X$ .
- **Problem:** Find a coloring in **red** and **blue** such that for every  $S_i \in \mathbb{S}$ ,

$$|(\# \text{ of } x \text{ colored } \textcolor{red}{red} \text{ in } S_i) - (\# \text{ of } x \text{ colored } \textcolor{blue}{blue} \text{ in } S_i)|$$

is small.

- A huge number of Applications!
- **CS:** Computational Geometry, Comb. Opt. , Monte Carlo Simulation, Machine Learning, Complexity, Communication Complexity, Pseudorandomness...
- **Math (but not so CS) :** Dynamical Systems, Combinatorics, Math. Finance, Number Thy, Ramsey Thy, Algebra, Measure Theory, Numerical Calculus...



# Get to the Business: Discrepancy Minimization

- Given:** A set system  $(X, \mathbb{S})$  **Universe**  $X:[n]$   
**Set of Subsets** of  $X$ ,  $\mathbb{S} = \{S_1, S_2, \dots, S_m\}$
- Problem:** Find  $\chi : X \rightarrow \{-1, 1\}^n$  to  
**minimize**  $|\chi(S)|_\infty = \max_{S_i \in \mathbb{S}} |\sum_{x \in S_i} \chi(x)|$

## Best Known *Algorithm*: Random Coloring

- $T = (X, \mathbb{S})$  : A set system with  $|X| = n$  and  $|\mathbb{S}| = m$ .
- $\chi : X \rightarrow \{-1, 1\}^n$  a random coloring.
- Chernoff Bound:  $\text{Prob}[|\chi(S)| \geq c\sqrt{n}] \leq e^{-\frac{c^2}{4}}$

## Best Known *Algorithm*: Random Coloring

- $T = (X, \mathbb{S})$  : A set system with  $|X| = n$  and  $|\mathbb{S}| = m$ .
- $\chi : X \rightarrow \{-1, 1\}^n$  a random coloring.
- Chernoff Bound:  $\text{Prob}[|\chi(S)| \geq c\sqrt{n}] \leq e^{-\frac{c^2}{4}}$
- Choose  $c = \sqrt{4 \log 4m}$  to obtain

$$\text{Prob}[|\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2m}$$

## Best Known *Algorithm*: Random Coloring

- $T = (X, \mathbb{S})$  : A set system with  $|X| = n$  and  $|\mathbb{S}| = m$ .
- $\chi : X \rightarrow \{-1, 1\}^n$  a random coloring.
- Chernoff Bound:  $\text{Prob}[|\chi(S)| \geq c\sqrt{n}] \leq e^{-\frac{c^2}{4}}$
- Choose  $c = \sqrt{4 \log 4m}$  to obtain

$$\text{Prob}[|\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2m}$$

- Union Bound over all sets:

$$\text{Prob}[\exists S \in \mathbb{S} : |\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2}$$

## Best Known *Algorithm*: Random Coloring

- $T = (X, \mathbb{S})$  : A set system with  $|X| = n$  and  $|\mathbb{S}| = m$ .
- $\chi : X \rightarrow \{-1, 1\}^n$  a random coloring.
- Chernoff Bound:  $\text{Prob}[|\chi(S)| \geq c\sqrt{n}] \leq e^{-\frac{c^2}{4}}$
- Choose  $c = \sqrt{4 \log 4m}$  to obtain

$$\text{Prob}[|\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2m}$$

- Union Bound over all sets:

$$\text{Prob}[\exists S \in \mathbb{S} : |\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2}$$

- Thus with probability at least  $\frac{1}{2}$ ,  $\chi$  achieves a discrepancy of  $\mathbf{O}(\sqrt{n \log m})$ .

## Best Known *Algorithm*: Random Coloring

- $T = (X, \mathbb{S})$  : A set system with  $|X| = n$  and  $|\mathbb{S}| = m$ .
- $\chi : X \rightarrow \{-1, 1\}^n$  a random coloring.
- Chernoff Bound:  $\text{Prob}[|\chi(S)| \geq c\sqrt{n}] \leq e^{-\frac{c^2}{4}}$
- Choose  $c = \sqrt{4 \log 4m}$  to obtain

$$\text{Prob}[|\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2m}$$

- Union Bound over all sets:

$$\text{Prob}[\exists S \in \mathbb{S} : |\chi(S)| \geq \sqrt{4n \log 4m}] \leq \frac{1}{2}$$

- Thus with probability at least  $\frac{1}{2}$ ,  $\chi$  achieves a discrepancy of  $\mathbf{O}(\sqrt{n \log m})$ .
- Analysis is **tight** !

# Surprise! Better Colorings *Exist!*

- Spencer (1985) : **“Six Standard Deviations Suffice”**  
*There always exists a coloring with discrepancy at most  $6\sqrt{n}$ .*  
(Here,  $m = n$ )

# Surprise! Better Colorings *Exist!*

- Spencer (1985) : **“Six Standard Deviations Suffice”**  
*There always exists a coloring with discrepancy at most  $6\sqrt{n}$ .*  
(Here,  $m = n$ )
- For arbitrary  $m$ , discrepancy =  $O\left(\sqrt{n \log \frac{m}{n}}\right)$



# Surprise! Better Colorings *Exist!*

- Spencer (1985) : “**Six Standard Deviations Suffice**”  
*There always exists a coloring with discrepancy at most  $6\sqrt{n}$ .*  
(Here,  $m = n$ )
- For arbitrary  $m$ , discrepancy =  $O(\sqrt{n \log \frac{m}{n}})$
- **Problem** : “Highly” non constructive Proof. (PHP on exponentially large universe)
- **Challenge**: Can we find such a coloring **algorithmically** ?  
[ Spencer 1985 ]

# Surprise! Better Colorings *Exist!*

- Spencer (1985) : “**Six Standard Deviations Suffice**”  
*There always exists a coloring with discrepancy at most  $6\sqrt{n}$ .*  
 (Here,  $m = n$ )
- For arbitrary  $m$ , discrepancy =  $O(\sqrt{n \log \frac{m}{n}})$
- **Problem** : “Highly” non constructive Proof. (PHP on exponentially large universe)
- **Challenge:** Can we find such a coloring **algorithmically** ?  
 [ Spencer 1985 ]
- **Conjecture:** Not Possible! [Alon and Spencer]

# Approximating Discrepancy

- **Question** : Suppose a set system has low discrepancy (much lower than  $\sqrt{n}$  ). Is it possible to find it?
- **Answer:**

# Approximating Discrepancy

- **Question** : Suppose a set system has low discrepancy (much lower than  $\sqrt{n}$  ). Is it possible to find it?
- **Answer:** **NO!** [Charikar-Newman-Nikolov 2011] :  
Distinguishing between discrepancy of 0 and  $\sqrt{n}$  is NP-Complete

# Approximating Discrepancy

- **Question** : Suppose a set system has low discrepancy (much lower than  $\sqrt{n}$ ). Is it possible to find it?
- **Answer:** **NO!** [Charikar-Newman-Nikolov 2011] : Distinguishing between discrepancy of 0 and  $\sqrt{n}$  is NP-Complete
- What if a set system has low *hereditary discrepancy* ? [Matousek]

- 

$$\text{Herdisc}(X, \mathbb{S}) = \max_{Y \subseteq X} \text{disc}(Y, \mathbb{S}_Y)$$

- Why Hereditary Discrepancy ?  
Well studied, more robust notion of discrepancy, much used in Computational Geometry PLUS  
*fits naturally into the technique!*

# The Results

## Theorem (Constructive Spencer)

*Given a set system  $(X, \mathbb{S})$  with  $|X| = |\mathbb{S}| = n$ , there exists a  $\text{poly}(m, n)$  algorithm to construct a coloring  $\chi$  with discrepancy  $O(\sqrt{n})$ .*

## Theorem ( Constructive Low Hereditary Discrepancy )

*Given a set system  $(X, \mathbb{S})$  with  $|X| = n$ ,  $|\mathbb{S}| = m$ , and a guarantee that  $\text{herdisc}(X, \mathbb{S}) \leq \lambda$ , there exists an algorithm running in time  $\text{poly}(m, n)$  to construct a coloring with discrepancy  $O(\lambda \log(mn))$ .*

- A neat technique! May have applications waiting to be discovered in making several other existential results algorithmic.

# A first Attempt: Convex Relaxations

## Theorem ( Constructive Low Hereditary Discrepancy )

*Given a set system  $(X, \mathbb{S})$  with  $|X| = n$ ,  $|\mathbb{S}| = m$ , and a guarantee that  $\text{herdisc}(X, \mathbb{S}) \leq \lambda$ , there exists an algorithm running in time  $\text{poly}(m, n)$  to construct a coloring with discrepancy  $O(\lambda \log(mn))$ .*

# A first Attempt: Convex Relaxations

## Theorem ( Constructive Low Hereditary Discrepancy )

Given a set system  $(X, \mathbb{S})$  with  $|X| = n$ ,  $|\mathbb{S}| = m$ , and a guarantee that  $\text{herdisc}(X, \mathbb{S}) \leq \lambda$ , there exists an algorithm running in time  $\text{poly}(m, n)$  to construct a coloring with discrepancy  $O(\lambda \log(mn))$ .

- **A first attempt: LP**

$$-\sqrt{n} \leq \sum_{x \in S_i} u_x^+ - \sum_{x \in S_i} u_x^- \leq \sqrt{n} \quad \forall S_i \in \mathbb{S}$$

$$u_x^+ + u_x^- = 1 \quad \forall u \in X$$

$$u_x^+, u_x^- \geq 0 \quad \forall u \in X$$



# A first Attempt: Convex Relaxations

## Theorem ( Constructive Low Hereditary Discrepancy )

Given a set system  $(X, \mathbb{S})$  with  $|X| = n$ ,  $|\mathbb{S}| = m$ , and a guarantee that  $\text{herdisc}(X, \mathbb{S}) \leq \lambda$ , there exists an algorithm running in time  $\text{poly}(m, n)$  to construct a coloring with discrepancy  $O(\lambda \log(mn))$ .

- **A first attempt: LP**

$$-\sqrt{n} \leq \sum_{x \in S_i} u_x^+ - \sum_{x \in S_i} u_x^- \leq \sqrt{n} \quad \forall S_i \in \mathbb{S}$$

$$u_x^+ + u_x^- = 1 \quad \forall u \in X$$

$$u_x^+, u_x^- \geq 0 \quad \forall u \in X$$

- **Problem:** Absolutely No Information!  $u_x^+ = u_x^- = \frac{1}{2}$

# A modified first attempt: SDPs

- SDP :

$$\left\| \sum_{x \in S_i} v_x \right\|^2 \leq n \quad \forall S_i \in \mathbb{S}$$

$$\|v_x\|^2 = 1 \quad \forall x \in X$$

$$v_x \in \mathbb{R}^n \quad \forall x \in X$$

# A modified first attempt: SDPs

- SDP :

$$\left\| \sum_{x \in S_i} v_x \right\|^2 \leq n \quad \forall S_i \in \mathcal{S}$$

$$\|v_x\|^2 = 1 \quad \forall x \in X$$

$$v_x \in \mathbb{R}^n \quad \forall x \in X$$

- **Problem** :  $v_x = e_x \quad \forall x \in X$  always a solution!

# A modified first attempt: SDPs

- SDP :

$$\left\| \sum_{x \in S_i} v_x \right\|^2 \leq n \quad \forall S_i \in \mathcal{S}$$

$$\|v_x\|^2 = 1 \quad \forall x \in X$$


$$v_x \in \mathbb{R}^n \quad \forall x \in X$$

- **Problem :**  $v_x = e_x \quad \forall x \in X$  always a solution!
- **Observation:** Tighten Some Constraints a Little Bit for some sets, and the trivial solution disappears...
- Example :

$$\left\| \sum_{x \in S_i} v_x \right\|^2 \leq \frac{n}{\log n}$$

# The Algorithm

- A random walk over *fractional* colorings of  $X$



- Each move adds an increment generated by a suitable SDP  $\delta_t$  to the current coloring.
- Whenever a label hits  $\{-1, 1\}^n$ , we freeze the label.
- Random Walk over different *directions* is (nicely!) correlated.

# The Algorithm: Analysis

- Random Walk over each direction has a *high* variance.  $\Rightarrow$  converge to  $\{-1, 1\}^n$  fast.
- Walk over Discrepancies of each set has *low* variance  $\Rightarrow$  Discrepancy is bounded!

# The Algorithm: Generating a Step

- Let  $U \subseteq X$  be the set of vertices which haven't hit  $\{-1, 1\}^n$  at the start of a fixed iteration.
- The following SDP is feasible:

$$\left\| \sum_{x \in S_i \cap U} v_x \right\|^2 \leq \lambda \quad \forall S_i \in \mathbb{S}$$

$$\|v_x\|^2 = 1 \quad \forall x \in U$$

- Rounding:** Choose a **random** vector of **independent**  $\mathcal{N}(0, 1)$  variables  $(g_1, g_2, \dots, g_n)$
- For each  $x$ , let the update be:  $\eta_x = \langle v_x, g \rangle$

# Why the rounding works

## Fact (Additivity of Gaussians)

If  $(g_1, g_2, \dots, g_n)$  is a vector of independent gaussians of variance 1, then  $\langle v_x, g \rangle \sim \mathcal{N}(0, \|v_x\|^2)$

- **High Variance Walk on coordinates :**

$$\eta_x = \langle v_x, g \rangle \sim \mathcal{N}(0, 1)$$

- **Low Variance Walk on Discrepancies** For each set  $S_i \in \mathbb{S}$ ,  $\sum_{x \in S_i} \eta_x \sim \mathcal{N}(0, \sigma^2)$  with  $\sigma \leq \lambda$ .



# Algorithm

- Start with the coloring  $\chi_0 = (0, 0, \dots, 0)$
- Update each coloring:  $\chi_t = \chi_{t-1} + \gamma \cdot \eta$   
( $\gamma$  is tiny  $\sim \frac{1}{n^2}$  step size, chosen to control error due to rounding)
- Whenever  $\chi(x) \geq (1 - 1/n)$  or  $\chi(x) \leq (-1 + 1/n)$  round  $\chi(x)$  and fix.
- Thus, color of an element does a random walk with each step  $\gamma \mathcal{N}(0, 1)$   
discrepancy of each set does a random walk with each step  $\gamma \mathcal{N}(0, \lambda^2)$

# Analysis

Claim (Half the vertices are colored in each iteration)

*With probability at least  $\frac{1}{2}$ ,  $\frac{n}{2}$  coordinates are colored in the first iteration.*

# Analysis

Claim (Half the vertices are colored in each iteration)

*With probability at least  $\frac{1}{2}$ ,  $\frac{n}{2}$  coordinates are colored in the first iteration.*

A round  $T = O(\frac{1}{\gamma^2})$  steps.

Claim ( Low Discrepancy Rise)

*Each set has discrepancy  $O(\lambda)$  in expectation in each round.*

# Analysis

Claim (Half the vertices are colored in each iteration)

*With probability at least  $\frac{1}{2}$ ,  $\frac{n}{2}$  coordinates are colored in the first iteration.*

A round  $T = O(\frac{1}{\gamma^2})$  steps.

Claim ( Low Discrepancy Rise)

*Each set has discrepancy  $O(\lambda)$  in expectation in each round.*

Claim

*Every element is colored in  $O(\log n)$  rounds.*

# Analysis

Claim (Half the vertices are colored in each iteration)

*With probability at least  $\frac{1}{2}$ ,  $\frac{n}{2}$  coordinates are colored in the first iteration.*

A round  $T = O(\frac{1}{\gamma^2})$  steps.

Claim ( Low Discrepancy Rise)

*Each set has discrepancy  $O(\lambda)$  in expectation in each round.*

Claim

*Every element is colored in  $O(\log n)$  rounds.*

Claim

*Expected discrepancy for any set in all rounds is at most  $O(\lambda \log n)$*

# Analysis

Claim (Half the vertices are colored in each iteration)

*With probability at least  $\frac{1}{2}$ ,  $\frac{n}{2}$  coordinates are colored in the first iteration.*

A round  $T = O(\frac{1}{\gamma^2})$  steps.

Claim ( Low Discrepancy Rise)

*Each set has discrepancy  $O(\lambda)$  in expectation in each round.*

Claim

*Every element is colored in  $O(\log n)$  rounds.*

Claim

*Expected discrepancy for any set in all rounds is at most  $O(\lambda \log n)$*

# Summary

- Each Step generated by some **SDP** which introduces **correlation** between updates on colors of different elements.
- Use Hereditary Discrepancy to guarantee **feasibility**.
- **High** Variance of the Walk along each coordinate.
- **Low** Variance walk on the set of discrepancies.

- Spencer's Setting:  $(X, \mathbb{S})$ ,  $|X| = n$  and  $|\mathbb{S}| = m$ .
- Suppose we use  $\lambda = \sqrt{n}$  we obtain : discrepancy of  $O(\sqrt{n}\sqrt{\log n \log m})$  .



- Spencer's Setting:  $(X, \mathbb{S})$ ,  $|X| = n$  and  $|\mathbb{S}| = m$ .
- Suppose we use  $\lambda = \sqrt{n}$  we obtain : discrepancy of  $O(\sqrt{n}\sqrt{\log n \log m})$  .
- **Idea!** : Tighten bounds on constraints in successive SDPs by using Spencer's bound itself!

- Spencer's Setting:  $(X, \mathbb{S})$ ,  $|X| = n$  and  $|\mathbb{S}| = m$ .
- Suppose we use  $\lambda = \sqrt{n}$  we obtain : discrepancy of  $O(\sqrt{n}\sqrt{\log n \log m})$  .
- **Idea!** : Tighten bounds on constraints in successive SDPs by using Spencer's bound itself!
- Nope! :-( Doesn't work. Gives  $O(\sqrt{n \log m})$
- **Issue:** The notorious Union Bound.

# Cool Idea: SDP from Spencer's Proof

**Want:**

- 1) Tighter Constraints on SDPs to guarantee lower discrepancy.
- 2) Forbid the nasty union bound.

# Open Questions

## Problem ( Beck Fiala Conjecture )

*Given a set system  $(X, \mathbb{S})$  such that each  $x \in X$  appears in at most  $t$  sets in  $\mathbb{S}$ , construct a coloring of  $X$  with minimum discrepancy.*

- History: Discrepancy  $2t - 1$  [Beck and Fiala, 1981],  $O(C\sqrt{t \log m \log n})$  [Beck, 1981],  $2t - 3$  by [Bednarchak and Helm, 1997 and Helm, 1999]
- $O(\sqrt{t \log n})$  [Banaszczyk, 1998], and a constructive version by [Srinivasan, ]
- Beck and Fiala conjectured a discrepancy of  $O(\sqrt{t})$  –  
WIDELY OPEN!

# Thank You!

## QUESTIONS?