The same controls, a different history

Recorded Qwen loops · every fraction, dwell and starting history



$P_f = H + f(E_b - H)$; the stated dwell applies to all four visited stages.

Each direction starts separately from H0 and H1 at reference phase zero.

Direction / original lane	P_f periods out / return	E_b period	Final H bit
f = 0.50 · 240 updates			
E0 / H0	6 / 6	6	**0**
E0 / H1	6 / 6	6	**0**
E1 / H0	6 / U	U	**1**
E1 / H1	U / U	U	**1**
f = 0.50 · 480 updates			
E0 / H0	6 / 6	6	**0**
E0 / H1	6 / 6	6	**0**
E1 / H0	6 / 6	U	**1**
E1 / H1	6 / 6	U	**1**
f = 0.25 · 480 updates			
E0 / H0	6 / 6	6	**0**
E0 / H1	6 / 6	6	**0**
E1 / H0	6 / 6	U	**1**
E1 / H1	6 / 6	U	**1**
f = 0.75 · 480 updates			
E0 / H0	6 / 6	6	**0**
E0 / H1	6 / 6	6	**0**
E1 / H0	6 / 6	U	**1**
E1 / H1	U / 6	U	**1**



6: qualified primitive period six. U: no qualified period from 1 to 64.

A gap compares one lane across two visits, never H0 against H1.

Finite-dwell history dependence; branch loss and long-term stability remain unproved.