



हैदराबाद विश्वविद्यालय  
University of Hyderabad

# My Progress

MonoHiggs to  $b\bar{b}$

---

Prayag Yadav

Last updated: 2023-10-20 14:59:45Z

University of Hyderabad

# Table of contents

1. Thu, 5<sup>th</sup> October 2023

Basic kinematic plots (Without any scale factors or corrections)

2. Thu, 26<sup>th</sup> October 2023


MET Filters / MET Flags

3. 2023-10-20

Thu, 5<sup>th</sup> October 2023


## Basic kinematic plots

---




- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**MonoHTobb\_ZpBaryonic**
- Lots of bjets in Signal  
MC

**Figure 1:** BTag score for signal MC sample



- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**Run2018A/MET**
- Less number of bjets in  
Data


**Figure 2:** BTag score for Data samples



- Basic selections :  
 $p_t > 25\text{GeV}$  and  
 $|\eta| < 2.5$
- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**MonoHTobb\_ZpBaryonic**
- Medium Weight  
Parameter used for  
ak4bjets : 0.3040


**Figure 3:** Jet  $p_t$  of signal MC samples

# Jet $p_t$ : Data



- Basic selections :  
 $p_t > 25\text{GeV}$  and  
 $|\eta| < 2.5$
- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**Run2018A/MET**
- Medium Weight  
Parameter used for  
ak4bjets : 0.3040
- Not as predictable as  
signal MC

**Figure 4:** Jet  $p_t$  of Data samples




- Basic selections :  
 $p_t > 25\text{GeV}$  and  
 $|\eta| < 2.5$  for each jet
- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**MonoHTobb\_ZpBaryonic**
- Medium Weight  
Parameter used for  
ak4bjets selection :  
0.3040
- Peaks around SM Higgs  
mass

Figure 5: DiJet mass of signal MC samples




# Dijet mass : Data




- Basic selections :  
 $p_t > 25\text{GeV}$  and  $|\eta| < 2.5$  for each jet
- Btagger used :  
**btagDeepFlavB**
- Sample used:  
**Run2018A/MET**
- Medium Weight  
Parameter used for  
ak4bjets selection :  
0.3040
- Lot of noise, no clear  
structure

**Figure 6:** Dijet mass of Data samples



- No filters or Trigger applied

**Figure 7:** MET  $p_t$  for signal MC samples




- No filters or Trigger applied
- Looks similar to the Jet data

**Figure 8:** MET  $p_t$  for Data samples

## MET Filters


---



- Compared how the MET  $p_t$  looks with and without MET triggers on Data

• .


Figure 9: MET  $p_t$  for MET2018A



- Compared how the MET  $\phi$  looks with and without MET triggers
- .jf


Figure 10: MET  $\phi$  for MET2018A

# MET $p_t$ : MonoHtobb\_ZpBaryonic



- Compared how the MET  $p_t$  looks with and without MET triggers on Signal MC
- .jf

Figure 11: MET  $p_t$  for MonoHtobb\_ZpBaryonic



- Compared how the MET  $\phi$  looks with and without MET triggers on Signal MC
- .

Figure 12: MET  $\phi$  for MC



## Section label

---



- point 1
- point 2

Figure 13: Your caption

