



हैदराबाद विश्वविद्यालय
University of Hyderabad

My Progress

MonoHiggs to $b\bar{b}$

Prayag Yadav

Last updated: 2024-01-04 20:56:38+05:30

University of Hyderabad

Table of contents

1. Thu, 5th October 2023

Basic kinematic plots (Without any scale factors or corrections)


2. Thu, 26th October 2023

MET Filters / MET Flags

3. Tue, 2nd January 2024


Contribution of various backgrounds: Resolved; 2018

Basic kinematic plots




- Btagger used :
btagDeepFlavB
- Sample used:
MonoHTobb_ZpBaryonic
- Lots of bjets in Signal
MC

Figure 1: BTag score for signal MC sample




- Btagger used :
btagDeepFlavB
- Sample used:
Run2018A/MET
- Less number of bjets in
Data

Figure 2: BTag score for Data samples




- Basic selections :
 $p_t > 25\text{GeV}$ and
 $|\eta| < 2.5$
- Btagger used :
btagDeepFlavB
- Sample used:
MonoHTobb_ZpBaryonic
- Medium Weight
Parameter used for
ak4bjets : 0.3040

Figure 3: Jet p_t of signal MC samples



- Basic selections :
 $p_t > 25\text{GeV}$ and
 $|\eta| < 2.5$
- Btagger used :
btagDeepFlavB
- Sample used:
Run2018A/MET
- Medium Weight
Parameter used for
ak4bjets : 0.3040
- Not as predictable as
signal MC


Figure 4: Jet p_t of Data samples



- Basic selections :
 $p_t > 25\text{GeV}$ and
 $|\eta| < 2.5$ for each jet
- Btagger used :
btagDeepFlavB
- Sample used:
MonoHTobb_ZpBaryonic
- Medium Weight
Parameter used for
ak4bjets selection :
0.3040
- Peaks around SM Higgs
mass


Figure 5: DiJet mass of signal MC samples

Dijet mass : Data




- Basic selections :
 $p_t > 25\text{GeV}$ and $|\eta| < 2.5$ for each jet
- Btagger used :
btagDeepFlavB
- Sample used:
Run2018A/MET
- Medium Weight
Parameter used for
ak4bjets selection :
0.3040
- Lot of noise, no clear
structure

Figure 6: Dijet mass of Data samples



- No filters or Trigger applied


Figure 7: MET p_t for signal MC samples



- No filters or Trigger applied
- Looks similar to the Jet data


Figure 8: MET p_t for Data samples

MET Filters



- Compared how the MET p_t looks with and without MET triggers on Data


Figure 9: MET p_t for MET2018A



- Compared how the MET ϕ looks with and without MET triggers
- .jf


Figure 10: MET ϕ for MET2018A

MET p_t : MonoHtobb_ZpBaryonic



- Compared how the MET p_t looks with and without MET triggers on Signal MC
- .jf


Figure 11: MET p_t for MonoHtobb_ZpBaryonic



- Compared how the MET ϕ looks with and without MET triggers on Signal MC
- .

Figure 12: MET ϕ for MC

Contribution of various
backgrounds:Resolved;2018




Selections Applied: Event Selections:

- $MET > 200$ GeV
- no leptons
- no photons

Object Selections:

- jet $p_t > 30$ GeV
- jet $|\eta| < 2.5$
- $\Delta\phi(Jet, MET)$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50$ GeV
- subleading bjet $p_t > 30$ GeV
- atmost 2 additional jets
- dijet = leading bjet + subleading bjet
- dijet mass between (100 GeV;150 GeV)
- dijet $p_t > 100$ GeV

Figure 13: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018




Selections Applied: Event Selections:

- $MET > 200$ GeV
- no leptons
- no photons

Object Selections:

- jet $p_t > 30$ GeV
- jet $|\eta| < 2.5$
- $\Delta\phi(Jet, MET)$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50$ GeV
- subleading bjet $p_t > 30$ GeV
- atmost 2 additional jets
- dijet = leading bjet + subleading bjet
- dijet mass between (100 GeV;150 GeV)
- dijet $p_t > 100$ GeV

Figure 14: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018




Selections Applied: Event Selections:

- $MET > 200$ GeV
- no leptons
- no photons

Object Selections:

- jet $p_t > 30$ GeV
- jet $|\eta| < 2.5$
- $\Delta\phi(Jet, MET)$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50$ GeV
- subleading bjet $p_t > 30$ GeV
- atmost 2 additional jets
- dijet = leading bjet + subleading bjet
- dijet mass between (100 GeV;150 GeV)
- dijet $p_t > 100$ GeV

Figure 15: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018




Selections Applied: Event Selections:

- MET > 200 GeV
- no leptons
- no photons

Object Selections:

- jet $p_t > 30$ GeV
- jet $|\eta| < 2.5$
- $\Delta\phi(\text{Jet}, \text{MET})$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50$ GeV
- subleading bjet $p_t > 30$ GeV
- atmost 2 additional jets
- dijet = leading bjet + subleading bjet
- dijet mass between (100 GeV;150 GeV)
- dijet $p_t > 100$ GeV

Figure 16: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018




Selections Applied: Event Selections:

- $\text{MET} > 200$ GeV
- no leptons
- no photons

Object Selections:

- jet $p_t > 30$ GeV
- jet $|\eta| < 2.5$
- $\Delta\phi(\text{Jet}, \text{MET})$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50$ GeV
- subleading bjet $p_t > 30$ GeV
- atmost 2 additional jets

Figure 17: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018



Selections Applied:

Event Selections:

- $\text{MET} > 200 \text{ GeV}$
- no leptons
- no photons

Object Selections:

- jet $p_t > 30 \text{ GeV}$
- jet $|\eta| < 2.5$
- $\Delta\phi(\text{Jet}, \text{MET})$
- at least 2 tight bjets (algorithm:DeepFlavB)
- leading bjet $p_t > 50 \text{ GeV}$
- subleading bjet $p_t > 30 \text{ GeV}$
- atmost 2 additional jets

Figure 18: Simulated contribution of various backgrounds to the signal in the resolved $b\bar{b}$ bar case for 2018

