

A new family of distributions: properties and applications

Gauss M. Cordeiro^a, Maria do Carmo S. Lima^a and Pedro Rafael D. Marinho^b

Abstract

This article introduces a new family of distributions by combining the Marshall and Olkin-G and Gamma-G classes. The family has only two extra shape parameters and can be a better model than other well-known classes of distributions. Simulations are performed to illustrate the consistency of the estimators. Its flexibility is shown using two real data sets.

Key Words: Adequacy model package; Marshall and Olkin distribution; R language; Weibull distribution

^a Department of Statistics, Federal University of Pernambuco, Recife, Brazil.

^b Department of Statistics, Federal University of Paraíba, João Pessoa, Brazil.

E-mail: *gauss@de.ufpe.br*, *maria@de.ufpe.br*, *pedro.rafael.marinho@gmail.com*

1 Introduction

The mechanism by adding shape parameters to a baseline distribution has proved to be useful to generate more flexible distributions, study their mathematical properties and improve the goodness-of-fit statistics to the data under study. Many special distributions in these recent families are discussed by Tahir and Nadarajah (2015).

Let $G(x)$ be the cumulative distribution function (CDF) of a baseline distribution and $g(x) = dG(x)/dx$ be the corresponding probability density function (PDF) depending on a parameter vector η . A generalized family is introduced here with two additional shape parameters by transforming the CDF $G(x)$ according to two sequential important generators, which is important for modeling data in several applied areas.

The CDF of the Marshall and Olkin's (1997) (MO-G) family (for $\theta > 0$) is

$$F_{\text{MO-G}}(x) = \frac{G(x)}{\theta + (1 - \theta)G(x)} = \frac{G(x)}{1 - (1 - \theta)[1 - G(x)]}, \quad x \in \mathbb{R}. \quad (1)$$

The density function corresponding to (1) has the form

$$f_{\text{MO-G}}(x) = \frac{\theta g(x)}{[\theta + (1 - \theta)G(x)]^2}. \quad (2)$$

For $\theta = 1$, $f_{\text{MO-G}}(x)$ is just equal to $g(x)$. Equation (2) represents the PDF of the minimum of n independent and identically distributed (iid) random variables having density $g(x)$, say T_1, \dots, T_N , where N has a geometric distribution with probability parameters θ and θ^{-1} if $0 < \theta < 1$ and $\theta > 1$, respectively.

Tahir and Nadarajah (2015, Table 2) presented thirty distributions belonging to this family. It can be easily generated via the quantile function (QF) of G as $Q_{\text{MO-G}}(u) = Q_G(\theta u [\theta u + 1 - u])$ for $u \in (0, 1)$.

Marshall and Olkin considered the exponential and Weibull distributions for the baseline G and derived some structural properties of the generated distributions. The special case that G is an exponential distribution refers to a two-parameter competitive model to the Weibull and gamma distributions.

The CDF of the gamma- G (Γ - G) family (Zografos and Balakrishnan, 2009) is

$$F_{\Gamma-G}(x) = \gamma_1(a, -\log[1 - G(x)]), \quad x \in \mathbb{R}, \quad (3)$$

where $a > 0$ is an extra shape parameter, $\gamma_1(a, z) = \gamma(a, z)/\Gamma(a)$ is the incomplete gamma function ratio and $\gamma(a, z) = \int_0^z t^{a-1} e^{-t} dt$.

Then, the PDF of the Γ - G family can be expressed as

$$f_{\Gamma-G}(x) = \frac{1}{\Gamma(a)} \{-\log[1 - G(x)]\}^{a-1} g(x). \quad (4)$$

Each new Γ - G distribution follows from a specified baseline G . For $a = 1$, the Γ - G family reduces to G . Let Z be a gamma random variable with unit scale and shape parameter $a > 0$; then $W = Q_G(1 - e^{-Z})$ has density (4). So, the Γ - G distribution is easily generated from the gamma distribution and the QF of G .

The remainder of this paper is organized as follows. Section 2 introduces the *Marshall and Olkin-Gamma-G* (MO- Γ - G) families and presents some special models. The maximum likelihood estimates (MLEs) is addressed in Section 3. Some simulations are reported in Section 4 which provide numerical evidence on the biases of the estimators in finite samples. Two empirical applications show the potentiality of the new family in Section 5. Some structural properties are derived in Section 6. Conclusions remarks are offered in Section 7.

2 The New Family

By combining Equations (1) and (3), the CDF of the random variable $X \sim \text{MO-}\Gamma\text{-}G$ of the new family is defined by

$$F_X(x) = \frac{\gamma_1(a, -\log[1 - G(x)])}{\theta + (1 - \theta)\gamma_1(a, -\log[1 - G(x)])}, \quad x \in \mathbb{R}. \quad (5)$$

By differentiating (5), the PDF of X can be expressed as

$$f_X(x) = \frac{\theta \{-\log[1 - G(x)]\}^{a-1} g(x)}{\Gamma(a) \{\theta + (1 - \theta)\gamma_1(a, -\log[1 - G(x)])\}^2}. \quad (6)$$

The density (6) can be interpreted from a sequence of N iid random variables, say Z_1, \dots, Z_N , each one having a gamma density unit scale and shape $a > 0$, assuming that N (is not fixed) has a geometric distribution with probabilities θ and θ^{-1} for $0 < \theta < 1$ and $\theta > 1$, respectively. By transforming the Z_i 's via the baseline QF by $W_i = Q_G(1 - e^{-Z_i})$ (for $i = 1, \dots, N$), Equation (2) defines the PDF of the minimum W_1, \dots, W_n . Making this double composition of the two generators, the proposed family absorbs the impacts of two different flexibilities on applications.

Table 2 provides some special cases of (6), where $\Phi(x)$ and $\phi(x)$ are the CDF and PDF of the standard normal distribution. The density and hazard functions of the Marshall-Olkin- Γ -Weibull (MO- Γ -W) are displayed in Figure 1, which provide more flexibility for these functions in relation to the baseline ones.




Figure 1: (A) MO- Γ -W density. (B) MO- Γ -W hazard.

The CDF (5) can be easily inverted to calculate the QF of the MO- Γ -G distribution, say $x = Q_X(u) = F_X^{-1}(u)$ in terms of $Q_G(\cdot)$, where u is a uniform number in $(0, 1)$. The inverse of $F_X(x) = u$ is easily obtained by combining the inverses of Equations (1) and (5). Equation $F_X(x) = u$ leads to $z = z(u) = \theta u / [1 - (1 - \theta)u]$ and $\gamma_1(a, -\log[1 - G(x)]) = z(u)$, and then the QF of X has a simple form

$$x = Q_G(v(u)),$$

where

$$v(u) = 1 - \exp[-\gamma_1^{-1}(a, z(u))],$$

and $\gamma_1^{-1}(a, w) = Q^{-1}(a, 1 - w)$ is the inverse function of $\gamma_1(a, w)$. Some formulae for $Q^{-1}(a, 1 - w)$ are given in <http://functions.wolfram.com/GammaBetaErf/InverseGammaRegularized/>.

Distribution	Baseline CDF	Generated PDF
Normal	$G(x) = \Phi(x)$	$f_X(x) = \frac{\theta \{-\log[1-\Phi(x)]\}^{a-1} \phi(x)}{\Gamma(a) \{\theta+(1-\theta)\gamma_1(a, -\log[1-\Phi(x)])\}^2}$
Logistic	$G(x) = \frac{1}{1+e^{-x}}$	$f_X(x) = \frac{\theta e^{-x} \{-\log[1-(1+e^{-x})^{-1}]\}^{a-1}}{\Gamma(a) (1+e^{-x})^2 \{\theta+(1-\theta)\gamma_1(a, -\log[1-(1+e^{-x})^{-1}])\}^2}$
Gumbel	$G(x) = 1 - \exp(-e^x)$	$f_X(x) = \frac{\theta \exp(ax - e^x)}{\Gamma(a) \{\theta+(1-\theta)\gamma_1(a, e^x)\}^2}$
Log-Normal	$G(x) = \Phi(\log x)$	$f_X(x) = \frac{\theta \phi(\log x) \{-\log[1-\Phi(\log x)]\}^{a-1}}{\Gamma(a) x \{\theta+(1-\theta)\gamma_1(a, -\log[1-\Phi(\log x)])\}^2}$
Exponential	$G(x) = 1 - \exp(-\lambda x), \lambda > 0$	$f_X(x) = \frac{\theta \lambda^a x^{(a-1)}}{\Gamma(a) \{\theta+(1-\theta)\gamma_1(a, \lambda x)\}^2}$
Weibull	$G(x) = 1 - \exp(-(x/\beta)^\alpha), \alpha, \beta > 0$	$f_X(x) = \frac{\alpha \theta x^{\alpha-1} [(x/\beta)^\alpha]^{a-1} \exp[-(x/\beta)^\alpha]}{\beta^\alpha \Gamma(a) \{\theta+(1-\theta)\gamma_1[a, (x/\beta)^\alpha]\}^2}$
Gamma	$G(x) = \gamma_1(\alpha, \beta x), \alpha, \beta > 0$	$f_X(x) = \frac{\theta \beta^\alpha x^{\alpha-1} e^{-\beta x} \{-\log[1-\gamma_1(\alpha, \beta x)]\}^{a-1}}{\Gamma(a) \{\theta+(1-\theta)\gamma_1(a, -\log[1-\gamma_1(\alpha, \beta x)])\}^2}$
Pareto	$G(x) = 1 - \frac{1}{(1+x)^\nu}, \nu > 0$	$f_X(x) = \frac{\theta e^{-x} [\nu \log(1+x)]^{a-1} g(x)}{\Gamma(a) (1+e^{-x})^2 \{\theta+(1-\theta)\gamma_1(a, \nu \log[1+x])\}^2}$

Table 1: Special distributions in the MO-Γ-G family.

3 Estimation

The MO- Γ -G family can be fitted to real data using the **AdequacyModel** package in the R software. This package does not require the log-likelihood function and it computes the MLEs, their standard errors (SEs) and the formal statistics defined in Section 5. It is only necessary to provide the PDF and CDF of the distribution to be fitted to a data set.

If x_i is one observation from (6) and $\boldsymbol{\eta}$ is a q -parameter vector specifying $G(\cdot)$, the log-likelihood function for $\boldsymbol{\theta}^\top = (a, \theta, \boldsymbol{\eta}^\top)$ from n observations is

$$\begin{aligned} \ell(\boldsymbol{\theta}) = & n \log(\theta) + n \log(\gamma) + n a \gamma \log(\lambda) + (a\gamma - 1) \sum_{i=1}^n \log(x_i) - \lambda^\gamma \sum_{i=1}^n \log(x_i) \\ & - n \log[\Gamma(a)] - 2 \sum_{i=1}^n \log\{\theta + (1 - \theta)\gamma_1[a, (\lambda x_i)^\gamma]\}. \end{aligned} \quad (7)$$

Due to the impossibility of obtaining the MLEs in closed form, numerical methods to maximize $\ell(\cdot)$ are necessary. Several programming languages and statistical software provides functions and routines that make it easy to obtain numerical estimates by various interactive methods. In practice, obtaining the MLEs for the parameters that index a distribution are commonly obtained in this way, since the Newton and quasi-Newton methods produce satisfactory results under reasonable conditions of the object function, that is, when they do not impose conditions that disturb the convergence of the algorithms.

To obtain the MLEs, the **AdequacyModel** package of the programming language R was used, see R Core Team (2020). This library, created and updated by one of the authors of this paper, is widely cited by several papers in Statistics and serves as a basis for other library implementations available on the Comprehensive R Archive Network - CRAN. By using the goodness.fit function, it provides an implementation of (6) in R giving $\ell(\cdot)$ by returning the MLEs and some adequacy measures of fit. Further details regarding this package can be obtained from Marinho et al. (2019).

4 Simulations

Due to the probable absence of MLEs in closed-form for distributions belonging to the MO- Γ -G family, it is necessary to examine the precision of the estimates calculated numerically. For doing that, the biases of the estimators of the parameters of the MO- Γ -Dagum($\theta, a, \alpha, \beta, p$) distribution are calculated, where $G \sim \text{Dagum}(\alpha, \beta, p)$ is the baseline distribution. All parameters are taken equal to one for different sample sizes reported in Table 2.

Ten thousand Monte Carlo simulations are performed for each sample size to calculate the estimates by the BFGS method. The figures in Table 2 indicate that this method behaves well when the sample size increases. This is theoretically expected. However, in practice, difficulties can be faced in other families of distributions due to the flatness of the log-likelihood function.

All simulations can be reproduced using the script in Appendix A. The simulations are parallelized and able to use all threads available by a multicore processor, thus making them more computationally efficient and consequently requiring less time to complete. The simulations are performed on a computer with an Intel Core i5-8265U processor with

8 threads working at a maximum frequency of 3.90 GHz, requiring, on these hardware, a time of 14.36 hours to perform all simulations. The figures in Table 2 reveal that the average biases of the MLEs could be very reduced only for $n > 1,000$.

To generate observations from the random variable X with density f , the well-known Acceptance-Rejection Algorithm for continuous random variables, which is very useful when the quantile function involves complex functions that can lead to some numerical inaccuracies. For doing this, another random variable Y is chosen such that it can generate observations from a PDF h with the same support as f . Then, the acceptance and rejection algorithm is defined by the following steps:

1. Generate an outcome y from Y ;
2. Generate an observation u from a uniform random variable U in $(0, 1)$;
3. If $u < \frac{f(y)}{cg(y)}$, where c is a real constant, accept $x = y$; otherwise reject y as an outcome from X and return to 1.

The constant c should be chosen in such a way that $\frac{f(y)}{cg(y)} \leq 1$. Thus, to minimize the computational cost of generating observations from X through the generated observations from Y , c is chosen as the lowest possible value to maximize the likelihood of acceptance. Further details of this method can be found in Rizzo (2019).

Table 2: Mean biases of the MLEs obtained using the BFGS method calculated from the Monte-Carlo simulation.

n	$B(\hat{\theta})$	$B(\hat{a})$	$B(\hat{\alpha})$	$B(\hat{\beta})$	$B(\hat{p})$	Time (mins)
10	0.2376	2.1635	2.7557	1.6282	1.3057	1.1430
20	0.4154	2.4639	1.5728	1.8082	0.7383	1.6248
60	0.7214	2.2432	0.5667	1.8815	0.2872	3.3954
100	0.6146	1.9579	0.3253	1.6148	0.2651	4.9628
200	0.3838	1.3894	0.1827	1.1773	0.3701	8.1457
400	0.2166	0.9635	0.1076	0.6181	0.3957	13.7370
600	0.1269	0.7242	0.0772	0.3968	0.3637	17.9310
1,000	0.0553	0.4885	0.0521	0.2328	0.2636	22.8784
2,000	0.0456	0.3087	0.0334	0.0990	0.1722	38.4593
5,000	-0.0058	0.1307	0.0146	0.0117	0.0171	52.8098
10,000	-0.0146	0.0842	0.0095	0.0095	0.0031	95.8380
20,000	-0.0090	0.0330	0.0038	0.0005	-0.0099	126.4260
30,000	-0.0028	0.0183	0.0012	-0.0036	-0.0029	182.0760
50,000	-0.0057	0.0124	0.0015	0.0016	-0.0021	291.9300

5 Applications

Consider the Weibull baseline. Two applications are provided to compare the new generated model with seven extended Weibull distributions, namely the beta-Weibull (β -W) (Famoye *et al.*, 2005), Kumaraswamy Weibull (Kw-W) (Cordeiro and Nadarajah, 2010), Marshall-Olkin Weibull (MO-W) (Ahmed *et al.*, 2017), Marshall-Olkin Extended Weibull (MOE-W) (Cordeiro *et al.*, 2019), exponentiated Weibull (exp-W) (Mudholkar and Srivastava, 1993), gamma Weibull (Γ -W) (Cordeiro *et al.*, 2016) and exponentiated generalized Weibull (EG-W) (Oguntunde *et al.*, 2015) (with $a = 1$). Some of these distributions are widely used in practice.

The log-likelihood for the Marshall-Olkin-Gamma-Weibull (MO-Γ-W) from one observation is

$$\begin{aligned}\ell(\boldsymbol{\theta}) = & \log(\theta) + \log(\gamma) + (a\gamma) \log(\lambda) + (a\gamma - 1) \log(x) - (\gamma x)^\gamma - \log[\Gamma(a)] \\ & - 2 \log\{\theta + (1 - \theta)\gamma_1[a, (\lambda x)^\gamma]\},\end{aligned}\quad (8)$$

where $\boldsymbol{\theta} = (a, \theta, \lambda, \gamma)^\top$. The components of the score function are

$$U_a(\boldsymbol{\theta}) = \gamma \log(\lambda) + \gamma \log(x) - \psi^{(0)}(a) - \frac{2 \{(1 - \theta)A - (1 - \theta)\psi^{(0)}(a)\gamma_1[a, (x\lambda)^\gamma]\}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (x\lambda)^\gamma]},$$

$$U_\theta(\boldsymbol{\theta}) = \frac{1}{\theta} - \frac{2 \{\Gamma(a) - \gamma_1[a, (\lambda x)^\gamma]\}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (\lambda x)^\gamma]},$$

$$U_\lambda(\boldsymbol{\theta}) = \frac{\gamma}{\lambda} [a - (\lambda x)^\gamma] + \frac{2\gamma \lambda^{-1}(\lambda x)^a \gamma (1 - \theta) \exp\{-(\lambda x)^\gamma\}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (x\lambda)^\gamma]}$$

and

$$U_\gamma \boldsymbol{\theta} = \frac{1}{\gamma} + a \log(\lambda) + a \log(x) - (\lambda x)^\gamma \log(\lambda x) + \frac{2(1 - \theta)(\lambda x)^{\gamma a} \log(\lambda x) \exp\{-(\lambda x)^\gamma\}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (x\lambda)^\gamma]},$$

where

$$A = G_{2,3}^{3,0} \left[(x\lambda)^\gamma \middle| \begin{matrix} 1, 1 \\ 0, 0, a \end{matrix} \right] + \log [(\lambda x)^\gamma] \gamma_1[a, (x\lambda)^\gamma],$$

$\psi^{(n)}(x)$ is the n -th derivative of the digamma function,

$$A = \psi^{(0)}(a) - \log [(\lambda x)^\gamma] - G_{2,3}^{3,0} \left((\lambda x)^\gamma \middle| \begin{matrix} 1, 1 \\ 0, 0, a \end{matrix} \right),$$

and $G_{p,q}^{m,n} \left(z \middle| \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \right)$ is the Meijer G function.

The **AdequacyModel** is used to fit the distributions cited before to two real data sets. The SANN method, which is a variant of simulated annealing (Belisle, 1992), is considered. The distributions are compared via the Anderson Darling (A*) and Cramér Von Mises (W*) statistics reported in the `goodness.fit` function.

For the first data set, a modification of the “FoodExpenditure” data from the **betareg** package is considered, which refers to the proportions of income spent on food for a random sample of 38 households in a large US city. Here, the household expenditures for food are given by

$$data = FoodExpenditure_{food} / \#(FoodExpenditure_{food}),$$

where $FoodExpenditure_{food}$ is the random variable corresponding to the household expenditures for food and $\#(\cdot)$ indicates the number of observations on this variable. The MLEs, their standard errors (SEs) (in parentheses) and the adequacy statistics are listed in Table 3. The proposed model has a better fit to the data than the others.

As a second application, the systolic blood pressure of persons collected in a pilot study about hypertension in the Dominican Republic in 1997 found in <http://biostat.mc.vanderbilt.edu/wiki/Main/DataSets> are analyzed. The MLEs, their SEs and the values of the statistics are reported in Table 4. The proposed distribution outperforms the rest of them by comparing the measures of these formal statistics.

Table 3: Application 1

Model	a	θ	λ	γ	W^*	A^*
MO- Γ -W($a, \theta, \lambda, \gamma$)	0.926134 (0.02626926)	1.379664 (0.22381086)	33.323073 (0.28530971)	25.398809 (0.08259864)	0.0339	0.2376
β -W($a, \theta, \lambda, \gamma$)	9.92882 (0.02908181)	0.1700880 (0.02049663)	9.759469 (<0.0001)	1.530541 (<0.0001)	0.043567	0.2594618
KW-W($a, \theta, \lambda, \gamma$)	0.04987575 (0.008090352)	99.99989793 (16.225905058)	1.07602954 (0.003156126)	23.40287099 (0.014646102)	1.330915	6.742609
MOE-W($a, \theta, \lambda, \gamma$)	0.1366666 (0.1599182)	2.020436 (<0.0001)	62.72201 (<0.0001)	4.2956659 (0.7365535)	0.03541554	0.2579222
EGW(a, b, λ, γ)	5.6189861421 (0.0028147823)	6.1833138579 (0.0009807091)	1.2870816760 (0.1159810838)	1.3798562942 (0.1480747352)	0.03715255	0.2518879
MO-W(a, λ, γ)	0.15920715 (0.07170351)	- (-)	1.58609458 (0.13353716)	4.26713785 (0.16650667)	0.03456333	0.257339
exp-W(a, λ, γ)	6.1102948 (0.4222173)	- (-)	4.4680469 (0.3175876)	1.3858740 (0.1677722)	0.03726684	0.2523749
γ -W(a, λ, γ)	5.751546 (0.0015006)	- (-)	10.0000 (0.0001163185)	1.208773 (0.00822782)	0.0879	0.6599

Table 4: Application 2

Model	a	θ	λ	γ	W^*	A^*
MO- Γ -W($a, \theta, \lambda, \gamma$)	9.629304 (0.006217876)	3.640779 (0.182495785)	6.260826 (0.024446563)	12.823429 (0.007965221)	0.5093	2.8076
β -W($a, \theta, \lambda, \gamma$)	31.08471 (0.01279570)	47.14636 (<0.0001)	0.01698383 (0.00014535)	2.054037 (<0.0001)	0.7540428	4.279418
KW-W($a, \theta, \lambda, \gamma$)	7363.281 (0.04194304)	0.03925762 (0.0004194304)	1.467640 (<0.0001)	0.6146940 (<0.0001)	0.5351	2.9617
MOE-W($a, \theta, \lambda, \gamma$)	101.13471834 (46.782882038)	0.42386507 (0.100832102)	0.03095935 (0.003644092)	1.70240332 (0.168142275)	1.14418	6.644617
EGW(a, b, λ, γ)	0.2351691 (0.002540837)	140.0000 (3.278565)	0.4576370 (<0.0001)	0.7425738 (<0.0001)	0.8925	5.3338
MO-W(a, λ, γ)	173.2139 (0.00016394)	- (-)	0.02125455 (0.000212794)	1.601970 (0.0001398109)	1.476088	8.58705
exp-W(a, λ, γ)	69.02916 (0.08389090)	- (-)	0.02405120 (0.0002249256)	1.345536 (<0.0001)	0.8899261	5.094141
Γ -W(a, λ, γ)	9.11229459 (0.96330688)	- (-)	0.02617729 (0.00291875)	1.74641178 (0.08030313)	0.6227098	3.488268

6 Mathematical properties

Some main mathematical properties are addressed for the MO- Γ -G family based on a general linear representation for its density function and those properties of exponentiated-G (exp-G) distributions.

6.1 Linear Representation

For an arbitrary CDF $G(x)$, the CDF and PDF of the exponentiated-G (exp-G) distribution with power parameter $a > 0$ are

$$\Pi_a(x) = G(x)^a \quad \text{and} \quad \pi_a(x) = a g(x) G(x)^{a-1},$$

respectively. This class of distributions is quite useful in several applications. In fact, Tahir and Nadarajah (2015) cited more than seventy papers on exponentiated distributions in their Table 1.

First, the MO-G cumulative distribution (2) admits the linear combination (Barreto-Souza *et al.*, 2013)

$$F_{\text{MO-G}}(x) = \sum_{i=0}^{\infty} w_i^{\text{MO-G}} \Pi_{i+1}(x) = \sum_{i=0}^{\infty} w_i^{\text{MO-G}} G(x)^{i+1}, \quad (9)$$

where the coefficients are (for $i = 0, 1, \dots$)

$$w_i^{\text{MO-G}} = w_i^{\text{MO-G}}(\theta) = \begin{cases} \frac{(-1)^i \theta}{(i+1)} \sum_{j=i}^{\infty} (j+1) \binom{j}{i} \bar{\theta}^j, & \theta \in (0, 1), \\ \theta^{-1} (1 - \theta^{-1})^i, & \theta > 1, \end{cases}$$

and $\bar{\theta} = 1 - \theta$.

Second, the linear combination for the Γ -G cumulative distribution (4) follows from Castellares and Lemonte (2015) as

$$F_{\Gamma\text{-G}}(x) = \sum_{j=0}^{\infty} w_j^{\Gamma\text{-G}} \Pi_{a+j}(x). \quad (10)$$

Here,

$$w_j^{\Gamma\text{-G}} = w_j^{\Gamma\text{-G}}(a) = \frac{\varphi_j(a)}{(a+j)},$$

$$\varphi_0(a) = \frac{1}{\Gamma(a)}, \quad \varphi_j(a) = \frac{(a-1)}{\Gamma(a)} \psi_{j-1}(j+a-2), \quad j \geq 1,$$

and

$$\psi_{n-1}(x) = \frac{(-1)^{n-1}}{(n+1)!} \left[H_n^{n-1} - \frac{x+2}{n+2} H_n^{n-2} + \frac{(x+2)(x+3)}{(n+2)(n+3)} H_n^{n-3} - \dots \right. \\ \left. + (-1)^{n-1} \frac{(x+2)(x+3) \cdots (x+n)}{(n+2)(n+3) \cdots (2n)} H_n^0 \right],$$

is the Stirling polynomial, $H_{n+1}^m = (2n+1-m)H_n^m + (n-m+1)H_n^{m-1}$ is a positive integer, $H_0^0 = 1$, $H_{n+1}^0 = 1 \times 3 \times 5 \times \cdots \times (2n+1)$ and $H_{n+1}^n = 1$.

By inserting (10) in Equation (9) and via an equation for a power series raised to a positive integer (Gradshteyn and Ryzhik, 2000), the expansion for the cdf of the MO- Γ -G distribution reduces to

$$F_{\text{MO-}\Gamma\text{-G}}(x) = \sum_{i=0}^{\infty} w_i^{\text{MO-}\Gamma} G(x)^{(i+1)a} \left[\sum_{j=0}^{\infty} w_j^{\Gamma\text{-G}} G(x)^j \right]^{i+1} \\ = \sum_{i=0}^{\infty} w_i^{\text{MO-}\Gamma} G(x)^{(i+1)a} \sum_{j=0}^{\infty} c_{i+1,j} G(x)^j = \sum_{i,j=0}^{\infty} d_{i,j} \Pi_{(i+1)a+j}(x),$$

where $d_{i,j} = d_{i,j}(a, \theta) = w_i^{\text{MO-G}} c_{i+1,j}(a)$, $c_{i+1,0}(a) = (w_0^{\Gamma\text{-G}})^{i+1}$ and, for $m \geq 1$, $c_{i+1,m}(a) = \frac{1}{mw_0^{\Gamma\text{-G}}} \sum_{r=1}^m [r(i+2) - m] w_r^{\Gamma\text{-G}} c_{i+1,m-r}(a)$.

By differentiating the last equation, the expansion for the MO- Γ -G density follows as

$$f_{\text{MO-}\Gamma\text{-G}}(x) = \sum_{i,j=0}^{\infty} d_{i,j} \pi_{(i+1)a+j}(x). \quad (11)$$

So, some structural properties of the proposed family can be found from the double linear combination (11) and those properties of the exp-G distribution. In most applications, the indices i and j can vary up to five.

6.2 Some quantities

Hereafter, let $T_{i,j} \sim \text{exp-G}[(i+1)a+j]$. The n th moment of X follows from (11) as

$$\mu'_n = E(X^n) = \sum_{i,j=0}^{\infty} d_{i,j} E(T_{i,j}) = \sum_{i,j=0}^{\infty} [(i+1)a+j-1] d_{i,j} \tau[n, (i+1)a+j-1], \quad (12)$$

where

$$\tau(n, a) = \int_{-\infty}^{\infty} x^n G(x)^a g(x) dx = \int_0^1 Q_G(u)^n u^a du.$$

Expressions for moments of several exponentiated distributions can be found in the papers cited in Tahir and Nadarajah (2015, Table 1). We give just one example from Equation (12) by taking the exponential distribution with rate $\lambda > 0$ for the baseline G. It follows easily as

$$\mu'_n = n! \lambda^n \sum_{i,j,m=0}^{\infty} \frac{(-1)^{n+m} [(i+1)a+j] d_{i,j}}{(m+1)^{n+1}} \binom{(i+1)a+j-1}{m}.$$

A general expansion for the moment generating function (MGF) $M(t) = E(e^{tX})$ of X can be expressed from (11)

$$M(t) = \sum_{i,j=0}^{\infty} d_{i,j} M_{i,j}(t) = \sum_{i,j=0}^{\infty} [(i+1)a+j] d_{i,j} \rho(t, (i+1)a+j-1), \quad (13)$$

where $M_{i,j}(t)$ is the MGF of $Y_{i,j}$ and

$$\rho(t, a) = \int_{-\infty}^{\infty} e^{tx} G(x)^a g(x) dx = \int_0^1 \exp\{t Q_G(u)\} u^a du.$$

The MGFs of many MO- Γ -G distributions can be determined from Equation (13). For example, the generating function of the MO- Γ -exponential with parameter λ (if $t < \lambda^{-1}$) is

$$M(t) = \sum_{i,j=0}^{\infty} [(i+1)a+j] d_{i,j} B((i+1)a+j, 1-\lambda t).$$

7 Conclusions

A new family of distributions called the Marshall and Olkin-Gamma-G family with two shape parameters is introduced. The estimation of the unknown parameters is done via maximum likelihood and a simulation study is conducted to verify its adequacy. Additionally, the usefulness of the proposed family is shown empirically using two real data sets.

A Appendix: Simulation code

The script below is used to perform the simulations whose results are given in Table 2 of Section 4. It is developed a script using the programming language R to be used in other simulations or even to check the results presented in this paper.

Listing 1: Monte-Carlo simulations for different sample sizes.

```
1 # Title: Marshall Olkin Gamma-G (MOGG)
2 # Author: Pedro Rafael D. Marinho
3
4 # Loading libraries. -----
5 library(parallel)
6 library(tibble)
7 library(pbmccapply)
8 library(magrittr)
9 library(purrr)
10 library(xtable)
11
12 # Baseline functions. -----
13 pdf_dagum <- function(x, alpha, beta, p)
14   alpha * p / x * (x / beta) ^ (alpha * p) / ((x / beta) ^ alpha + 1) ^ (p + 1)
15 # integrate(f = pdf_dagum, lower = 0, upper = Inf, alpha = 1.2,
16 #           beta = 1.6, p = 2.2)
17
18 cdf_dagum <- function(x, alpha, beta, p)
19   (1 + (x / beta) ^ (-alpha)) ^ (-p)
20 # cdf_dagum(x = Inf, alpha = 1, beta = 4, p = 1)
21
22 # This function creates MOGG functions. -----
23 pdf_mogg <- function(g, G) {
24   # Using Closures.
25   function(x, theta, a, ...) {
26     if (theta <= 0 || a <= 0)
27       warning("The \"a\" and \"theta\" parameters must be greater than zero.")
28     num <-
29       theta * (-log(1 - G(x = x, ...))) ^ (a - 1) * g(x = x, ...)
30     den <-
31       gamma(a) *
32       (theta + (1 - theta) *
33         pgamma(-log(1 - G(x = x, ...)), a, 1L)) ^ 2
34     num / den
35   }
36 }
37
38 rdagum <- function(n = 1L, alpha, beta, p) {
39   beta * (runif(n = n, min = 0, max = 1) ^ (-1 / p) - 1) ^ (-1 / alpha)
40 }
41
42 pdf_mogdagum <- pdf_mogg(g = pdf_dagum, G = cdf_dagum)
43
44 # MOG-Dagum -----
45 rmogdagum <- function(n = 1L, theta, a, alpha, beta, p) {
46   cond_c <- function(x, theta, a, alpha, beta, p) {
47     num <- pdf_mogdagum(x, theta, a, alpha, beta, p)
48     den <- pdf_dagum(x,
49                       alpha = alpha,
50                       beta = beta,
51                       p = p)
52     - num / den
```

```

53 }
54
55 x_max <-
56   optim(
57     fn = cond_c,
58     method = "BFGS",
59     par = 1,
60     theta = theta,
61     a = a,
62     alpha = alpha,
63     beta = beta,
64     p = p
65   )$par
66
67 c <-
68   pdf_mogdagum(x_max, theta, a, alpha, beta, p) / pdf_dagum(x_max,
69                                                             alpha = alpha,
70                                                             beta = beta,
71                                                             p = p)
72
73 criterion <- function(y, u) {
74   num <- pdf_mogdagum(y, theta, a, alpha, beta, p)
75   den <- pdf_dagum(y,
76                   alpha = alpha,
77                   beta = beta,
78                   p = p)
79   u < num / (c * den)
80 }
81
82 values <- double(n)
83 i <- 1L
84 repeat {
85   y <- rdagum(
86     n = 1L,
87     alpha = alpha,
88     beta = beta,
89     p = p
90   )
91   u <- runif(n = 1L, min = 0, max = 1)
92
93   if (criterion(y, u)) {
94     values[i] <- y
95     i <- i + 1L
96   }
97   if (i > n)
98     break
99 }
100 values
101 }
102
103 # Testing the rmogw Function -----
104 theta = 5
105 a = 1
106 alpha = 5
107 beta = 1
108 p = 1
109 pdf_mogdagum <- pdf_mogg(g = pdf_dagum, G = cdf_dagum)
110 sample_data <- rmogdagum(n = 250L, theta, a, alpha, beta, p)
111 x <- seq(0, max(sample_data), length.out = 500L)
112 hist(sample_data,
113       probability = TRUE,
114       xlab = "",
115       main = "")
116 lines(x, pdf_mogdagum(x, theta, a, alpha, beta, p))
117
118 # Monte Carlo simulations. -----
119 mc <- function(n = 250L,
120               M = 1e3L,
121               par_true,
122               method = "BFGS") {
123   theta <- par_true[1L]
124   a <- par_true[2L]
125   alpha <- par_true[3L]
126   beta <- par_true[4L]

```

```

127 p <- par_true[5L]
128
129 # Log-likelihood function. -----
130 pdf_mogw <- pdf_mogg(g = pdf_dagum, G = cdf_dagum)
131 log_likelihood <- function(x, par) {
132   theta <- par[1L]
133   a <- par[2L]
134   alpha <- par[3L]
135   beta <- par[4L]
136   p <- par[5L]
137
138   - sum(log(
139     pdf_mogdagum(
140       x,
141       theta = theta,
142       a = a,
143       alpha = alpha,
144       beta = beta,
145       p = p
146     )
147   ))
148 }
149
150 myoptim <-
151   function(...)
152     tryCatch(
153       expr = optim(...),
154       error = function(e)
155         NA
156     )
157
158 one_step_mc <- function(i) {
159   sample_data <- rmogdagum(n, theta, a, alpha, beta, p)
160
161   result <- myoptim(
162     fn = log_likelihood,
163     par = c(1, 1, 1, 1, 1),
164     x = sample_data,
165     method = method
166   )
167
168   while (is.na(result) || result$convergence != 0) {
169     sample_data <- rmogdagum(n, theta, a, alpha, beta, p)
170     result <- myoptim(
171       fn = log_likelihood,
172       par = c(1, 1, 1, 1, 1),
173       method = method,
174       x = sample_data
175     )
176   }
177
178   result$par
179 }
180
181 result_vector <-
182   unlist(
183     pbmcapply::pbmclapply(
184       X = 1L:M,
185       FUN = one_step_mc,
186       mc.cores = parallel::detectCores()
187     )
188   )
189
190 result <-
191   tibble::as_tibble(matrix(result_vector, byrow = TRUE, ncol = 5L))
192
193 names(result) <- c("theta", "a", "alpha", "beta", "p")
194
195 result
196 }
197
198
199 bias_function <- function(x, par_true) {
200   x - par_true

```

```

201 }
202
203 mse_function <- function(x, par_true) {
204   (x - par_true) ^ 2
205 }
206
207 simulate <- function(n) {
208   # True parameters (theta, a, alpha, beta and p) -----
209   true_parameters <- c(1, 1, 1, 1, 1)
210
211   set.seed(1L, kind = "L'Ecuyer-CMRG")
212   t0 <- Sys.time()
213   result_mc <-
214     mc(
215       n = n,
216       M = 1e4L,
217       par_true = true_parameters,
218       method = "BFGS"
219     )
220   total_time <- Sys.time() - t0
221
222   mc.reset.stream()
223
224   # Average Bias of Estimators -----
225   eval(parse(
226     text = glue(
227       "bias_{n} <- apply(X = result_mc, MARGIN = 1L, FUN = bias_function,
228         par_true = true_parameters) %>%
229         apply(MARGIN = 1L, FUN = mean)"
230     )
231   ))
232   eval(parse(text = glue(
233     "save(file = \"bias_{n}.RData\", bias_{n})"
234   )))
235
236   # Mean Square Error -----
237   eval(parse(
238     text = glue(
239       "mse_{n} <- apply(X = result_mc, MARGIN = 1L, FUN = mse_function,
240         par_true = true_parameters) %>%
241         apply(MARGIN = 1L, FUN = mean)"
242     )
243   ))
244   eval(parse(text = glue(
245     "save(file = \"mse_{n}.RData\", mse_{n})"
246   )))
247
248   # Total Time -----
249   eval(parse(text = glue("time_{n} <- total_time")))
250   eval(parse(text = glue(
251     "save(file = \"time_{n}.RData\", time_{n})"
252   )))
253
254   # Result MC
255   eval(parse(text = glue("result_{n} <- result_mc")))
256   eval(parse(text = glue(
257     "save(file = \"result_{n}.RData\", result_{n})"
258   )))
259
260 }
261
262 walk(
263   .x = c(
264     10,
265     20,
266     60,
267     100,
268     200,
269     400,
270     600,
271     1000,
272     2000,
273     5000,
274     10000,

```

```

275     20000,
276     30000,
277     50000
278   ),
279   .f = simulate
280 )
281
282 first_col <-
283   c(10, 20, 60, 100, 200, 400, 600, 1000, 5000, 10000, 20000, 30000, 50000)
284
285 tabela <-
286   rbind(
287     bias_10,
288     bias_20,
289     bias_60,
290     bias_100,
291     bias_200,
292     bias_400,
293     bias_600,
294     bias_1000,
295     bias_5000,
296     bias_10000,
297     bias_20000,
298     bias_30000,
299     bias_50000
300   )
301 tabela <- cbind(n = first_col, tabela)
302 rownames(tabela) <- NULL
303
304 tabela <- tibble::as_tibble(tabela)
305
306 latex <-
307   print.xtable(
308     xtable(tabela,
309             caption = "Mean bias of EMV obtained by the BFGS method in 10,000
310                       Monte Carlo repetitions.",
311             digits = 4L),
312     print.results = FALSE
313   )
314
315 writeLines(
316   c(
317     "\\documentclass[12pt]{article}",
318     "\\begin{document}",
319     "\\thispagestyle{empty}",
320     latex,
321     "\\end{document}"
322   ),
323   "mc_simulation.tex"
324 )
325
326 tools::texi2pdf("mc_simulation.tex", clean = TRUE)

```

References

- Alzaatreh, A., Famoye, F. and Lee, C. A new method for generating families of continuous distributions. *Metron* **71**(1) 63-79 (2013).
- Barreto-Souza, W., Lemonte, A.J. and Cordeiro, G.M. General results for the Marshall and Olkin's family of distributions. *Anais da Academia Brasileira de Ciências* **85**(1) 3-21 (2013).
- Birnbaum, Z.W. and S.C. Saunders. A new family of life distributions. *Journal of Applied Probability* **6**(2) 319-327 (1969).
- Castellares, F. and Lemonte, A.J. A new generalized Weibull distribution generated by gamma random variables. *Journal of the Egyptian Mathematical Society* **23**(2)

- 382-390 (2015).
- Cordeiro, G.M. and de Castro, M. A new family of generalized distributions. *Journal of Statistical Computation and Simulation* **81**(7) 883-898 (2011).
- Eugene, N., Lee, C. and Famoye, F. Beta-normal distribution and its applications. *Communications in Statistics – Theory and Methods* **31**(4) 497-512 (2002).
- Gradshteyn, I.S. and Ryzhik, I.M. Table of Integrals, Series and Products. Sixth edition. San Diego: Academic Press (2000).
- Gupta, R.C., Gupta, R.D. and Gupta, P.L. Modeling failure time data by Lehmann alternatives. *Communications in Statistics – Theory and Methods* **27**(4) 887-904 (1998).
- Gupta, R.D. and Kundu, D. Exponentiated exponential family: an alternative to gamma and Weibull distributions. *Biometrical Journal* **43**(1) 117-130 (2001).
- Lemonte, A.J., Cordeiro, G.M. and Moreno-Arenas, G. A new useful three-parameter extension of the exponential distribution. *Statistics* **50**(2) 312-337 (2016).
- Marinho, P.R.D., Silva, R.B., Bourguignon, M., Cordeiro, G. M. and Nadarajah. S. Adequacy Model: An R package for probability distributions and general purpose optimization. *PloS one*, **14**(8), e0221487 (2019).
- Marshall, A.W. and Olkin, I. A new method for adding a parameter to a family of distributions with application to the exponential and Weibull families. *Biometrika* **84**(3) 641-652 (1997).
- Mudholkar, G.S. and Srivastava, D.K. The exponentiated Weibull family: a reanalysis of the bus-motor-failure data. *Technometrics* **37**(4) 436-445 (1995).
- Nadarajah, S., Cordeiro, G.M. and Ortega, E.M.M. Some general results for the beta-modified Weibull distribution. *Journal of Statistical Computation and Simulation* **81**(10) 1211-1232 (2011).
- Nadarajah, S., Cordeiro, G.M. and Ortega, E.M.M. General results for the Kumaraswamy G distribution. *Journal of Statistical Computation and Simulation* **87**(7) 951-979 (2012).
- Nadarajah, S. and Gupta, A.K. The exponentiated gamma distribution with application to drought data. *Calcutta Statistical Association Bulletin* **59**(1-2) 29-54 (2007).
- Nadarajah, S. and Haghighi, F. An extension of the exponential distribution. *Statistics* **45**(6) 543-558 (2011).
- Nadarajah, S. and Kotz, S. The exponentiated-type distributions. *Acta Applicandae Mathematicae* **92**(2) 97-111 (2006).
- Nichols, M.D. and Padgett, W.J. A Bootstrap control chart for Weibull percentiles. *Quality and Reliability Engineering International* **22**(2) 141-151 (2006).
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/> (2020).

- Proschan, F. Theoretical explanation of observed decreasing failure rate. *Technometrics* **5**(3) 375-383 (1963).
- Ristić, M.M. and Balakrishnan, N. The gamma-exponentiated exponential distribution. *Journal of Statistical Computation and Simulation* **82**(8) 1191-1206 (2012).
- Rizzo, M. L. Statistical computing with R. CRC Press (2019).
- Stacy, E.W. A Generalization of the Gamma Distribution. *Annals of Mathematical Statistics* **33**(3) 1187-1192 (1962).
- Tahir, M.H. and Nadarajah, S. Parameter induction in continuous univariate distributions: Well-established G families. *Anais da Academia Brasileira de Ciências* **87**(2) 539-568 (2015).
- Zografos, K. and Balakrishnan, N. On families of beta-and generalized gamma-generated distribution and associate inference. *Statistical Methodology* **6**(4) 344-362 (2009).