




The Marshall and Olkin-G and Gamma-G family of distributions: properties and applications

Authors: MARIA DO CARMO S. LIMA 
– Department of Statistics, Federal University of Pernambuco,
Pernambuco, Recife, Brazil (maria@de.ufpe.br)
GAUSS M. CORDEIRO 
– Department of Statistics, Federal University of Pernambuco,
Pernambuco, Recife, Brazil (gauss@de.ufpe.br)
PEDRO RAFAEL D. MARINHO  
– Department of Statistics, Federal University of Paraíba,
Paraíba, João Pessoa, Brazil (pedro.rafael.marinho@gmail.com)

1 Received: December 2020 Revised: May 2022 Accepted: Month 0000

2 Abstract:

- 3 • The article introduces a new family by combining the Marshall and Olkin-G and
4 Gamma-G classes. It has only two extra shape parameters and can be a better model
5 than other existing classes of distributions. Simulations are performed to verify the
6 consistency of the estimators. Its flexibility is shown by means of two real data sets.

7 Key-Words:

- 8 • *Applications; distribution family; mathematical properties; simulations.*

9 AMS Subject Classification:

- 10 • 60E05, 62E10, 62E15.

1. INTRODUCTION

11 The advances in the field of Data Science requires the search for new families
12 of distributions that adequately model real data has been increasing steadily in the

*Corresponding author

last years. The construction of different generation methods and even generators of families has made it difficult to compare new proposals. In the midst of a huge set of existing families in the literature of new distributions, to find a proposal that is in fact an excellent competitor when compared to other existing ones, in terms of adjustment to real data sets and also that does not present estimation problems, is a major challenge.

The classes of distributions in the early 1980s were based on the simple idea of adding parameters to a baseline distribution. The mechanism by adding shape parameters to a baseline distribution has proved to be useful to make the generated distributions more flexible especially for studying tail properties than existing ones and for improving their goodness-of-fit statistics to real data. Many special distributions in these families are discussed by Tahir and Nadarajah (2015).

The addition of parameters in the construction of new distributions/families was improved by the inclusion of mathematical functions known in the literature, such as beta and gamma functions, for example, which produce new generators with more flexible properties than their baselines. Two well-known examples are the beta-G (Eugene et al., 2002) and gamma-G (Zografos and Balakrishnan, 2009) generators.

However, the inclusion of such functions for generating new families brought, in some cases, problems for parameter estimation. So, despite the fit being more suitable for some types of data and, therefore, having a superior performance when compared to other generators, the estimation process can often be a problem.

In this context, this work presents a new family obtained by composing a very competitive class in the literature with another class that has the gamma function in its structure.

Let $G(x)$ be the cumulative distribution function (CDF) of a baseline distribution and $g(x) = dG(x)/dx$ be the corresponding probability density function (PDF) depending on a parameter vector η . A generalized family is presented with two extra shape parameters by transforming the CDF $G(x)$ according to two sequential important classes. These classes, called Marshall and Olkin-G and Gamma-G, are important for modeling data in several areas, and they are reviewed below.

The CDF of the Marshall and Olkin's (1997) (MO-G) class (for $\theta > 0$) is

$$(1.1) \quad F_{\text{MO-G}}(x) = \frac{G(x)}{\theta + (1 - \theta)G(x)} = \frac{G(x)}{1 - (1 - \theta)[1 - G(x)]}, \quad x \in \mathbb{R}.$$

The density function corresponding to (1.1) has the form

$$(1.2) \quad f_{\text{MO-G}}(x) = \frac{\theta g(x)}{[\theta + (1 - \theta)G(x)]^2}.$$

For $\theta = 1$, $f_{\text{MO-G}}(x)$ is equal to $g(x)$. Equation (1.2) represents the PDF

of the minimum of n iid random variables having density $g(x)$, say T_1, \dots, T_N , where N has a geometric distribution with probability parameters θ and θ^{-1} if $0 < \theta < 1$ and $\theta > 1$, respectively.

Tahir and Nadarajah (2015, Table 2) presented thirty distributions belonging to this family. It is easily generated from the baseline quantile function (QF) by $Q_{\text{MO-G}}(u) = Q_G(\theta u / [\theta u + 1 - u])$ for $u \in (0, 1)$.

Marshall and Olkin considered the exponential and Weibull distributions for the baseline G and derived some structural properties of the generated distributions. If G is an exponential distribution, the special case refers to a two-parameter competitive model to the Weibull and gamma distributions.

The CDF of the gamma-G (Γ -G) class (Zografos and Balakrishnan, 2009) is

$$(1.3) \quad F_{\Gamma\text{-G}}(x) = \gamma_1(a, -\log[1 - G(x)]), \quad x \in \mathbb{R},$$

where $a > 0$ is an extra shape parameter, $\gamma_1(a, z) = \gamma(a, z)/\Gamma(a)$ is the incomplete gamma function ratio, and $\gamma(a, z) = \int_0^z t^{a-1} e^{-t} dt$.

Then, the PDF of the Γ -G class can be expressed as

$$(1.4) \quad f_{\Gamma\text{-G}}(x) = \frac{1}{\Gamma(a)} \{-\log[1 - G(x)]\}^{a-1} g(x).$$

Each new Γ -G distribution follows from a given baseline G. For $a = 1$, the Γ -G class reduces to G. If Z is a gamma random variable with unit scale and shape $a > 0$, then $W = Q_G(1 - e^{-Z})$ has density (1.4). So, the Γ -G distribution is easily generated from the gamma distribution and the QF of G.

The remaining of the paper is addressed as follows. Section 2 introduces the *Marshall and Olkin-Gamma-G* (MOGa-G) family, and provides some special models. The maximum likelihood estimates (MLEs) of its parameters is addressed in Section 3. Some simulations are performed in Section 4 to estimate the biases of the MLEs. Two empirical applications illustrate the potentiality of the proposed family in Section 5. A variety of theoretical properties are obtained in Section 6. Some conclusions remarks are offered in Section 7.

2. THE NEW FAMILY

By combining Equations (1.1) and (1.3), the CDF of the random variable $X \sim \text{MOGa-G}$ representing the new family has the form

$$(2.1) \quad F_X(x) = \frac{\gamma_1(a, -\log[1 - G(x)])}{\theta + (1 - \theta)\gamma_1(a, -\log[1 - G(x)])}, \quad x \in \mathbb{R}.$$

1 By differentiating (2.1), the PDF of X follows as

$$(2.2) \quad f_X(x) = \frac{\theta \{-\log[1 - G(x)]\}^{a-1} g(x)}{\Gamma(a) \{\theta + (1 - \theta)\gamma_1(a, -\log[1 - G(x)])\}^2}.$$

2 The density (2.2) can be interpreted from a sequence of N iid random
 3 variables, say Z_1, \dots, Z_N , each one having a gamma density with unit scale and
 4 shape $a > 0$, assuming that N (is not fixed) has a geometric distribution with
 5 probabilities θ and θ^{-1} for $0 < \theta < 1$ and $\theta > 1$, respectively. By transforming
 6 the Z_i 's via the baseline QF by $W_i = Q_G(1 - e^{-Z_i})$ (for $i = 1, \dots, N$), Equation
 7 (1.2) defines the PDF of the minimum W_1, \dots, W_n . The proposed family from
 8 the double composition of the two classes absorbs the impacts of their different
 9 flexibilities on real applications.

10 Table 1 provides some special cases of (2.2), where $\Phi(x)$ and $\phi(x)$ are the
 11 CDF and PDF of the standard normal distribution. The density and hazard
 12 ($h(x) = f(x)/[1 - F(x)]$) functions of the MOGa-Weibull (MOGa-W) model are
 13 displayed in Figure 1, which provide more flexibility to both functions for both
 14 classes applied separately to the Weibull model.




Figure 1: The density and hazard functions of the MOGa-W model.

The CDF (2.1) can be easily inverted to calculate the QF of the MOGa-G distribution, say $x = Q_X(u) = F_X^{-1}(u)$ (for $u \in (0, 1)$), in terms of the baseline QF $Q_G(\cdot)$. The inverse of $F_X(x) = u$, where u is a uniform number in $(0, 1)$, follows by combining the inverses of Equations (1.1) and (2.1). So, $F_X(x) = u$ gives $z = z(u) = \theta u / [1 - (1 - \theta)u]$ and $\gamma_1(a, -\log[1 - G(x)]) = z(u)$. Hence, the QF of X can be expressed as

$$x = Q_G(v(u)),$$

where

$$v(u) = 1 - \exp \left[-\gamma_1^{-1}(a, z(u)) \right],$$

- 1 and $\gamma_1^{-1}(a, w) = Q^{-1}(a, 1 - w)$ is the inverse function of $\gamma_1(a, w)$. Some formulae
 2 for $Q^{-1}(a, 1 - w)$ are given in [http://functions.wolfram.com/GammaBetaErf/](http://functions.wolfram.com/GammaBetaErf/InverseGammaRegularized/)
 3 [InverseGammaRegularized/](http://functions.wolfram.com/GammaBetaErf/InverseGammaRegularized/).

Table 1: Special Distributions in the MOGa-G family.

Distribution	Baseline CDF	Generated PDF
Normal	$G(x) = \Phi(x)$	$f_X(x) = \frac{\theta \{-\log[1-\Phi(x)]\}^{a-1} \phi(x)}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, -\log[1-\Phi(x)])\}^2}$
Logistic	$G(x) = \frac{1}{1+e^{-x}}$	$f_X(x) = \frac{\theta e^{-x} \{-\log[1-(1+e^{-x})^{-1}]\}^{a-1}}{\Gamma(a) (1+e^{-x})^2 \{\theta + (1-\theta)\gamma_1(a, -\log[1-(1+e^{-x})^{-1}])\}^2}$
Gumbel	$G(x) = 1 - \exp(-e^x)$	$f_X(x) = \frac{\theta \exp(ax - e^x)}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, e^x)\}^2}$
Log-Normal	$G(x) = \Phi(\log x)$	$f_X(x) = \frac{\theta \phi(\log x) \{-\log[1-\Phi(\log x)]\}^{a-1}}{\Gamma(a) x \{\theta + (1-\theta)\gamma_1(a, -\log[1-\Phi(\log x)])\}^2}$
Exponential	$G(x) = 1 - \exp(-\lambda x), \lambda > 0$	$f_X(x) = \frac{\theta \lambda^a x^{a-1}}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, \lambda x)\}^2}$
Weibull	$G(x) = 1 - \exp(-(\lambda x)^\gamma), \lambda, \gamma > 0$	$f_X(x) = \frac{\theta \gamma \lambda^a \gamma x^{a\gamma-1} \exp\{-(\lambda x)^\gamma\}}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, (\lambda x)^\gamma)\}^2}$
Gamma	$G(x) = \gamma_1(a, \beta x), a, \beta > 0$	$f_X(x) = \frac{\theta \beta^a x^{a-1} e^{-\beta x} \{-\log[1-\gamma_1(a, \beta x)]\}^{a-1}}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, -\log[1-\gamma_1(a, \beta x)])\}^2}$
Pareto	$G(x) = 1 - \frac{1}{(1+x)^\nu}, \nu > 0$	$f_X(x) = \frac{\theta e^{-x} [\nu \log(1+x)]^{a-1} g(x)}{\Gamma(a) (1+e^{-x})^2 \{\theta + (1-\theta)\gamma_1(a, \nu \log[1+x])\}^2}$
Dagum	$G(x) = [1 + (x/\beta)^{-\alpha}]^{-p}, \alpha, \beta, p > 0$	$f_X(x) = \frac{\theta \{-\log[1 - [1 + (x/\beta)^{-\alpha}]^{-p}]\}^{a-1} g(x)}{\Gamma(a) \{\theta + (1-\theta)\gamma_1(a, -\log(1 - ((x/\beta)^{-\alpha} + 1)^{-p}))\}^2}$

3. ESTIMATION

- 4 The MOGa-G family can be fitted to real data using the **AdequacyModel**
 5 package (Marinho et al., 2019) in the R software. This package does not require
 6 to define the log-likelihood function, and it computes the MLEs, their standard
 7 errors (SEs), and the formal statistics defined in Section 5. It is only necessary
 8 to provide the PDF and CDF of the distribution to be fitted to a data set.

For example, if x_i is one observation from (2.2) and $\boldsymbol{\eta}$ is a q -parameter vector specifying $G(\cdot)$ as the Weibull CDF, the log-likelihood function for $\boldsymbol{\theta}^\top = (a, \theta, \boldsymbol{\eta}^\top)$

from n observations can be expressed as

$$\begin{aligned} \ell(\boldsymbol{\theta}) = & n \log(\theta) + n \log(\gamma) + n a \gamma \log(\lambda) + (a\gamma - 1) \sum_{i=1}^n \log(x_i) - \lambda^\gamma \sum_{i=1}^n \log(x_i) \\ (3.1) \quad & - n \log[\Gamma(a)] - 2 \sum_{i=1}^n \log\{\theta + (1 - \theta)\gamma_1[a, (\lambda x_i)^\gamma]\}. \end{aligned}$$

Due to the impossibility of obtaining the MLEs in closed form, numerical methods to calculate the estimates that maximize $\ell(\cdot)$ are necessary. Several programming languages and statistical software provides functions and routines that make it easy to obtain numerical estimates by various interactive methods. In practice, these estimates are commonly found in this way, since the Newton and quasi-Newton methods produce satisfactory results under reasonable conditions of the object function, i.e., when they do not impose restrictions that disturb the convergence of the algorithms.

The **AdequacyModel** package of the programming language R is used to obtain the MLEs, see R Core Team (2020). This library, created and maintained by one of the authors of this paper, is widely cited by several works in statistics, and serves as a basis for other library implementations available on the Comprehensive R Archive Network (CRAN). By using the `goodness.fit` function, it is possible to provide an implementation of (2.2), and obtain $\ell(\cdot)$ by returning the MLEs, and some measures of adequacy of fit. Further details regarding this package can be obtained from Marinho et al. (2019).

4. SIMULATIONS

Due to the probable absence of MLEs in closed-form for distributions belonging to the MOGa-G family, it is necessary to examine the precision of the estimates calculated numerically.

In order to do that, the biases of the estimators of the parameters of the MOGa-Dagum($\theta, a, \alpha, \beta, p$) and MOGa-Weibull($\theta, a, \lambda, \gamma$) distributions are determined, where $G \sim \text{Dagum}(\alpha, \beta, p)$ and $\text{Weibull}(\gamma, \lambda)$ are the baseline models, respectively. All parameters are taken equal to one for different sample sizes reported in Tables 2 and 3.

The numbers in Tables 2 and 3 indicate that the estimation method behaves well when the sample size increases. This is theoretically expected. However, in practice, difficulties can be faced to other families due to the flatness of the log-likelihood function.

All Monte Carlo simulations can be reproduced using the script in <https://github.com/prdm0/MOGG>. The simulations are parallelized and able to use all

1 threads available by a multicore processor, thus making them more computatio-
 2 nally efficient, and consequently requiring less time to complete.

3 The simulations are performed on a computer with an Intel(R) Core(TM) i5-
 4 9500 CPU processor with 6 threads working at a maximum frequency of 3.00GHz,
 5 requiring, on this hardware, a time of 15.4828 hours to perform all simulations,
 6 7.7414 hours for the MOGa-Dagum($\theta, a, \alpha, \beta, p$) distribution, and 4.9688 hours for
 7 the MOGa-Weibull($\theta, a, \lambda, \gamma$) distribution. Tables 2 and 3 reveal that the average
 8 biases of the MLEs could be very small for $n > 2,000$.

9 To generate observations from the random variable X with density f , the
 10 well-known Acceptance-Rejection Algorithm for continuous random variables is
 11 very useful when the QF involves complex functions that can lead to some numer-
 12 ical inaccuracies. For doing this, another random variable Y is chosen such that
 13 it can generate observations from a PDF h with the same support as f . Then,
 14 the acceptance and rejection algorithm is defined by the following steps:

- 15 1. Generate an outcome y from Y ;
- 16 2. Generate an observation u from a random variable $U \sim \mathcal{U}(0, 1)$;
- 17 3. If $u < \frac{f(y)}{cg(y)}$, where c is a real constant, accept $x = y$; otherwise reject y as
 18 an outcome from X and return to 1.

19 The constant c must be chosen in such a way that $\frac{f(y)}{cg(y)} \leq 1$. Thus, to
 20 minimize the computational cost of generating observations from X through the
 21 generated observations from Y , c is chosen as the lowest possible value to maximize
 22 the likelihood of acceptance. Further details of this method can be found in Rizzo
 23 (2019).

Table 2: Average biases of the MLEs of the MOGa-Dagum($\theta, a, \alpha, \beta, p$) distribu-
 tion calculated by the BFGS method from simulations.

n	$B(\hat{\theta})$	$B(\hat{a})$	$B(\hat{\alpha})$	$B(\hat{\beta})$	$B(\hat{p})$	Time (mins)
10	0.2213	2.1944	2.6971	1.5803	1.3190	0.6960
20	0.4240	2.4793	1.5591	1.8083	0.7414	0.9819
60	0.7458	2.2661	0.5598	1.8495	0.2812	1.9417
100	0.6194	1.9438	0.3312	1.6142	0.2935	2.7208
200	0.3950	1.4262	0.1856	1.1556	0.3611	4.4534
400	0.2077	0.9599	0.1082	0.6157	0.4076	7.4698
600	0.1200	0.7213	0.0767	0.4024	0.3572	9.4975
1000	0.0629	0.4791	0.0503	0.2123	0.2584	12.4221
2000	0.0362	0.2958	0.0298	0.1145	0.1878	20.7251
5000	-0.0040	0.1325	0.0159	0.0144	0.0167	28.3380
10000	-0.0133	0.0815	0.0096	0.0081	0.0039	50.9298
20000	-0.0111	0.0349	0.0037	0.0006	-0.0109	68.6320
30000	-0.0036	0.0191	0.0006	-0.0041	-0.0034	97.3046
50000	-0.0057	0.0129	0.0016	0.0015	-0.0026	158.3737

Table 3: Average biases of the MLEs of the MOGa-Weibull($\theta, a, \lambda, \gamma$) distribution calculated by the BFGS method from simulations.

n	$B(\hat{\theta})$	$B(\hat{a})$	$B(\hat{\lambda})$	$B(\hat{\gamma})$	Time (mins)
10	0.0818	0.1362	4.9274	1.2407	0.6716
20	0.3404	-0.0177	3.4160	1.4117	0.8077
60	0.7037	-0.0677	1.8806	1.3385	1.1773
100	0.6698	-0.0535	1.3684	1.1796	1.2643
200	0.5371	-0.0299	0.8265	0.9110	1.7886
400	0.3371	-0.0047	0.4205	0.5967	2.8386
600	0.2457	0.0076	0.2685	0.4306	3.6867
1000	0.1476	0.0093	0.1553	0.2818	5.0944
2000	0.0731	0.0035	0.0758	0.1530	8.8577
5000	0.0264	0.0007	0.0283	0.0618	15.8586
10000	0.0128	-0.0007	0.0142	0.0318	29.8629
20000	0.0053	-0.0012	0.0071	0.0160	48.7417
30000	0.0023	-0.0014	0.0053	0.0119	64.7387
50000	0.0023	-0.0004	0.0028	0.0063	112.7422

5. APPLICATIONS

Two applications compare the MOGa-Weibull (MOGa-W for short) model with seven extended Weibull distributions: the beta-Weibull (β -W) (Famoye et al., 2005), Kumaraswamy Weibull (Kw-W) (Cordeiro et al., 2010), Marshall-Olkin Weibull (MO-W) (Ahmed et al., 2017), Marshall-Olkin Extended Weibull (MOE-W) (Cordeiro et al., 2019), exponentiated Weibull (exp-W) (Mudholkar and Srivastava, 1993), gamma Weibull (Γ -W) (Cordeiro et al., 2016), and exponentiated generalized Weibull (EG-W) (Oguntunde et al., 2015) (with $a = 1$). Some of these distributions are widely used in practice.

The log-likelihood for $\boldsymbol{\theta}$ from the MOGa-W distribution from one observation can be expressed as

$$\begin{aligned} \ell(\boldsymbol{\theta}) = & \log(\theta) + \log(\gamma) + (a\gamma) \log(\lambda) + (a\gamma - 1) \log(x) - (\gamma x)^\gamma - \log[\Gamma(a)] \\ & - 2 \log\{\theta + (1 - \theta)\gamma_1[a, (\lambda x)^\gamma]\}, \end{aligned} \quad (5.1)$$

where $\boldsymbol{\theta} = (a, \theta, \lambda, \gamma)^\top$. The components of the score function are

$$U_a(\boldsymbol{\theta}) = \gamma \log(\lambda) + \gamma \log(x) - \psi^{(0)}(a) - \frac{2 \{ (1 - \theta)A - (1 - \theta)\psi^{(0)}(a)\gamma_1[a, (x\lambda)^\gamma] \}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (\lambda x)^\gamma]},$$

$$U_\theta(\boldsymbol{\theta}) = \frac{1}{\theta} - \frac{2 \{ \Gamma(a) - \gamma_1[a, (\lambda x)^\gamma] \}}{\theta \Gamma(a) + (1 - \theta)\gamma_1[a, (\lambda x)^\gamma]},$$

$$U_\lambda(\boldsymbol{\theta}) = \frac{\gamma}{\lambda} [a - (\lambda x)^\gamma] + \frac{2\gamma \lambda^{-1} (\lambda x)^{a\gamma} (1 - \theta) \exp\{-(\lambda x)^\gamma\}}{\theta \Gamma(a) + (1 - \theta) \gamma_1 [a, (\lambda x)^\gamma]}$$

and

$$U_\gamma(\boldsymbol{\theta}) = \frac{1}{\gamma} + a \log(\lambda) + a \log(x) - (\lambda x)^\gamma \log(\lambda x) + \frac{2(1 - \theta)(\lambda x)^{a\gamma} \log(\lambda x) \exp\{-(\lambda x)^\gamma\}}{\theta \Gamma(a) + (1 - \theta) \gamma_1 [a, (\lambda x)^\gamma]},$$

where $\psi^{(n)}(x)$ is the n th derivative of the digamma function,

$$A = \log [(\lambda x)^\gamma] \gamma_1 [a, (\lambda x)^\gamma] + G_{2,3}^{3,0} \left((\lambda x)^\gamma \middle| \begin{matrix} 1, 1 \\ 0, 0, a \end{matrix} \right),$$

1 and $G_{p,q}^{m,n} \left(z \middle| \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \right)$ is the Meijer G function.

2 The **AdequacyModel** package is used to fit the previous distributions to
 3 two real data sets. The SANN method, which is a variant of simulated annealing
 4 algorithm (Belisle, 1992), is used here. The distributions are compared via the
 5 Anderson Darling (A*) and Cramér Von Mises (W*) statistics reported in the
 6 `goodness.fit` function.

7 For the first case, the **betareg** package is applied to a modification of the
 8 “FoodExpenditure” data, which refer to the proportions of income spent on food
 9 for 38 households in a large US city (according to the package information). The
 10 household expenditures for food are given by

$$data = FoodExpenditure_{food} / \#(FoodExpenditure_{food}),$$

11 where $FoodExpenditure_{food}$ is the random variable corresponding to the house-
 12 hold expenditures for food, and $\#(\cdot)$ indicates the number of observations on this
 13 variable. The observations for the first data set are given below:

14 0.4210000, 0.4382105, 0.5721316, 0.1955526, 0.2758158, 0.3565263, 0.6120000,
 15 0.4730526, 0.3726579, 0.2322368, 0.3732632, 0.5158947, 0.3612632, 0.5563421,
 16 0.4591053, 0.2533947, 0.3685526, 0.2410526, 0.4955526, 0.2010789, 0.3653158,
 17 0.2544737, 0.5685263, 0.2859474, 0.7626316, 0.2863684, 0.4884474, 0.3060263,
 18 0.4754474, 0.3826053, 0.5050526, 0.6820526, 0.7587632, 0.4176053, 0.3923684,
 19 0.2513158, 0.6070000, 0.3881842.

20 Some descriptive statistics are reported in Table 4. The minimum value
 21 refers to a family of 3 people and income of 39,151, which does not represent the
 22 lowest family income for the current data, as expected, occupying only the fifth
 23 position among those with the lowest income. The maximum value corresponds to
 24 a family of 6 people with an income of 69,929, the second-largest number among
 25 the number of people per family in the group in question. Furthermore, we can
 26 note that the current data present positive asymmetry and negative kurtosis.

27 In addition, the standard deviation is relatively low. Figure 2 displays the
 28 total time on test (TTT) plot for the first data set, which shows that the failure

1 function is decreasing. So, the MOGa-W distribution is appropriate to fit these
 2 data, since its hazard function presents this shape (see, Figure 1b).

Table 4: Descriptive statistics for the food data.

Minimum	0.1956
1st Qu.	0.2913
Median	0.3903
Mean	0.4198
3rd Qu.	0.5027
Maximum	0.7626
Standard Deviation	0.1480
Skewness	0.5250
Kurtosis	-0.4440




Figure 2: TTT plot for the food data.

3 The MLEs, their standard errors (SEs) (in parentheses), and the statistics
 4 W^* and A^* for the fitted models to the current data are listed in Table 5. The
 5 results indicate that the proposed model has better performance than the other
 6 seven fitted models.

Table 5: Estimation results for food data.

Model	\hat{a}	$\hat{\theta}$	$\hat{\lambda}$	$\hat{\gamma}$	W^*	A^*
MOGa-W($a, \theta, \lambda, \gamma$)	0.9261 (0.0262)	1.3796 (0.2238)	33.3230 (0.2853)	25.3988 (0.0825)	0.0339	0.2376
β -W($a, \theta, \lambda, \gamma$)	9.9288 (0.0290)	0.1700 (0.0204)	9.7594 (<0.0001)	1.5305 (<0.0001)	0.0435	0.2594
KW-W($a, \theta, \lambda, \gamma$)	0.0498 (0.0080)	99.9998 (16.2259)	1.0760 (0.0031)	23.4028 (0.0146)	1.3309	6.7426
MOE-W($a, \theta, \lambda, \gamma$)	0.1366 (0.1599)	2.0204 (<0.0001)	62.7220 (<0.0001)	4.2956 (0.7365)	0.0354	0.2579
EGW(a, b, λ, γ)	5.6189 (0.0028)	6.1833 (0.0009)	1.2870 (0.1159)	1.3798 (0.1480)	0.0371	0.2518
MO-W(a, λ, γ)	0.1592 (0.0717)	- (-)	1.5860 (0.1335)	4.2671 (0.1665)	0.0345	0.2573
exp-W(a, λ, γ)	6.1102 (0.4222)	- (-)	4.4680 (0.3175)	1.3858 (0.1677)	0.0372	0.2523
γ -W(a, λ, γ)	5.7515 (0.0015)	- (-)	10.0000 (0.0001)	1.2087 (0.0082)	0.0879	0.6599

1 A data set collected in a pilot study about hypertension in the Dominican
 2 Republic in 1997 refers to the second application. The observations are the systolic
 3 blood pressure of persons who came to medical clinics in several villages for a
 4 variety of complaints. **The observations for the data set in question are:** 150, 120,
 5 120, 180, 138, 115, 130, 150, 200, 120, 190, 90, 130, 120, 200, 140, 110, 134, 160,
 6 140, 105, 126, 129, 120, 100, 130, 118, 144, 180, 138, 110, 140, 120, 118, 110, 110,
 7 130, 140, 130, 165, 180, 130, 140, 112, 130, 158, 112, 150, 140, 142, 110, 140, 130,
 8 132, 140, 140, 122, 128, 90, 118, 120, 110, 122, 200, 110, 140, 150, 120, 150, 120,
 9 164, 122, 112, 130, 140, 102, 122, 130, 102, 130, 122, 200, 140, 180, 124, 110, 124,
 10 90, 120, 159, 142, 140, 118, 122, 108, 170, 120, 140, 100, 118, 110, 114, 150, 160,
 11 140, 190, 118, 120, 150, 120, 200, 150, 168, 110, 142, 150, 160, 142, 160, 150, 110,
 12 128, 122, 150, 140, 122, 120, 130, 100, 130, 150, 130, 100, 120, 105, 100, 150, 196,
 13 130, 110, 140, 122, 110, 164, 120, 120, 150, 160, 150, 135, 124, 110, 100, 95, 130,
 14 120, 108, 118, 170, 105, 120, 95, 95, 120, 140, 142, 160, 110, 190, 180, 130, 130,
 15 120, 204, 150, 150, 120, 122, 120, 130, 140, 148, 118, 126, 136, 140, 130, 102, 110,
 16 110, 130, 126, 142, 140, 128, 130, 124, 162, 130, 130, 110, 80, 166, 140, 160, 160,
 17 140, 98, 138, 120, 112, 112, 134, 140, 115, 140, 98, 115, 120, 80, 160, 126, 110,
 18 130, 104, 236, 118, 120, 140, 120, 98, 164, 150, 110, 120, 130, 170, 180, 110, 120,
 19 130, 118, 130, 190, 158, 90, 99, 210, 180, 140, 184, 105, 120, 150, 140, 130, 160,
 20 118, 210, 100, 170, 150, 130, 170, 150, 120, 134, 90, 125, 170, 140, 150, 110, 105,
 21 140, 120, 100, 124, 112, 160, 140, 118, 190, 110, 118, 160, 150, 124, 128, 150, 120,
 22 125, 118, 132, 110, 143, 170, 98, 124, 180, 178, 110, 98, 159, 110, 140, 130, 122,

1 110, 98, 180, 90, 118, 165, 138, 138, 170, 106, 170, 140, 90, 118, 110, 102, 102,
 2 180, 100, 110, 162, 140, 110, 98, 140, 140, 110, 170, 112, 90, 102, 106, 124, 110,
 3 180, 138, 90, 150, 126, 110, 130, 150, 145, 140, 156, 110, 150, 160, 120, 140, 120,
 4 110, 120, 140, 160, 160, 110, 150, 118, 110, 120, 120, 146, 124, 170, 124, 170, 159,
 5 120, 120, 118, 152, 190.

6 Table 6 provides the descriptive statistics for the second data set. Consi-
 7 dering that the systolic blood pressure represents the highest number presented in
 8 the pressure measuring equipment, the maximum (236) found in this table should
 9 represent an individual with serious heart problems. This is due to the fact that
 10 the normal systolic pressure is 120.

11 The minimum value (80) should be for an individual who probably suffers
 12 from low blood pressure. Also note that the data set has positive skewness. This
 13 indicates that few people have high pressure values and, in this case, the mode is
 14 120, which represents a good value for systolic pressure.

15 Figure 3 provides the TTT plot for the second clinic data, which shows that
 16 the hazard function is decreasing, thus supporting the MOGa-W model.




Figure 3: TTT plot for the clinic data.

Table 6: Descriptive statistics for the clinic data.

Minimum	80
1st Qu.	118
Median	130
Mean	133
3rd Qu.	150
Maximum	236
Standard Deviation	25.7157
Skewness	0.7893
Kurtosis	0.5908

1 The MLEs of the parameters, their SEs and the values of the adequacy
2 measures for the fitted models to the clinic data are reported in Table 7. By
3 comparing the measure values, the proposed distribution outperforms all other
4 fitted models.

Table 7: Estimation results for clinic data.

Model	\hat{a}	$\hat{\theta}$	$\hat{\lambda}$	$\hat{\gamma}$	W^*	A^*
MOGa-W($a, \theta, \lambda, \gamma$)	9.6293 (0.0062)	3.6407 (0.1824)	6.2608 (0.0244)	12.8234 (0.0079)	0.5093	2.8076
β -W($a, \theta, \lambda, \gamma$)	31.0847 (0.0127)	47.1463 (<0.0001)	0.0169 (0.0001)	2.0540 (<0.0001)	0.7540	4.2794
KW-W($a, \theta, \lambda, \gamma$)	7363.2810 (0.0419)	0.0392 (0.0004)	1.4676 (<0.0001)	0.6146 (<0.0001)	0.5351	2.9617
MOE-W($a, \theta, \lambda, \gamma$)	101.1347 (46.7828)	0.4238 (0.1008)	0.0309 (0.0036)	1.7024 (0.1681)	1.1441	6.6446
EGW(a, b, λ, γ)	0.2351 (0.0025)	140.0000 (3.2785)	0.4576 (<0.0001)	0.7425 (<0.0001)	0.8925	5.3338
MO-W(a, λ, γ)	173.2139 (0.0001)	- (-)	0.0212 (0.0002)	1.6019 (0.0001)	1.4760	8.5870
exp-W(a, λ, γ)	69.0291 (0.0838)	- (-)	0.0240 (0.0002)	1.3455 (<0.0001)	0.8899	5.0941
Γ -W(a, λ, γ)	9.1122 (0.9633)	- (-)	0.0261 (0.0029)	1.7464 (0.0803)	0.6227	3.4882

6. MATHEMATICAL PROPERTIES

5 Here, some mathematical properties for the MOGa-G family are presented
6 based on a linear representation for its density function in terms of “exponentiated-

1 G'' (exp-G) densities.

6.1. Linear Representation

2 For an arbitrary CDF $G(x)$, the CDF and PDF of the exp-G distribution
 3 with power parameter $a > 0$ are

$$\Pi_a(x) = G(x)^a \quad \text{and} \quad \pi_a(x) = a g(x) G(x)^{a-1},$$

4 respectively. This class of distributions is quite useful in several applications. In
 5 fact, Tahir and Nadarajah (2015) cited more than seventy papers on exponenti-
 6 ated distributions in their Table 1.

7 First, the CDF of the MO-G distribution (1.2) admits the linear represen-
 8 tation (Barreto-Souza et al., 2013)

$$(6.1) \quad F_{\text{MO}-\Gamma}(x) = \sum_{i=0}^{\infty} w_i^{\text{MO-G}} \Pi_{i+1}(x) = \sum_{i=0}^{\infty} w_i^{\text{MO-G}} G(x)^{i+1},$$

9 where the coefficients are (for $i = 0, 1, \dots$)

$$w_i^{\text{MO}-\Gamma} = w_i^{\text{MO}-\Gamma}(\theta) = \begin{cases} \frac{(-1)^i \theta}{(i+1)} \sum_{j=i}^{\infty} (j+1) \binom{j}{i} \bar{\theta}^j, & \theta \in (0, 1), \\ \theta^{-1} (1 - \theta^{-1})^i, & \theta > 1, \end{cases}$$

10 and $\bar{\theta} = 1 - \theta$.

11 Second, the linear combination for the Γ -G cumulative distribution (1.4)
 12 follows from Castellares and Lemonte (2015) as

$$(6.2) \quad F_{\Gamma\text{-G}}(x) = \sum_{j=0}^{\infty} w_j^{\Gamma\text{-G}} \Pi_{a+j}(x).$$

Here,

$$w_j^{\Gamma\text{-G}} = w_j^{\Gamma\text{-G}}(a) = \frac{\varphi_j(a)}{(a+j)},$$

$$\varphi_0(a) = \frac{1}{\Gamma(a)}, \quad \varphi_j(a) = \frac{(a-1)}{\Gamma(a)} \psi_{j-1}(j+a-2), \quad j \geq 1,$$

and

$$\begin{aligned} \psi_{n-1}(x) = \frac{(-1)^{n-1}}{(n+1)!} & \left[H_n^{n-1} - \frac{x+2}{n+2} H_n^{n-2} + \frac{(x+2)(x+3)}{(n+2)(n+3)} H_n^{n-3} - \dots \right. \\ & \left. + (-1)^{n-1} \frac{(x+2)(x+3) \cdots (x+n)}{(n+2)(n+3) \cdots (2n)} H_n^0 \right], \end{aligned}$$

1 is the Stirling polynomial, $H_{n+1}^m = (2n+1-m)H_n^m + (n-m+1)H_n^{m-1}$ is a
 2 positive integer, $H_0^0 = 1$, $H_{n+1}^0 = 1 \times 3 \times 5 \times \cdots \times (2n+1)$ and $H_{n+1}^n = 1$.

By inserting (6.2) in Equation (6.1) and via a result for a power series raised to a positive integer (Gradshteyn and Ryzhik, 2000), the expansion for the cdf of the MOGa-G distribution reduces to

$$\begin{aligned} F_{\text{MO}-\Gamma-\text{G}}(x) &= \sum_{i=0}^{\infty} w_i^{\text{MO}-\Gamma} G(x)^{(i+1)a} \left[\sum_{j=0}^{\infty} w_j^{\Gamma-\text{G}} G(x)^j \right]^{i+1} \\ &= \sum_{i=0}^{\infty} w_i^{\text{MO}-\Gamma} G(x)^{(i+1)a} \sum_{j=0}^{\infty} c_{i+1,j} G(x)^j = \sum_{i,j=0}^{\infty} d_{i,j} \Pi_{(i+1)a+j}(x), \end{aligned}$$

3 where $d_{i,j} = d_{i,j}(a, \theta) = w_i^{\text{MO}-\Gamma} c_{i+1,j}(a)$, $c_{i+1,0}(a) = (w_0^{\Gamma-\text{G}})^{i+1}$ and, for $m \geq 1$,
 4 $c_{i+1,m}(a) = \frac{1}{mw_0^{\Gamma-\text{G}}} \sum_{r=1}^m [r(i+2) - m] w_r^{\Gamma-\text{G}} c_{i+1,m-r}(a)$.

5 By differentiating the last equation, the linear representation for the MOGa-
 6 G density holds

$$(6.3) \quad f_{\text{MO}-\Gamma-\text{G}}(x) = \sum_{i,j=0}^{\infty} d_{i,j} \pi_{(i+1)a+j}(x).$$

7 So, some structural properties of the proposed family can be determined
 8 from the double linear combination (6.3) and those properties of the exp-G distri-
 9 bution. In most applications, the indices i and j can vary up to a small number
 10 of terms.

6.2. Some quantities

11 Hereafter, let $T_{i,j} \sim \text{exp-G}[(i+1)a + j]$. The n th moment of X can be
 12 determined from (6.3) as

$$(6.4) \quad \mu'_n = E(X^n) = \sum_{i,j=0}^{\infty} d_{i,j} E(T_{i,j}) = \sum_{i,j=0}^{\infty} [(i+1)a + j - 1] d_{i,j} \tau[n, (i+1)a + j - 1],$$

13 where

$$\tau(n, a) = \int_{-\infty}^{\infty} x^n G(x)^a g(x) dx = \int_0^1 Q_G(u)^n u^a du.$$

14 Expressions for moments of several exponentiated distributions can be found
 15 in the papers cited in Tahir and Nadarajah (2015, Table 1). We give just one
 16 example from Equation (6.4) by taking the exponential distribution with rate
 17 $\lambda > 0$ for the baseline G. It follows easily as

$$\mu'_n = n! \lambda^n \sum_{i,j,m=0}^{\infty} \frac{(-1)^{n+m} [(i+1)a + j] d_{i,j}}{(m+1)^{n+1}} \binom{(i+1)a + j - 1}{m}.$$

For empirical purposes, the shape of many distributions can be usefully described by the incomplete moments. These moments play an important role for measuring inequality. For example, the mean deviations and Lorenz and Bonferroni curves depend upon the first incomplete moment of the distribution. The n th incomplete moment of X can be expressed as

$$(6.5) \quad m_n(y) = \int_{-\infty}^y x^n f_X(x) dx = \sum_{i,j=0}^{\infty} [(i+1)a+j] d_{i,j} \int_0^{G(y)} Q_G(u)^n u^{(i+1)a+j-1} du.$$

The definite integral in (6.5) can be evaluated for most baseline G distributions.

The moment generating function (MGF) $M(t) = E(e^{tX})$ of X can be expressed from (6.3)

$$(6.6) \quad M(t) = \sum_{i,j=0}^{\infty} d_{i,j} M_{i,j}(t) = \sum_{i,j=0}^{\infty} [(i+1)a+j] d_{i,j} \rho(t, (i+1)a+j-1),$$

where $M_{i,j}(t)$ is the MGF of $Y_{i,j}$ and

$$\rho(t, a) = \int_{-\infty}^{\infty} e^{tx} G(x)^a g(x) dx = \int_0^1 \exp\{t Q_G(u)\} u^a du.$$

The MGFs of several MOGa-G distributions can be determined from Equation (6.6). For example, the generating function of the MOGa-exponential with parameter λ (if $t < \lambda^{-1}$) is

$$M(t) = \sum_{i,j=0}^{\infty} [(i+1)a+j] d_{i,j} B((i+1)a+j, 1-\lambda t).$$

7. CONCLUSIONS

A new family of distributions called the Marshall and Olkin-Gamma-G family with two shape parameters is introduced. The estimation of the unknown parameters is done via the maximum likelihood method and a simulation study is conducted to verify its adequacy. Additionally, the usefulness of the proposed family is shown empirically by means of two applications to real data. In fact, the new family can generate very competitive distributions with the same number of parameters than others constructed by existing classes.

REFERENCES

- [1] M.H. Tahir, S. Nadarajah: Parameter induction in continuous univariate distributions: Well-established G families. *Anais da Academia Brasileira de Ciências* **87** 539-568 (2015).

- 1 [2] A.W. Marshall, I. Olkin: A new method for adding a parameter to a family of
2 distributions with application to the exponential and Weibull families. *Biometrika*
3 **84** 641-652 (1997).
- 4 [3] K. Zografos, N. Balakrishnan: On families of beta-and generalized gamma-
5 generated distribution and associate inference. *Statistical Methodology* **6** 344-362
6 (2009).
- 7 [4] N. Eugene, C. Lee, F. Famoye: Beta-normal distribution and its applications.
8 *Communications in Statistics – Theory and Methods* **31** 497-512 (2002).
- 9 [5] P. R. D. Marinho, R. B. Silva, M. Bourguignon, G. M. Cordeiro, S. Nadarajah.
10 *AdequacyModel: An R package for probability distributions and general purpose*
11 *optimization. PloS one*, **14**, e0221487 (2019).
- 12 [6] R Core Team. R: A language and environment for statistical computing. R Founda-
13 tion for Statistical Computing, Vienna, Austria. URL [https://www.R-project.](https://www.R-project.org/)
14 [org/](https://www.R-project.org/) (2022).
- 15 [7] F. Famoye, C. Lee and O. Olumolade: The beta-Weibull distribution. *Journal of*
16 *Statistical Theory and Applications* **4** 121-136 (2005).
- 17 [8] G.M. Cordeiro, E. M. M. Ortega and S. Nadarajah: The Kumaraswamy Weibull
18 distribution with application to failure data **347** [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.jfranklin.2010.06.010)
19 [jfranklin.2010.06.010](https://doi.org/10.1016/j.jfranklin.2010.06.010) (2010).
- 20 [9] H. H. Ahmed, O. M. Bdair and M. Ahsanullah: On Marshall-Olkin Extended
21 Weibull Distribution **16** 1-17 (2017).
- 22 [10] G. S. Mudholkar and D. K. Srivastava: Exponentiated Weibull Family for An-
23 alyzing Bathtub Failure-Rate Data *IEEE Transactions on Reliability* **42** 299-302
24 <http://dx.doi.org/10.1109/24.229504> (1993)
- 25 [11] G. M. Cordeiro, M. C. S. Lima, A. E. Gomes, C. Q. da Silva and E. M. M. Ortega:
26 The gamma extended Weibull distribution *Journal of Statistical Distributions and*
27 *Applications Berlin* **3** 1-19 (2016).
- 28 [12] P. Oguntunde, O. Odetunmbi and A. O. Adejumo On the Exponentiated Gener-
29 alized Weibull Distribution: A Generalization of the Weibull Distribution *Indian*
30 *Journal of Science and Technology* **8** DOI: 10.17485/ijst/2015/v8i35/67611 (2015).
- 31 [13] C. J. P. Belisle: Convergence theorems for a class of simulated annealing algo-
32 rithms on *Rd J Applied Probability* **29** 885-895 1992.
- 33 [14] W. Barreto-Souza, A.J. Lemonte, G.M. Cordeiro: General results for the Marshall
34 and Olkin’s family of distributions. *Anais da Academia Brasileira de Ciências* **85**
35 3-21 (2013).
- 36 [15] F. Castellares, A.J. Lemonte: A new generalized Weibull distribution generated
37 by gamma random variables. *Journal of the Egyptian Mathematical Society* **23**
38 382-390 (2015).
- 39 [16] Gradshteyn IS, Ryzhik IM: Table of Integrals, Series and Products. Sixth edition.
40 San Diego: Academic Press (2000).
- 41 [17] Rizzo, Maria L. Statistical computing with R. CRC Press (2019).