


Solutions: Midterm 2 Practice Problems

Problem 1

In the following three parts, choose the figure with the appropriate regression line(s).

Part A


Choose the figure with the solid line that represents the simple regression line ($y \sim x$).



A3

Part B


Choose the figure with the two dashed lines associated with the additive multiple regression ($y \sim x + z$).



B1

Part C

Choose the figure with the two dotted lines associated with the interactive multiple regression ($y \sim x * z$).



C2

Problem 2

Suppose we have a sample $Y_1 \dots Y_n$ stored on an $\mathbf{R}$ vector Y of length n . We run this code.

```

B=10000
Y.bar = rep(NA,B)
for(b in 1:B) {
  Y.bar[b]= mean(sample(Y,size=n,replace=TRUE))
}

```




Figure 1: Histogram of the vector Y's elements Figure 2: Histogram of the Y.bar's elements

Part a

What does this code calculate? State it in words and mark it with an 'a', and an arrow or bracket if necessary for clarity, on the appropriate plot above.

```
mean(Y)
```

The sample mean. It's indicated by the solid vertical line on the left histogram.

Part b

What does this code calculate? State it in words and mark it with a 'b', and an arrow or bracket if necessary for clarity, on the appropriate plot above.

```
sd(Y)
```

The sample standard deviation. It's the distance between the solid vertical and either dashed vertical line on the left histogram.

Part c

What does this code calculate? State it in words and mark it with an ‘c’, and an arrow or bracket if necessary for clarity, on the appropriate plot above.

```
sd(Y.bar)
```

The bootstrap sampling distribution’s standard deviation. It’s the distance between the solid vertical and either dashed vertical line on the left histogram.

Part d

What does this code calculate? State it in words and mark it with an ‘d’, and an arrow or bracket if necessary for clarity, on the appropriate plot above.

```
mean(Y) - 1.96* sd(Y)/sqrt(n)
mean(Y) + 1.96* sd(Y)/sqrt(n)
```

A 95% confidence interval — based on normal approximation — for the mean of the population our sample $Y_1 \dots Y_n$ is drawn from. This is indicated by the horizontal segment (with a dot in the middle) on the right histogram.

Part e

How could you calculate roughly the same thing as in part d using ‘sd(Y.bar)’ and not ‘sd(Y)’?

We can use the standard deviation of the bootstrap sampling distribution, rather than ‘sd(Y)/sqrt(n)’, as our estimate of standard error.

```
mean(Y) - 1.96 * sd(Y.bar)
mean(Y) + 1.96 * sd(Y.bar)
```

Problem 3

Consider the following R code:

```
1. lm(residuals(lm(y~z)) ~ residuals(lm(z~x)))
2. lm(y ~ residuals(lm(z~x)))
3. lm(residuals(lm(y~z)) ~ residuals(lm(x~z)))
4. lm(residuals(lm(y~z)) ~ x)
5. lm(residuals(lm(y~x)) ~ residuals(lm(x~z)))
6. lm(residuals(lm(y~x)) ~ z)
7. lm(residuals(lm(y~x)) ~ residuals(lm(z~x)))
8. lm(y ~ residuals(lm(x~z)))
```

- a) Which one or more of the above will produce the coefficient on x from the multiple regression of $y \sim x + z$?

3 and 8

- b) Which one or more of the above will produce the same residuals as the residuals from the multiple regression of $y \sim x + z$?


3 and 7

Problem 4

Y_i is a individual i 's income and X_i is that individual's number of years of education. We take a random sample of individuals and record their years of education and income. Define $\hat{\mu}(16)$ as the subsample mean income among those with 16 years of education. Define an indicator function for $x = 16$ and show that $\hat{\mu}(16)$ is an unbiased estimator of $\mu(16)$, the subpopulation mean income for those with 16 years of education. Justify your steps.

Problem 5

In the below visualization, the x-axis is an individual's years of education, the y-axis is income, the red line is the least squares line, and the black dots indicate the mean income within each year of education. You have three options to summarize the change in income across an additional year of education: the average over years, the average weighted by the proportion of individuals with the years of education P_x , and the slope of the least squares line. Rank these options smallest to largest regarding the magnitude of the estimate you expect and explain your reasoning.



Problem 6

Suppose we want γ_1 from $y_i = \gamma_0 + \gamma_1 x_i + \gamma_2 w_i + e_i$ but we run the regression $lm(y \sim x)$ without w . We have de-meaned our variables x_i and w_i such that $\bar{x} = 0$ and $\bar{w} = 0$. Will the coefficient on x from $lm(y \sim x)$ be equal to γ_1 ? Derive what your coefficient on x will be in terms of γ s and justify your steps.