

Applied Analytics and Predictive Modeling

Spring 2020

Lecture-4

Lydia Manikonda
manikl@rpi.edu



Rensselaer

Today's agenda

- Class Presentations – Groups 1, 2 and 10
- Data Quality
- Linear Regression

Presentations of Case Study-1

Data Quality

- Poor data quality negatively affects many data processing efforts

“The most important point is that poor data quality is an unfolding disaster.

- Poor data quality costs the typical company at least ten percent (10%) of revenue; twenty percent (20%) is probably a better estimate.”

Thomas C. Redman, DM Review, August 2004


- Data mining example: a classification model for detecting people who are loan risks is built using poor data
 - Some credit-worthy candidates are denied loans
 - More loans are given to individuals that default

Data Quality ...


- What kinds of data quality problems?
- How can we detect problems with the data?
- What can we do about these problems?
- Examples of data quality problems:
 - Noise and outliers
 - Missing values
 - Duplicate data
 - Wrong data

Noise

- For objects, noise is an extraneous object
- For attributes, noise refers to modification of original values
 - Examples: distortion of a person's voice when talking on a poor phone and "snow" on television screen




Two Sine Waves



Two Sine Waves + Noise

Outliers

- **Outliers** are data objects with characteristics that are considerably different than most of the other data objects in the data set
 - **Case 1:** Outliers are noise that interferes with data analysis
 - **Case 2:** Outliers are the goal of our analysis
 - Credit card fraud
 - Intrusion detection
- Causes?



Missing Values

- Reasons for missing values
 - Information is not collected
(e.g., people decline to give their age and weight)
 - Attributes may not be applicable to all cases
(e.g., annual income is not applicable to children)
- Handling missing values
 - Eliminate data objects or variables
 - Estimate missing values
 - Example: time series of temperature
 - Example: census results
 - Ignore the missing value during analysis

Missing Values ...

- Missing completely at random (MCAR)
 - Missingness of a value is independent of attributes
 - Fill in values based on the attribute
 - Analysis may be unbiased overall
- Missing at Random (MAR)
 - Missingness is related to other variables
 - Fill in values based other values
 - Almost always produces a bias in the analysis
- Missing Not at Random (MNAR)
 - Missingness is related to unobserved measurements
 - Informative or non-ignorable missingness
- Not possible to know the situation from the data

Duplicate Data

- Data set may include data objects that are duplicates, or almost duplicates of one another
 - Major issue when merging data from heterogeneous sources
- Examples:
 - Same person with multiple email addresses
- Data cleaning
 - Process of dealing with duplicate data issues
- When should duplicate data not be removed?

Similarity and Dissimilarity Measures

- Similarity measure
 - Numerical measure of how alike two data objects are.
 - Is higher when objects are more alike.
 - Often falls in the range $[0,1]$
- Dissimilarity measure
 - Numerical measure of how different two data objects are
 - Lower when objects are more alike
 - Minimum dissimilarity is often 0
 - Upper limit varies
- Proximity refers to a similarity or dissimilarity

Similarity/Dissimilarity for Simple Attributes

The following table shows the similarity and dissimilarity between two objects, x and y , with respect to a single, simple attribute.

Attribute Type	Dissimilarity	Similarity
Nominal	$d = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$	$s = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{if } x \neq y \end{cases}$
Ordinal	$d = x - y / (n - 1)$ (values mapped to integers 0 to $n-1$, where n is the number of values)	$s = 1 - d$
Interval or Ratio	$d = x - y $	$s = -d, s = \frac{1}{1+d}, s = e^{-d},$ $s = 1 - \frac{d - \min_d}{\max_d - \min_d}$

Euclidean Distance


- Euclidean Distance

$$d(\mathbf{x}, \mathbf{y}) = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}$$

where n is the number of dimensions (attributes) and x_k and y_k are, respectively, the k^{th} attributes (components) or data objects \mathbf{x} and \mathbf{y} .

- Standardization is necessary, if scales differ.

Euclidean Distance



point	x	y
p1	0	2
p2	2	0
p3	3	1
p4	5	1

	p1	p2	p3	p4
p1	0	2.828	3.162	5.099
p2	2.828	0	1.414	3.162
p3	3.162	1.414	0	2
p4	5.099	3.162	2	0

Distance Matrix

Minkowski Distance

- Minkowski Distance is a generalization of Euclidean Distance

$$d(\mathbf{x}, \mathbf{y}) = \left(\sum_{k=1}^n |x_k - y_k|^r \right)^{1/r}$$

Where r is a parameter, n is the number of dimensions (attributes) and x_k and y_k are, respectively, the k^{th} attributes (components) or data objects \mathbf{x} and \mathbf{y} .

Minkowski Distance: Examples

- $r = 1$. City block (Manhattan, taxicab, L_1 norm) distance.
 - A common example of this is the Hamming distance, which is just the number of bits that are different between two binary vectors
- $r = 2$. Euclidean distance
- $r \rightarrow \infty$. “supremum” (L_{\max} norm, L_{∞} norm) distance.
 - This is the maximum difference between any component of the vectors
- Do not confuse r with n , i.e., all these distances are defined for all numbers of dimensions.

Minkowski Distance

point	x	y
p1	0	2
p2	2	0
p3	3	1
p4	5	1

L1	p1	p2	p3	p4
p1	0	4	4	6
p2	4	0	2	4
p3	4	2	0	2
p4	6	4	2	0


L2	p1	p2	p3	p4
p1	0	2.828	3.162	5.099
p2	2.828	0	1.414	3.162
p3	3.162	1.414	0	2
p4	5.099	3.162	2	0

L_{∞}	p1	p2	p3	p4
p1	0	2	3	5
p2	2	0	1	3
p3	3	1	0	2
p4	5	3	2	0

Distance Matrix

Mahalanobis Distance


$$\text{mahalanobis}(\mathbf{x}, \mathbf{y}) = (\mathbf{x} - \mathbf{y})^T \Sigma^{-1} (\mathbf{x} - \mathbf{y})$$



Σ is the covariance matrix

For red points, the Euclidean distance is 14.7, Mahalanobis distance is 6.

Mahalanobis Distance



**Covariance
Matrix:**

$$\Sigma = \begin{bmatrix} 0.3 & 0.2 \\ 0.2 & 0.3 \end{bmatrix}$$

A: (0.5, 0.5)

B: (0, 1)

C: (1.5, 1.5)

Mahal(A,B) = 5

Mahal(A,C) = 4

Common Properties of a Distance

- Distances, such as the Euclidean distance, have some well known properties.

1. $d(\mathbf{x}, \mathbf{y}) \geq 0$ for all \mathbf{x} and \mathbf{y} and $d(\mathbf{x}, \mathbf{y}) = 0$ only if $\mathbf{x} = \mathbf{y}$. (Positive definiteness)
2. $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$ for all \mathbf{x} and \mathbf{y} . (Symmetry)
3. $d(\mathbf{x}, \mathbf{z}) \leq d(\mathbf{x}, \mathbf{y}) + d(\mathbf{y}, \mathbf{z})$ for all points \mathbf{x} , \mathbf{y} , and \mathbf{z} . (Triangle Inequality)

where $d(\mathbf{x}, \mathbf{y})$ is the distance (dissimilarity) between points (data objects), \mathbf{x} and \mathbf{y} .

- A distance that satisfies these properties is a **metric**

Common Properties of a Similarity

- Similarities, also have some well known properties.

1. $s(\mathbf{x}, \mathbf{y}) = 1$ (or maximum similarity) only if $\mathbf{x} = \mathbf{y}$.

2. $s(\mathbf{x}, \mathbf{y}) = s(\mathbf{y}, \mathbf{x})$ for all \mathbf{x} and \mathbf{y} . (Symmetry)

where $s(\mathbf{x}, \mathbf{y})$ is the similarity between points (data objects), \mathbf{x} and \mathbf{y} .

Similarity Between Binary Vectors

- Common situation is that objects, p and q , have only binary attributes
- Compute similarities using the following quantities
 - f_{01} = the number of attributes where p was 0 and q was 1
 - f_{10} = the number of attributes where p was 1 and q was 0
 - f_{00} = the number of attributes where p was 0 and q was 0
 - f_{11} = the number of attributes where p was 1 and q was 1
- Simple Matching and Jaccard Coefficients
 - SMC = number of matches / number of attributes
 - = $(f_{11} + f_{00}) / (f_{01} + f_{10} + f_{11} + f_{00})$
 - J = number of 11 matches / number of non-zero attributes
 - = $(f_{11}) / (f_{01} + f_{10} + f_{11})$

SMC versus Jaccard: Example

$$\mathbf{x} = 1\ 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0\ 0$$

$$\mathbf{y} = 0\ 0\ 0\ 0\ 0\ 0\ 0\ 1\ 0\ 0\ 1$$

$$f_{01} = 2 \quad (\text{the number of attributes where } p \text{ was } 0 \text{ and } q \text{ was } 1)$$

$$f_{10} = 1 \quad (\text{the number of attributes where } p \text{ was } 1 \text{ and } q \text{ was } 0)$$

$$f_{00} = 7 \quad (\text{the number of attributes where } p \text{ was } 0 \text{ and } q \text{ was } 0)$$

$$f_{11} = 0 \quad (\text{the number of attributes where } p \text{ was } 1 \text{ and } q \text{ was } 1)$$

$$\begin{aligned} \text{SMC} &= (f_{11} + f_{00}) / (f_{01} + f_{10} + f_{11} + f_{00}) \\ &= (0+7) / (2+1+0+7) = 0.7 \end{aligned}$$

$$\text{J} = (f_{11}) / (f_{01} + f_{10} + f_{11}) = 0 / (2 + 1 + 0) = 0$$

Cosine Similarity

- If \mathbf{d}_1 and \mathbf{d}_2 are two document vectors, then

$$\cos(\mathbf{d}_1, \mathbf{d}_2) = \langle \mathbf{d}_1, \mathbf{d}_2 \rangle / \|\mathbf{d}_1\| \|\mathbf{d}_2\| ,$$

where $\langle \mathbf{d}_1, \mathbf{d}_2 \rangle$ indicates inner product or vector dot product of vectors, \mathbf{d}_1 and \mathbf{d}_2 , and $\|\mathbf{d}\|$ is the length of vector \mathbf{d} .

- Example:

$$\mathbf{d}_1 = 3 \ 2 \ 0 \ 5 \ 0 \ 0 \ 0 \ 2 \ 0 \ 0$$

$$\mathbf{d}_2 = 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 2$$

$$\langle \mathbf{d}_1, \mathbf{d}_2 \rangle = 3*1 + 2*0 + 0*0 + 5*0 + 0*0 + 0*0 + 0*0 + 2*1 + 0*0 + 0*2 = 5$$

$$\|\mathbf{d}_1\| = (3*3 + 2*2 + 0*0 + 5*5 + 0*0 + 0*0 + 0*0 + 2*2 + 0*0 + 0*0)^{0.5} = (42)^{0.5} = 6.481$$

$$\|\mathbf{d}_2\| = (1*1 + 0*0 + 0*0 + 0*0 + 0*0 + 0*0 + 0*0 + 1*1 + 0*0 + 2*2)^{0.5} = (6)^{0.5} = 2.449$$

$$\cos(\mathbf{d}_1, \mathbf{d}_2) = 0.3150$$

Correlation measures the linear relationship between objects

$$\text{corr}(\mathbf{x}, \mathbf{y}) = \frac{\text{covariance}(\mathbf{x}, \mathbf{y})}{\text{standard_deviation}(\mathbf{x}) * \text{standard_deviation}(\mathbf{y})} = \frac{s_{xy}}{s_x s_y}, \quad (2.11)$$

where we are using the following standard statistical notation and definitions

$$\text{covariance}(\mathbf{x}, \mathbf{y}) = s_{xy} = \frac{1}{n-1} \sum_{k=1}^n (x_k - \bar{x})(y_k - \bar{y}) \quad (2.12)$$


$$\text{standard_deviation}(\mathbf{x}) = s_x = \sqrt{\frac{1}{n-1} \sum_{k=1}^n (x_k - \bar{x})^2}$$

$$\text{standard_deviation}(\mathbf{y}) = s_y = \sqrt{\frac{1}{n-1} \sum_{k=1}^n (y_k - \bar{y})^2}$$

$$\bar{x} = \frac{1}{n} \sum_{k=1}^n x_k \text{ is the mean of } \mathbf{x}$$

$$\bar{y} = \frac{1}{n} \sum_{k=1}^n y_k \text{ is the mean of } \mathbf{y}$$

Visually Evaluating Correlation



**Scatter plots
showing the
similarity from
-1 to 1.**

Information Based Measures

- Information theory is a well-developed and fundamental discipline with broad applications
- Some similarity measures are based on information theory
 - Mutual information in various versions
 - Maximal Information Coefficient (MIC) and related measures
 - General and can handle non-linear relationships
 - Can be complicated and time intensive to compute

Information and Probability



- Information relates to possible outcomes of an event
 - transmission of a message, flip of a coin, or measurement of a piece of data
- The more certain an outcome, the less information that it contains and vice-versa
 - For example, if a coin has two heads, then an outcome of heads provides no information
 - More quantitatively, the information is related the probability of an outcome
 - The smaller the probability of an outcome, the more information it provides and vice-versa
 - Entropy is the commonly used measure

Entropy

- For
 - a variable (event), X ,
 - with n possible values (outcomes), x_1, x_2, \dots, x_n
 - each outcome having probability, p_1, p_2, \dots, p_n
 - the entropy of X , $H(X)$, is given by

$$H(X) = - \sum_{i=1}^n p_i \log_2 p_i$$

- Entropy is between 0 and $\log_2 n$ and is measured in bits
 - Thus, entropy is a measure of how many bits it takes to represent an observation of X on average

Entropy Examples

- For a coin with probability p of heads and probability $q = 1 - p$ of tails

$$H = -p \log_2 p - q \log_2 q$$

- For $p = 0.5$, $q = 0.5$ (fair coin) $H = 1$
 - For $p = 1$ or $q = 1$, $H = 0$
- What is the entropy of a fair four-sided die?

Entropy for Sample Data: Example

Hair Color	Count	p	$-p\log_2 p$
Black	75	0.75	0.3113
Brown	15	0.15	0.4105
Blond	5	0.05	0.2161
Red	0	0.00	0
Other	5	0.05	0.2161
Total	100	1.0	1.1540

Entropy for Sample Data

- Suppose we have
 - a number of observations (m) of some attribute, X , e.g., the hair color of students in the class,
 - where there are n different possible values
 - And the number of observation in the i^{th} category is m_i
 - Then, for this sample

$$H(X) = - \sum_{i=1}^n \frac{m_i}{m} \log_2 \frac{m_i}{m}$$

- For continuous data, the calculation is harder

Mutual Information

- Information one variable provides about another

Formally, $I(X, Y) = H(X) + H(Y) - H(X, Y)$, where

$H(X, Y)$ is the joint entropy of X and Y ,

$$H(X, Y) = - \sum_i \sum_j p_{ij} \log_2 p_{ij}$$

Where p_{ij} is the probability that the i^{th} value of X and the j^{th} value of Y occur together

- For discrete variables, this is easy to compute
- Maximum mutual information for discrete variables is $\log_2(\min(n_X, n_Y))$, where n_X (n_Y) is the number of values of X (Y)

Mutual Information Example

Student Status	Count	p	$-p\log_2 p$
Undergrad	45	0.45	0.5184
Grad	55	0.55	0.4744
Total	100	1.00	0.9928

Grade	Count	p	$-p\log_2 p$
A	35	0.35	0.5301
B	50	0.50	0.5000
C	15	0.15	0.4105
Total	100	1.00	1.4406

Student Status	Grade	Count	p	$-p\log_2 p$
Undergrad	A	5	0.05	0.2161
Undergrad	B	30	0.30	0.5211
Undergrad	C	10	0.10	0.3322
Grad	A	30	0.30	0.5211
Grad	B	20	0.20	0.4644
Grad	C	5	0.05	0.2161
Total		100	1.00	2.2710


Mutual information of Student Status and Grade = $0.9928 + 1.4406 - 2.2710 = 0.1624$

Density

- Measures the degree to which data objects are close to each other in a specified area
- The notion of density is closely related to that of proximity
- Concept of density is typically used for clustering and anomaly detection
- Examples:
 - Euclidean density
 - Euclidean density = number of points per unit volume
 - Probability density
 - Estimate what the distribution of the data looks like
 - Graph-based density
 - Connectivity

Euclidean Density: Grid-based Approach

- Simplest approach is to divide region into a number of rectangular cells of equal volume and define density as # of points the cell contains



Grid-based density.

0	0	0	0	0	0	0
0	0	0	0	0	0	0
4	17	18	6	0	0	0
14	14	13	13	0	18	27
11	18	10	21	0	24	31
3	20	14	4	0	0	0
0	0	0	0	0	0	0

Counts for each cell.

Euclidean Density: Center-Based

- Euclidean density is the number of points within a specified radius of the point




Illustration of center-based density.

Linear Regression

Linear Regression

The technique is used to **predict** the value of one variable (the dependent variable - y) **based on** the value of other variables (independent variables x_1, x_2, \dots, x_k)

$$\overline{y = \beta_0 + \beta_1 x + \varepsilon}$$

Modeling

- The first order linear model


y = dependent variable

x = independent variable

β_0 = y-intercept


β_1 = slope of the line

\mathcal{E} = error variable



Estimating the coefficients

- The estimates are determined by
 - drawing a sample from the population of interest,
 - calculating sample statistics.
 - producing a straight line that cuts into the data.



The question is:
Which straight line fits best?


The best line is the one that minimizes the sum of squared vertical differences between the points and the line.

Sum of squared differences $= (2 - 1)^2 + (4 - 2)^2 + (1.5 - 3)^2 + (3.2 - 4)^2 = 6.89$

Sum of squared differences $= (2 - 2.5)^2 + (4 - 2.5)^2 + (1.5 - 2.5)^2 + (3.2 - 2.5)^2 = 3.99$

Let us compare two lines

The second line is horizontal



The smaller the sum of squared differences the better the fit of the line to the data.

To calculate the estimates of the coefficients that minimize the differences between the data points and the line, use the formulas:

$$b_1 = \frac{\text{cov}(X, Y)}{s_x^2}$$
$$b_0 = \bar{y} - b_1 \bar{x}$$

The regression equation that estimates the equation of the first order linear model is:

$$\hat{y} = b_0 + b_1 x$$

Relationship between odometer reading and a used car's selling price.

- A car dealer wants to find the relationship between the odometer reading and the selling price of used cars.
- A random sample of 100 cars is selected, and the data recorded.
- Find the regression line.

Car	Odometer	Price
1	37388	5318
2	44758	5061
3	45833	5008
4	30862	5795
5	31705	5784
6	34010	5359
.	.	.
.	.	.
.	.	.

Independent variable x

Dependent variable y

Solution to calculate b_0 and b_1 we need to calculate several statistics first;

$$\bar{x} = 36,009.45; \quad s_x^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = 43,528,688$$

$$\bar{y} = 5,411.41; \quad \text{cov}(X, Y) = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{n-1} = -1,356,256$$

where $n = 100$.

$$b_1 = \frac{\text{cov}(X, Y)}{s_x^2} = \frac{-1,356,256}{43,528,688} = -.0312$$

$$b_0 = \bar{y} - b_1 \bar{x} = 5411.41 - (-.0312)(36,009.45) = 6,533$$

$$\hat{y} = b_0 + b_1 x = 6,533 - .0312x$$