

Experiment 5

The Laws of Radiation - Stefan-Boltzmann Law (Radiant Flux)

METHOD

If the temperature of the cavity is T , then we can ascribe a characteristic energy $\epsilon^* = kT$ to a typical photon. Such photon would have a de Broglie wavelength $l^* = h/p^* = hc/\epsilon^*$ and characteristic frequency $\nu^* = \epsilon^*/h$. Defining a dimensionless quantity $x = \epsilon/\epsilon^*$, the spectral energy density for a particular mode ($\epsilon = h\nu$) can be rewritten as

$$u(\nu) d\nu \rightarrow u(x) dx = \frac{8\pi\epsilon^*}{l^{*3}} \left(\frac{x^3}{e^x - 1} \right) dx = \frac{8\pi\epsilon^*}{l^{*3}} f_P(x) dx$$

where $f_P(x) = \frac{x^3}{e^x - 1}$ and

$$u = \int_0^\infty u(\nu) d\nu = \int_0^\infty \bar{\epsilon} \frac{G(\nu) d\nu}{V_0} = \frac{8\pi h}{c^3} \int_0^\infty \left(\frac{\nu^3}{e^{\frac{h\nu}{kT}} - 1} \right) d\nu = \frac{8\pi^2 k^4}{15c^3 h^3} T^4 = \frac{4}{c} \sigma T^4$$

with density of states given by

$$G(\nu) d\nu = \frac{8\pi\nu^2}{c^3} d\nu V_0$$

CODING AND PLOTS

1. (Wiens Displacement law, Energy of states)

The plot of $f_P(x)$ versus dimensionless variable x peaks at a certain $x = x_p$. Correspondingly, the spectral energy density will peak at x_p irrespective of the temperature of the cavity.

(a) Find the value of x_p .

(b) Since the dimensionless variable x can also be written as $x = \epsilon/\epsilon^* = hc/\lambda k_b T$, deduce the Wien's constant b

$$\lambda T = \frac{hc}{k x_p} = b$$

Prove that it is $b = 0.003 \text{ meter Kelvin}$ and hence the Weins displacement Law

2. (Stefan-Boltzmann Law)

With the characteristic parameters of the cavity, the total energy density u of the radiation in a cavity at temperature T is given by

$$u = \int_0^\infty u(\nu) d\nu = \frac{8\pi\epsilon^*}{l^{*3}} \int_0^\infty \left(\frac{x^3}{e^x - 1} \right) dx = \frac{8\pi\epsilon^*}{l^{*3}} \int_0^\infty f_P(x) dx = \frac{8\pi\epsilon^*}{l^{*3}} I_P$$

The integral I_P is independent of the parameters of the cavity.

With numerical integration show that $I_P = \pi^4/15$

The total energy density u of the radiation in a cavity at temperature T thus becomes

$$u = \frac{\pi^4}{15} \frac{8\pi\epsilon^*}{l^{*3}} = \frac{\pi^4}{15} C(T)$$

Calculate the factor $C(T)$ for temperatures $T = [100K, 10000K]$ in steps of $500 K$.

Find the energy density $u(T) = (\pi^4/15)C(T)$ for these temperatures. The total energy density $u(T)$ of the radiation in a cavity at temperature T and the radiant flux $F(T)$ are related by $F = \frac{c}{4}u$. Find the radiant flux $F(T) = (c/4)(\pi^4/15)C(T)$ for these temperatures.

Plot

(a) F versus T and also

(b) plot $\ln F$ versus $\ln T$. Find the value of slope and intercept from the graph in part

(c) interpret.

Hence, deduce the Stefan Boltzmann law

$$F = \sigma T^4 = \left(\frac{c}{4}\right) \frac{8\pi^2 k^4}{15c^3 h^3} T^4$$