

# The combinatorics of the leading root of Ramanujan's (and related) functions

Thomas Prellberg

School of Mathematical Sciences  
Queen Mary, University of London

Mathematics Colloquium, University of Kent  
January 28, 2014

# In Memoriam

Donald Preece

(† Jan 8, 2014)



Emeritus Professor of Mathematical Sciences, QMUL  
Emeritus Professor of Combinatorial Mathematics, Kent

# Topic Outline

- 1  $q$ -Airy (and related) Functions
- 2 More on Theta and Partial Theta Functions
- 3 Identities for the Leading Roots
- 4 Combinatorics
- 5 Outlook

# Outline

- 1  $q$ -Airy (and related) Functions
  - Ramanujan's Function
  - Painlevé Airy Function
  - Partial Theta Function
- 2 More on Theta and Partial Theta Functions
- 3 Identities for the Leading Roots
- 4 Combinatorics
- 5 Outlook

# Ramanujan's Function

$$A_q(x) = \sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} = \sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(1-q)(1-q^2)\dots(1-q^n)}$$

# Ramanujan's Function

$$A_q(x) = \sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} = \sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(1-q)(1-q^2)\dots(1-q^n)}$$

Rogers-Ramanujan Identities

$$A_q(-1) = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})}$$


and

$$A_q(-q) = \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q; q)_n} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})}$$

Related to partitions of integers into parts mod 5


# Area-weighted Dyck paths

Count Dyck paths with respect to steps and enclosed area



# Area-weighted Dyck paths

Count Dyck paths with respect to steps and enclosed area



Generating function

$$G(x, q) = \sum_{m,n} c_{m,n} x^n q^m = \frac{A_q(x)}{A_q(x/q)}$$

$x$  counts pairs of up/down steps,  $q$  counts enclosed area



# Ramanujan's Lost Notebook, page 57

$$\sum_{n=0}^{\infty} \frac{q^{n^2} (-x)^n}{(q; q)_n} = \prod_{n=1}^{\infty} \left( 1 - \frac{x q^{2n-1}}{1 - q^n y_1 - q^{2n} y_2 - q^{3n} y_3 - \dots} \right)$$

where

$$(t; q)_n = (1 - t)(1 - tq)(1 - tq^2) \dots (1 - tq^{n-1})$$

with  $y_1, y_2, y_3, y_4$  explicitly given

# Ramanujan's Lost Notebook, page 57

$$\sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} = \prod_{n=1}^{\infty} \left( 1 - \frac{xq^{2n-1}}{1 - q^n y_1 - q^{2n} y_2 - q^{3n} y_3 - \dots} \right)$$

where

$$(t; q)_n = (1 - t)(1 - tq)(1 - tq^2) \dots (1 - tq^{n-1})$$

with  $y_1, y_2, y_3, y_4$  explicitly given

More precisely,

$$y_1 = \frac{1}{(1 - q)\psi^2(q)}, \quad y_2 = 0$$

$$y_3 = \frac{q + q^3}{(1 - q)(1 - q^2)(1 - q^3)\psi^2(q)} - \frac{\sum_{n=0}^{\infty} \frac{(2n+1)q^{2n+1}}{1 - q^{2n+1}}}{(1 - q)^3\psi^6(q)}, \quad y_4 = y_1 y_3$$

and

$$\psi(q) = \sum_{n=0}^{\infty} q^{n(n+1)/2} = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}}$$

# The roots of $A_q(x)$

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} &= \prod_{n=1}^{\infty} \left( 1 - \frac{xq^{2n-1}}{1 - q^n y_1 - q^{2n} y_2 - q^{3n} y_3 - \dots} \right) \\ &= \prod_{n=0}^{\infty} \left( 1 - \frac{x}{x_n(q)} \right)\end{aligned}$$

# The roots of $A_q(x)$

$$\begin{aligned}
 \sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} &= \prod_{n=1}^{\infty} \left( 1 - \frac{xq^{2n-1}}{1 - q^n y_1 - q^{2n} y_2 - q^{3n} y_3 - \dots} \right) \\
 &= \prod_{n=0}^{\infty} \left( 1 - \frac{x}{x_n(q)} \right)
 \end{aligned}$$

- Roots are positive, real, and simple (Al-Salam and Ismail, 1983)
- Ramanujan's expansion is an asymptotic series (Andrews, 2005)
- Relation to Stieltjes-Wigert polynomials (Andrews, 2005)
- Integral equation for roots (Ismail and Zhang, 2007)
- Combinatorial interpretation of  $y_k$  (Huber, 2008, and Huber and Yee, 2010)

# The aim of this talk

$$\sum_{n=0}^{\infty} \frac{q^{n^2} (-x)^n}{(q; q)_n} = \prod_{n=0}^{\infty} \left( 1 - \frac{x}{x_n(q)} \right)$$

$$qx_0(-q) = 1 + q + q^2 + 2q^3 + 4q^4 + 8q^5 + 16q^6 + 33q^7 + 70q^8 + 151q^9 + \dots$$

# The aim of this talk

$$\sum_{n=0}^{\infty} \frac{q^{n^2}(-x)^n}{(q; q)_n} = \prod_{n=0}^{\infty} \left(1 - \frac{x}{x_n(q)}\right)$$

$$qx_0(-q) = 1 + q + q^2 + 2q^3 + 4q^4 + 8q^5 + 16q^6 + 33q^7 + 70q^8 + 151q^9 + \dots$$

## Goal

A combinatorial interpretation of the coefficients of the leading root

# Painlevé Airy Function

$$\text{Ai}_q(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(q^2; q^2)_n} = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(1 - q^2)(1 - q^4) \dots (1 - q^{2n})}$$

# Painlevé Airy Function

$$\text{Ai}_q(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(q^2; q^2)_n} = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(1 - q^2)(1 - q^4) \dots (1 - q^{2n})}$$

$$x_0(q) = 1 + q + q^2 + 2q^3 + 3q^4 + 6q^5 + 12q^6 + 25q^7 + 54q^8 + 120q^9 + \dots$$

The coefficients of the leading root also seem to be positive integers



# Painlevé Airy Function

$$\text{Ai}_q(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(q^2; q^2)_n} = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(1 - q^2)(1 - q^4) \dots (1 - q^{2n})}$$

# Painlevé Airy Function

$$\text{Ai}_q(x) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(q^2; q^2)_n} = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(1-q^2)(1-q^4)\dots(1-q^{2n})}$$

Connection Formula (Morita, 2011)

$$A_{q^2} \left( -\frac{q^3}{x^2} \right) = \frac{1}{(q; q)_{\infty} (-1; q)_{\infty}} \left\{ \Theta \left( -\frac{x}{q}, q \right) \text{Ai}_q(-x) + \Theta \left( \frac{x}{q}, q \right) \text{Ai}_q(x) \right\}$$

with Theta Function

$$\Theta(x, q) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-x)^n$$

# Partial Theta Function

The Theta Function has roots  $x_k(q) = q^k \quad k \in \mathbb{Z}$

$$\Theta(x, q) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-x)^n = (q; q)_{\infty} (x; q)_{\infty} (q/x; q)_{\infty}$$

# Partial Theta Function

The Theta Function has roots  $x_k(q) = q^k \quad k \in \mathbb{Z}$

$$\Theta(x, q) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-x)^n = (q; q)_{\infty} (x; q)_{\infty} (q/x; q)_{\infty}$$

The Partial Theta Function

$$\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (-x)^n$$

does not admit a “nice” product formula, but

$$x_0(q) = 1 + q + 2q^2 + 4q^3 + 9q^4 + 21q^5 + 52q^6 + 133q^7 + 351q^8 + 948q^9 + \dots$$

The coefficients of the leading root are positive integers (Sokal, 2012)

# Outline

- 1  $q$ -Airy (and related) Functions
- 2 More on Theta and Partial Theta Functions
  - Jacobi Theta Function
  - $q$ -Theta Function
  - Partial Theta Function
  - Rogers-Ramanujan Function
- 3 Identities for the Leading Roots
- 4 Combinatorics
- 5 Outlook

# Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} \exp(\pi i n^2 \tau + 2\pi i n z)$$

# Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} \exp(\pi i n^2 \tau + 2\pi i n z)$$

- Quasi-periodic function satisfying

$$\vartheta(z + a + b\tau; \tau) = \exp(-\pi i b^2 \tau - 2\pi i b z) \vartheta(z; \tau)$$

# Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} \exp(\pi i n^2 \tau + 2\pi i n z)$$

- Quasi-periodic function satisfying

$$\vartheta(z + a + b\tau; \tau) = \exp(-\pi i b^2 \tau - 2\pi i b z) \vartheta(z; \tau)$$

- Relation to modular group

$$\vartheta(z/\tau; -1/\tau) = (-i\tau)^{1/2} \exp(\pi i z^2/\tau) \vartheta(z; \tau)$$



# Roots of the Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} q^{n^2} x^{2n}$$

where  $q = \exp(\pi i \tau)$  and  $x = \exp(\pi i z)$

# Roots of the Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} q^{n^2} x^{2n}$$

where  $q = \exp(\pi i \tau)$  and  $x = \exp(\pi i z)$

- Jacobi triple product

$$\sum_{n=-\infty}^{\infty} q^{n^2} x^{2n} = \prod_{m=1}^{\infty} (1 - q^{2m})(1 + q^{2m-1} x^2)(1 + q^{2m-1}/x^2)$$

# Roots of the Jacobi Theta Function

- Jacobi Theta function

$$\vartheta(z; \tau) = \sum_{n=-\infty}^{\infty} q^{n^2} x^{2n}$$

where  $q = \exp(\pi i \tau)$  and  $x = \exp(\pi i z)$

- Jacobi triple product

$$\sum_{n=-\infty}^{\infty} q^{n^2} x^{2n} = \prod_{m=1}^{\infty} (1 - q^{2m})(1 + q^{2m-1} x^2)(1 + q^{2m-1}/x^2)$$

- Roots

$$x_k(q) = -q^{k+1/2} \quad k \in \mathbb{Z}$$

# $q$ -Theta Function

- Combinatorialists prefer

$$\Theta(x, q) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-x)^n = (q; q)_{\infty} (x; q)_{\infty} (q/x; q)_{\infty}$$

with  $q$ -product notation

$$(t; q)_n = \prod_{m=0}^{n-1} (1 - tq^m)$$

# *q*-Theta Function

- Combinatorialists prefer

$$\Theta(x, q) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-x)^n = (q; q)_{\infty} (x; q)_{\infty} (q/x; q)_{\infty}$$

with *q*-product notation

$$(t; q)_n = \prod_{m=0}^{n-1} (1 - tq^m)$$

- Roots

$$x_k(q) = q^k \quad k \in \mathbb{Z}$$

# Partial Theta Function

- Partial Theta Function

$$\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n$$

# Partial Theta Function

- Partial Theta Function

$$\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n$$

- No “nice” product formula

# Partial Theta Function

- Partial Theta Function

$$\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n$$

- No “nice” product formula
- Roots

$$x_k(q) = ?$$



# Partial Theta Function

- Partial Theta Function

$$\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n$$

- No “nice” product formula
- Roots

$$x_k(q) = ?$$

- Special case  $R(x, q, 0)$  of Rogers-Ramanujan Function

$$R(x, y, q) = \sum_{n=0}^{\infty} \frac{y^{\binom{n}{2}} x^n}{(q; q)_n}$$

# Rogers-Ramanujan Function

- Rogers-Ramanujan Function

$$R(x, y, q) = \sum_{n=0}^{\infty} \frac{y^{\binom{n}{2}} x^n}{(q; q)_n}$$

# Rogers-Ramanujan Function

- Rogers-Ramanujan Function

$$R(x, y, q) = \sum_{n=0}^{\infty} \frac{y^{\binom{n}{2}} x^n}{(q; q)_n}$$

- Euler identities

$$R(x, 1, q) = \sum_{n=0}^{\infty} \frac{x^n}{(q; q)_n} = \frac{1}{(x; q)_{\infty}}$$

and

$$R(x, q, q) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} x^n}{(q; q)_n} = (-x; q)_{\infty}$$

# Rogers-Ramanujan Function

- Rogers-Ramanujan Function

$$R(x, y, q) = \sum_{n=0}^{\infty} \frac{y^{\binom{n}{2}} x^n}{(q; q)_n}$$

- Euler identities

$$R(x, 1, q) = \sum_{n=0}^{\infty} \frac{x^n}{(q; q)_n} = \frac{1}{(x; q)_{\infty}}$$

and

$$R(x, q, q) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} x^n}{(q; q)_n} = (-x; q)_{\infty}$$

- $R(x, 1, q)$  has no roots, whereas  $R(x, q, q)$  has roots

$$x_k(1, q) = -q^{-k} \quad k \in \mathbb{N}_0$$

# Outline

- 1  $q$ -Airy (and related) Functions
- 2 More on Theta and Partial Theta Functions
- 3 **Identities for the Leading Roots**
  - The Leading Roots
  - Key Identities
  - Two Identities for  $\Theta_0(x, q)$
  - Two identities for  $\xi_0(q)$
  - Positivity
- 4 Combinatorics
- 5 Outlook

# The Leading Roots

Partial Theta Function  $\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (-x)^n$

$$\Theta_0(x, q) = 0$$

$$x = 1 + q + 2q^2 + 4q^3 + 9q^4 + 21q^5 + 52q^6 + 133q^7 + 351q^8 + 948q^9 + \dots$$

# The Leading Roots

Partial Theta Function  $\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (-x)^n$

$$\Theta_0(x, q) = 0$$

$$x = 1 + q + 2q^2 + 4q^3 + 9q^4 + 21q^5 + 52q^6 + 133q^7 + 351q^8 + 948q^9 + \dots$$

Ramanujan Function  $A_q(x) = \sum_{n=0}^{\infty} q^{n^2} (-x)^n / (q; q)_n$

$$A_{-q}(x/q) = 0$$

$$x = 1 + q + q^2 + 2q^3 + 4q^4 + 8q^5 + 16q^6 + 33q^7 + 70q^8 + 151q^9 + \dots$$

# The Leading Roots

Partial Theta Function  $\Theta_0(x, q) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (-x)^n$

$\Theta_0(x, q) = 0$

$x = 1 + q + 2q^2 + 4q^3 + 9q^4 + 21q^5 + 52q^6 + 133q^7 + 351q^8 + 948q^9 + \dots$

Ramanujan Function  $A_q(x) = \sum_{n=0}^{\infty} q^{n^2} (-x)^n / (q; q)_n$

$A_{-q}(x/q) = 0$

$x = 1 + q + q^2 + 2q^3 + 4q^4 + 8q^5 + 16q^6 + 33q^7 + 70q^8 + 151q^9 + \dots$

Painlevé Airy Function  $\text{Ai}_q(x) = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (-x)^n / (q^2; q^2)_n$

$\text{Ai}_q(x) = 0$

$x = 1 + q + q^2 + 2q^3 + 3q^4 + 6q^5 + 12q^6 + 25q^7 + 54q^8 + 120q^9 + \dots$



# Key Identities

$\Theta_0(x, q)$  satisfies (Andrews and Warnaar, 2007)

$$\Theta_0(x, q) = (x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (x; q)_n}$$

# Key Identities

$\Theta_0(x, q)$  satisfies (Andrews and Warnaar, 2007)

$$\Theta_0(x, q) = (x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (x; q)_n}$$

$A_q(x)$  satisfies (Gessel and Stanton, 1983)

$$A_{-q}(x/q) = (xq; q^2)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2-n} x^n}{(q^2; q^2)_n (xq; q^2)_n}$$

# Key Identities

$\Theta_0(x, q)$  satisfies (Andrews and Warnaar, 2007)

$$\Theta_0(x, q) = (x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (x; q)_n}$$

$A_q(x)$  satisfies (Gessel and Stanton, 1983)

$$A_{-q}(x/q) = (xq; q^2)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2-n} x^n}{(q^2; q^2)_n (xq; q^2)_n}$$

$\text{Ai}_q(x)$  satisfies (Gessel and Stanton, 1983)

$$\text{Ai}_q(x) = (x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2 + \binom{n}{2}} x^n}{(q^2; q^2)_n (x; q)_n}$$

# Identities for the Roots

Partial Theta Function

(Sokal, 2012)

$$\Theta_0(x, q) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (qx; q)_{n-1}}$$

# Identities for the Roots

## Partial Theta Function

(Sokal, 2012)

$$\Theta_0(x, q) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (qx; q)_{n-1}}$$

## Ramanujan Function

$$A_{-q}(x/q) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$

# Identities for the Roots

## Partial Theta Function

(Sokal, 2012)

$$\Theta_0(x, q) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (qx; q)_{n-1}}$$

## Ramanujan Function

$$A_{-q}(x/q) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$

## Painlevé Airy Function

$$\text{Ai}_q(x) = 0 \text{ if } x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2 + \binom{n}{2}} x^n}{(q^2; q^2)_n (qx; q)_{n-1}}$$

## Two identities for $\Theta_0(x, q)$

$\Theta_0(x, q)$  satisfies

$$\begin{aligned}\Theta_0(x, q) &= (q; q)_\infty (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^n}{(q; q)_n (-x; q)_n} \\ &= (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} (-x)^n}{(q; q)_n (-x; q)_n}\end{aligned}$$

## Two identities for $\Theta_0(x, q)$

$\Theta_0(x, q)$  satisfies

$$\begin{aligned}
 \Theta_0(x, q) &= (q; q)_\infty (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^n}{(q; q)_n (-x; q)_n} \\
 &= (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} (-x)^n}{(q; q)_n (-x; q)_n}
 \end{aligned}$$

- The first identity follows from Euler's identities



## Two identities for $\Theta_0(x, q)$

$\Theta_0(x, q)$  satisfies

$$\begin{aligned}\Theta_0(x, q) &= (q; q)_\infty (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^n}{(q; q)_n (-x; q)_n} \\ &= (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2} (-x)^n}{(q; q)_n (-x; q)_n}\end{aligned}$$

- The first identity follows from Euler's identities
- The first and second identity follow from Heine's transformations for  $q$ -deformed hypergeometric functions

# Proof of the first identity for $\Theta_0(x, q)$

$$\begin{aligned}
 \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n &= \sum_{n=0}^{\infty} q^{\binom{n}{2}} x^n \frac{(q; q)_{\infty}}{(q; q)_n (q^{n+1}; q)_{\infty}} \\
 &= (q; q)_{\infty} \sum_{n=0}^{\infty} q^{\binom{n}{2}} \frac{x^n}{(q; q)_n} \sum_{m=0}^{\infty} \frac{(q^{n+1})^m}{(q; q)_m} \\
 &= (q; q)_{\infty} \sum_{m=0}^{\infty} \frac{q^m}{(q; q)_m} \sum_{n=0}^{\infty} q^{\binom{n}{2}} \frac{(xq^m)^n}{(q; q)_n} \\
 &= (q; q)_{\infty} \sum_{m=0}^{\infty} \frac{q^m}{(q; q)_m} (-xq^m; q)_{\infty} \\
 &= (q; q)_{\infty} (-x; q)_{\infty} \sum_{m=0}^{\infty} \frac{q^m}{(q; q)_m (-x; q)_m}
 \end{aligned}$$

## Two functional equations for $\xi_0(q)$

Lemma [Sokal]

$\xi_0(q)$  satisfies

$$\xi_0(q) = 1 + \sum_{n=1}^{\infty} \frac{q^n}{(q; q)_n (\xi_0(q)q; q)_{n-1}}$$

and

$$\xi_0(q) = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} \xi_0(q)^n}{(q; q)_n (\xi_0(q)q; q)_{n-1}}$$

## Two functional equations for $\xi_0(q)$

Lemma [Sokal]

$\xi_0(q)$  satisfies

$$\xi_0(q) = 1 + \sum_{n=1}^{\infty} \frac{q^n}{(q; q)_n (\xi_0(q)q; q)_{n-1}}$$

and

$$\xi_0(q) = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} \xi_0(q)^n}{(q; q)_n (\xi_0(q)q; q)_{n-1}}$$

- This follows directly from the preceding identities for  $\Theta_0(x, q)$

# Proof of the first equation for $\xi_0(q)$

From

$$\Theta_0(x, q) = (q; q)_\infty (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^n}{(q; q)_n (-x; q)_n}$$

it follows that

$$\Theta_0(x, q) = (q; q)_\infty (-xq; q)_\infty \left[ 1 + x + \sum_{n=1}^{\infty} \frac{q^n}{(q; q)_n (-xq; q)_{n-1}} \right]$$

# Proof of the first equation for $\xi_0(q)$

From

$$\Theta_0(x, q) = (q; q)_\infty (-x; q)_\infty \sum_{n=0}^{\infty} \frac{q^n}{(q; q)_n (-x; q)_n}$$

it follows that

$$\Theta_0(x, q) = (q; q)_\infty (-xq; q)_\infty \left[ 1 + x + \sum_{n=1}^{\infty} \frac{q^n}{(q; q)_n (-xq; q)_{n-1}} \right]$$

Hence  $\Theta_0(-\xi_0(q), q) = 0$  implies that

$$0 = 1 - \xi_0(q) + \sum_{n=1}^{\infty} \frac{q^n}{(q; q)_n (\xi_0(q)q; q)_{n-1}}$$

# Positivity

The coefficients of the leading root of the partial theta function are positive integers (Sokal, 2012)

# Positivity

The coefficients of the leading root of the partial theta function are positive integers (Sokal, 2012)

- Letting  $x^{(0)} = 1$  and iterating

$$x^{(N+1)} = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(q; q)_n (qx^{(N)}; q)_{n-1}}$$

one can show coefficient-wise monotonicity of  $x^{(N)}$ , and hence positivity for the leading root of the Partial Theta Function



# Positivity

The coefficients of the leading root of the partial theta function are positive integers (Sokal, 2012)

- Letting  $x^{(0)} = 1$  and iterating

$$x^{(N+1)} = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(q; q)_n (qx^{(N)}; q)_{n-1}}$$

one can show coefficient-wise monotonicity of  $x^{(N)}$ , and hence positivity for the leading root of the Partial Theta Function

- The proof is easily adapted to Ramanujan's Function and the Painlevé Airy Function, hence proving positivity (and integrality) of the leading root for all three functions

# Positivity

The coefficients of the leading root of the partial theta function are positive integers (Sokal, 2012)

- Letting  $x^{(0)} = 1$  and iterating

$$x^{(N+1)} = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(q; q)_n (qx^{(N)}; q)_{n-1}}$$

one can show coefficient-wise monotonicity of  $x^{(N)}$ , and hence positivity for the leading root of the Partial Theta Function


- The proof is easily adapted to Ramanujan's Function and the Painlevé Airy Function, hence proving positivity (and integrality) of the leading root for all three functions

Is there an underlying combinatorial structure?

# Outline

- 1  $q$ -Airy (and related) Functions
- 2 More on Theta and Partial Theta Functions
- 3 Identities for the Leading Roots
- 4 **Combinatorics**
  - Ferrers Diagrams
  - Trees Decorated with Ferrers Diagrams
  - Changing the Area Weights
- 5 Outlook

# Ferrers Diagrams



a Ferrers diagram with Durfee square of size  $n$

# Why Ferrers Diagrams?

The generating function  $G(x, y, q)$  of Ferrers diagrams with  $n$ -th largest row having length  $n$  for some positive integer  $n$ , enumerated with respect to width ( $x$ ), height ( $y$ ), and total area ( $q$ ), is given by

$$G(x, y, q) = \sum_{n=1}^{\infty} \frac{(xy)^n q^{n^2}}{(qy; q)_n (qx; q)_{n-1}}$$

# Why Ferrers Diagrams?

The generating function  $G(x, y, q)$  of Ferrers diagrams with  $n$ -th largest row having length  $n$  for some positive integer  $n$ , enumerated with respect to width ( $x$ ), height ( $y$ ), and total area ( $q$ ), is given by

$$G(x, y, q) = \sum_{n=1}^{\infty} \frac{(xy)^n q^{n^2}}{(qy; q)_n (qx; q)_{n-1}}$$

Compare with


$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (qx; q)_{n-1}}$$

to get

$$x = 1 + G(x, 1, q)$$

# Enumerating Ferrers Diagrams

$$G(x, y, q) = \sum_{n=1}^{\infty} (xy)^n q^{n^2} \frac{1}{(yq; q)_n} \frac{1}{(xq; q)_{n-1}}$$



The sum is over all sizes  $n$  of Durfee squares

# Trees Decorated with Ferrers Diagrams

The functional equation

$$x = 1 + G(x, 1, q)$$

admits a combinatorial interpretation using the “theory of species”


## Theorem (TP, 2012)

Let  $F_q$  be the species of Ferrers diagrams with  $n$ -th largest row having length  $n$  for some integer  $n$ , weighted by area ( $q$ ), with size given by the width of the Ferrers diagram, augmented by the ‘empty polyomino’.

Then  $x$  enumerates  $F_q$ -enriched rooted trees ([trees decorated such that the out-degree of the vertex matches the width of the Ferrers diagram](#)) with respect to the total area of the Ferrers diagrams at the vertices of the tree.



# Trees Decorated with Ferrers Diagrams




Tree with area 15, contributing  $q^{15}$  to the Partial Theta Function root

# The Same Trees, But Different Area Weights

Partial Theta Function (TP, 2012)


$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q; q)_n (qx; q)_{n-1}}$$



# The Same Trees, But Different Area Weights

Ramanujan Function (TP, 2013)

$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$




Count dark area twice

# The Same Trees, But Different Area Weights


Painlevé Airy Function (TP, 2013)

$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2 + \binom{n}{2}} x^n}{(q^2; q^2)_n (qx; q)_{n-1}}$$



Count dark area twice

# Trees Decorated with Ferrers Diagrams



- contributing  $q^{15}$  to the Partial Theta Function root
- contributing  $q^{20}$  to the Ramanujan Function root
- contributing  $q^{19}$  to the Painlevé Airy Function root

# Outline

- 1  $q$ -Airy (and related) Functions
- 2 More on Theta and Partial Theta Functions
- 3 Identities for the Leading Roots
- 4 Combinatorics
- 5 Outlook
  - Why  $\Theta$ ,  $A_q$ ,  $Ai_q$ ?
  - Higher Roots

# Why are $\Theta$ , $A_q$ , $Ai_q$ special?

Can we understand and generalise?

# Why are $\Theta$ , $A_q$ , $Ai_q$ special?

Can we understand and generalise?

$$\sum_{n=0}^N \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n (q; q)_{N-n}} = 1$$

follows from the  $q$ -binomial theorem



# Why are $\Theta$ , $A_q$ , $Ai_q$ special?

Can we understand and generalise?

$$\sum_{n=0}^N \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n (q; q)_{N-n}} = 1$$

follows from the  $q$ -binomial theorem

$$\sum_{n=0}^N \frac{q^n}{(q^2; q^2)_n (q^2; q^2)_{N-n}} = \frac{(-q, q)_N}{(q^2; q^2)_N},$$

is given by Andrews (1976) and, more explicitly, by Cigler (1982)

# Why are $\Theta$ , $A_q$ , $Ai_q$ special?

Can we understand and generalise?

$$\sum_{n=0}^N \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n (q; q)_{N-n}} = 1$$

follows from the  $q$ -binomial theorem

$$\sum_{n=0}^N \frac{q^n}{(q^2; q^2)_n (q^2; q^2)_{N-n}} = \frac{(-q, q)_N}{(q^2; q^2)_N},$$

is given by Andrews (1976) and, more explicitly, by Cigler (1982)

$$\sum_{n=0}^N \frac{(-1)^n q^{n^2}}{(q^2; q^2)_n (q; q)_{N-n}} = \frac{1}{(q^2; q^2)_N},$$

follows from an identity by Cauchy

# Why are $\Theta$ , $A_q$ , $Ai_q$ special?

Can we understand and generalise?

$$\sum_{n=0}^N \frac{(-1)^n q^{\binom{n+1}{2}}}{(q; q)_n (q; q)_{N-n}} = 1$$

follows from the  $q$ -binomial theorem

$$\sum_{n=0}^N \frac{q^n}{(q^2; q^2)_n (q^2; q^2)_{N-n}} = \frac{(-q, q)_N}{(q^2; q^2)_N},$$

is given by Andrews (1976) and, more explicitly, by Cigler (1982)

$$\sum_{n=0}^N \frac{(-1)^n q^{n^2}}{(q^2; q^2)_n (q; q)_{N-n}} = \frac{1}{(q^2; q^2)_N},$$

follows from an identity by Cauchy

These look similar, but their proofs have nothing in common!

# Higher Roots for $A_q(x)$

Let  $x = y/q^{2m}$  in

$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$

## Higher Roots for $A_q(x)$

Let  $x = y/q^{2m}$  in

$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$

Then

$$A_{-q}(y/q^{2m+1}) = 0 \text{ if}$$

$$y = 1 + \frac{(-1)^m q^m}{\sum_{n=0}^m \frac{(-1)^n y^n q^n}{(q^2; q^2)_n} \prod_{k=1}^{m-n} (y - q^{2k})} \sum_{n=1}^{\infty} \frac{q^{n^2} y^{m+n}}{(q^2; q^2)_{m+n} (q^2 y; q^2)_{n-1}}$$

## Higher Roots for $A_q(x)$

Let  $x = y/q^{2m}$  in

$$x = 1 + \sum_{n=1}^{\infty} \frac{q^{n^2} x^n}{(q^2; q^2)_n (q^2 x; q^2)_{n-1}}$$

Then

$$A_{-q}(y/q^{2m+1}) = 0 \text{ if}$$

$$y = 1 + \frac{(-1)^m q^m}{\sum_{n=0}^m \frac{(-1)^n y^n q^n}{(q^2; q^2)_n} \prod_{k=1}^{m-n} (y - q^{2k})} \sum_{n=1}^{\infty} \frac{q^{n^2} y^{m+n}}{(q^2; q^2)_{m+n} (q^2 y; q^2)_{n-1}}$$

Numerically, the  $m$ -th root seems to satisfy

$$\frac{1}{q^{2m+1}} (1 + (-1)^m q^{m+1} (\text{positive terms}))$$

# Summary and Open Question

- Uncovered intriguing combinatorial content of roots of some special functions.

# Summary and Open Question

- Uncovered intriguing combinatorial content of roots of some special functions.
- Can one generalise this in any meaningful way?



# Summary and Open Question

- Uncovered intriguing combinatorial content of roots of some special functions.
- Can one generalise this in any meaningful way?

## The End