

# MTH5105 Differential and Integral Analysis

## 2010-2011

### Solutions 6

## 1 Exercise for Feedback

- 1) (a) Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be differentiable with bounded derivative. Show that  $f$  is uniformly continuous.  
[Hint: Use that if  $|f'(x)| \leq M$  for all  $x \in \mathbb{R}$  then  $|f(x) - f(y)| \leq M|x - y|$  for all  $x, y \in \mathbb{R}$  (similar to Exercise sheet 2).]
- (b) Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto x$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto \sin(x)$ . Prove or disprove:
- (i)  $f$  is uniformly continuous.
  - (ii)  $g$  is uniformly continuous.
  - (iii)  $fg$  is uniformly continuous.
  - (iv)  $x \mapsto \begin{cases} g(x)/f(x) & x \neq 0 \\ 1 & x = 0 \end{cases}$  is uniformly continuous.

Solution:

- (a) To say  $f'$  is bounded means that there exists an  $M > 0$  such that  $|f'(x)| \leq M$  for all  $x \in \mathbb{R}$ . Hence, as shown in exercise sheet 2,

$$\forall x, y \in \mathbb{R} : |f(x) - f(y)| \leq M|x - y|.$$

Now given  $\epsilon > 0$  choose  $\delta = \epsilon/M$ . Then

$$\forall x, y \in \mathbb{R} : |x - y| < \delta \Rightarrow |f(x) - f(y)| \leq M|x - y| < M\delta = \epsilon.$$

- (b) (i) TRUE:  $f$  is uniformly continuous.  
As  $f'(x) = 1$ ,  $|f'(x)| \leq M$  with  $M = 1$ , so  $f$  has bounded derivative, and the assertion follows from (a).
- (ii) TRUE:  $g$  is uniformly continuous.  
As  $g'(x) = \cos(x)$ ,  $|g'(x)| \leq M$  with  $M = 1$ , so  $g$  has bounded derivative, and the assertion follows from (a).
- (iii) FALSE:  $fg$  is not uniformly continuous.  
As  $(fg)'(x) = x \cos(x) + \sin(x)$ ,  $g'$  is not bounded. This alone is no proof, but it indicates that the reason for non-uniformity is that at  $x = 2n\pi$  we find  $(fg)'(x) = 2n\pi$  which is arbitrarily large.  
To turn this into a proof, we repeat the strategy from the example in the lecture: Given  $\delta > 0$  we need to pick  $x_n, y_n \in \mathbb{R}$  with  $|x_n - y_n| < \delta$  but  $|x_n \sin(x_n) - y_n \sin(y_n)| \geq 1$ . Taking  $x_n = 2n\pi$  and  $y_n = 2n\pi + \delta'$ , we estimate

$$|x_n \sin(x_n) - y_n \sin(y_n)| = |(2n\pi + \delta'/2) \sin(\delta'/2)| \geq 4n\delta'.$$

In the last step we need  $\delta' \leq \pi/2$ . Therefore, if we choose  $\delta' = \min(1/4n, \pi/2)$  then  $|x_n - y_n| < \delta$  and  $|x_n \sin(x_n) - y_n \sin(y_n)| \geq 1$ .

(iv) TRUE:  $x \mapsto h(x) = \begin{cases} f(x)/g(x) & x \neq 0 \\ 1 & x = 0 \end{cases}$  is uniformly continuous.

For  $x \neq 0$  the quotient rule gives  $h'(x) = (g/f)'(x) = (x \cos(x) - \sin(x))/x^2$ , and for  $x = 0$  we get

$$h'(0) = \lim_{x \rightarrow 0} \frac{\sin(x)/x - 1}{x} = \lim_{x \rightarrow 0} \frac{\sin(x) - x}{x^2} = \lim_{x \rightarrow 0} \frac{\cos(x) - 1}{2x} = \lim_{x \rightarrow 0} \frac{-\sin(x)}{2} = 0.$$

As  $\lim_{x \rightarrow 0} h'(x) = \lim_{x \rightarrow 0} \frac{x \cos(x) - \sin(x)}{x^2} = \lim_{x \rightarrow 0} \frac{\cos(x) + x \sin(x) - \cos(x)}{2x} = 0 = h'(0)$ ,  $h'$  is continuous and hence bounded on  $[-L, L]$  for any  $L > 0$ . Additionally, if  $|x| > L$  we estimate  $|h'(x)| \leq |\cos(x)/x| + |\sin(x)/x^2| < 1/L + 1/L^2$ . Therefore  $h'$  is bounded on  $\mathbb{R}$ , and the assertion follows from (a).

## 2 Extra Exercises

2) Let  $f : (0, 1) \rightarrow \mathbb{R}$  be continuous. Show that

- a)  $f$  is uniformly continuous if  $\lim_{x \rightarrow 0} f(x)$  and  $\lim_{x \rightarrow 1} f(x)$  exist.
- \*b) If  $f$  is uniformly continuous then  $\lim_{x \rightarrow 0} f(x)$  and  $\lim_{x \rightarrow 1} f(x)$  exist.

Solution:

- a) If  $A = \lim_{x \rightarrow 0} f(x)$  and  $B = \lim_{x \rightarrow 1} f(x)$  exist, then the function  $g : [0, 1] \rightarrow \mathbb{R}$  defined by

$$g(x) = \begin{cases} A & x = 0 \\ f(x) & 0 < x < 1 \\ B & x = 1 \end{cases}$$

is continuous on  $[0, 1]$  and therefore uniformly continuous on  $[0, 1]$ . The function  $f$  is a restriction of  $g$  to the smaller interval  $(0, 1)$  and therefore also uniformly continuous.

- \*b) This part is considerably harder. We start by showing that  $f$  is bounded. Let  $\varepsilon > 0$ . Then there exists a  $\delta > 0$  such that  $|f(x) - f(y)| < \varepsilon$  for any two points  $x, y \in (0, 1)$  that are less than distance  $\delta$  apart. Now two arbitrary points  $u, v \in (0, 1)$  are less than distance one apart and can therefore be connected by a chain of  $n = \lfloor 1/\delta \rfloor$  points such that two consecutive points are less than distance  $\delta$  apart. Therefore,  $|f(u) - f(v)| < (n + 1)\varepsilon$  is finite and  $f$  must be bounded.

We now show that  $\lim_{x \rightarrow 0} f(x)$  exists (the case of  $x \rightarrow 1$  is completely analogous). As we have established that  $f$  is bounded, we know that for  $0 < \delta < 1$

$$a(\delta) = \inf\{f(x) : 0 < x < \delta\} \quad \text{and} \quad b(\delta) = \sup\{f(x) : 0 < x < \delta\}$$

are well-defined, bounded functions of  $\delta$ . Moreover,  $a(\delta)$  increases as  $\delta \rightarrow 0$  and  $b(\delta)$  decreases as  $\delta \rightarrow 0$ . As  $a(\delta) \leq b(\delta)$ , both

$$a = \lim_{\delta \rightarrow 0} a(\delta) \quad \text{and} \quad b = \lim_{\delta \rightarrow 0} b(\delta)$$

exist. If we can show that  $a = b$  then it follows that  $\lim_{x \rightarrow 0} f(x) = a$ . We bound

$$\begin{aligned} b(\delta) - a(\delta) &= \sup\{f(x) : 0 < x < \delta\} - \inf\{f(y) : 0 < y < \delta\} \\ &= \sup\{f(x) - f(y) : 0 < x, y < \delta\} \leq \varepsilon, \end{aligned}$$

so that for any  $\varepsilon > 0$  there exists a  $\delta > 0$  such that

$$b(\delta) \leq a(\delta) + \varepsilon.$$

But this implies that  $b(\delta) \leq a(\delta)$ , whence equality follows.

3) Let  $\alpha \in \mathbb{R}$  and  $f : [0, 1] \rightarrow \mathbb{R}$  be given by

$$f(x) = \begin{cases} x^\alpha & x \in \{1/k; k \in \mathbb{N}\} , \\ 0 & \text{else.} \end{cases}$$

For which values of  $\alpha$  is  $f$  Riemann-integrable? If  $f$  is Riemann-integrable, what is the value of  $\int_0^1 f(x) dx$ ?

Solution:

If  $\alpha < 0$  then  $f$  is unbounded and therefore not Riemann-integrable.

Let now  $\alpha \geq 0$ , so that  $f$  is bounded by 1.

As  $f$  is zero on all irrational numbers,  $L(f, P) = 0$  for all  $P \in \mathcal{P}$ , and thus

$$\int_{*0}^1 f(x) dx = 0 .$$

Consider the partition of  $[0, 1]$  by

$$P_n = \{0, n/n^2, (n+1)/n^2, \dots, (n^2-1)/n^2, n^2/n^2\}$$

into one interval of width  $1/n$  and  $n^2 - n$  intervals of width  $1/n^2$ . (Many other choices would work here, as well.)

For  $x \geq 1/n$ ,  $f(x)$  is non-zero at precisely  $n$  points, so that  $\sup_{x \in I_i} f(x)$  is non-zero on the left-most interval of width  $1/n$  and at most  $2n$  intervals of width  $1/n^2$ . Thus,

$$U(f, P_n) \leq \frac{1}{n} + 2n \frac{1}{n^2} = \frac{3}{n} .$$

We thus have

$$0 = L(f, P_n) \leq U(f, P_n) \leq \frac{3}{n}$$

so that the  $f$  is Riemann-integrable and  $\int_0^1 f(x) dx = 0$ .

4) Let  $f : [a, b] \rightarrow \mathbb{R}$  be Riemann-integrable and  $c \in \mathbb{R}$ .

(a) Given a partition  $P$  of  $[a, b]$ , show that

$$U(cf, P) - L(cf, P) \leq |c|(U(f, P) - L(f, P)) .$$

(b) Deduce from (a) that  $cf$  is integrable and  $\int_a^b cf(x) dx = c \int_a^b f(x) dx$ . [This completes the proof of Theorem 7.4.]

Solution:

(a) For  $c \geq 0$ ,

$$\sup_{x \in I_i} cf(x) = c \sup_{x \in I_i} f(x) \quad \text{and} \quad \inf_{x \in I_i} cf(x) = c \inf_{x \in I_i} f(x) ,$$

so that

$$\sup_{x \in I_i} cf(x) - \inf_{x \in I_i} cf(x) = c \left( \sup_{x \in I_i} f(x) - \inf_{x \in I_i} f(x) \right) .$$

For  $c \leq 0$  this changes to

$$\sup_{x \in I_i} cf(x) = c \inf_{x \in I_i} f(x) \quad \text{and} \quad \inf_{x \in I_i} cf(x) = c \sup_{x \in I_i} f(x) .$$

so that

$$\sup_{x \in I_i} cf(x) - \inf_{x \in I_i} cf(x) = -c \left( \sup_{x \in I_i} f(x) - \inf_{x \in I_i} f(x) \right) .$$

Taken together, this implies

$$\sup_{x \in I_i} cf(x) - \inf_{x \in I_i} cf(x) = |c| \left( \sup_{x \in I_i} f(x) - \inf_{x \in I_i} f(x) \right) .$$

Multiplying by  $\Delta x_i$  and summing over all  $i$  gives the desired result.

(b) If  $U(f, P) - L(f, P) < \epsilon$  for some  $\epsilon > 0$ , then also

$$U(cf, P) - L(cf, P) \leq |c|(U(f, P) - L(f, P)) < |c|\epsilon .$$

By Riemann's integrability criterion,  $cf$  is integrable.

Finally, for  $c \geq 0$  we have

$$L(cf, P) = cL(f, P) \leq c \int_a^b f(x) dx \leq cU(f, P) = U(cf, P)$$

and for  $c \leq 0$  we have

$$L(cf, P) = cU(f, P) \leq c \int_a^b f(x) dx \leq cL(f, P) = U(cf, P)$$

so that in both cases

$$\int_a^b cf(x) dx = c \int_a^b f(x) dx$$

follows.