

# Simulating models of polymer collapse

Thomas Prellberg

School of Mathematical Sciences  
Queen Mary, University of London

COVLAT06


June 29 – July 1, 2006

- Polymers in solution:
  - Equilibrium statistical mechanics, lattice models, exponents
- Algorithm:
  - Stochastic growth & flat histogram (PERM/flatPERM)
- Simulations and results:
  - Interacting self-avoiding walks/trails (ISAW/ISAT)
  - Site-weighted random walks (SWRW): a tale of surprises

# Polymers in Solution


# Modelling of Polymers in Solution

- Polymers:  
long chains of monomers
- “Coarse-Graining”:  
beads on a chain
- “Excluded Volume”:  
minimal distance between beads
- Contact with solvent:  
effective short-range interaction
- Good/bad solvent:  
repelling/attracting interaction



# Modelling of Polymers in Solution

- Polymers:  
long chains of monomers
- “Coarse-Graining”:  
beads on a chain
- “Excluded Volume”:  
minimal distance between beads
- Contact with solvent:  
effective short-range interaction
- Good/bad solvent:  
repelling/attracting interaction




## A Model of a Polymer in Solution

Random Walk + Excluded Volume + Short Range Attraction

# Polymer Collapse, Coil-Globule Transition, $\Theta$ -Point

length  $N$ , spatial extension  $R \sim N^\nu$



$T > T_c$ : good solvent  
swollen phase (coil)



$T = T_c$ :  
 $\Theta$ -polymer

$T < T_c$ : bad solvent  
collapsed phase (globule)



# Critical Exponents

Length scale exponent  $\nu$ :  $R_N \sim N^\nu$

d	Coil	$\Theta$	Globule
2	$3/4$	$4/7$	$1/2$
3	$0.587 \dots$	$1/2(\log)$	$1/3$
4	$1/2(\log)$	$1/2$	$1/4$

# Critical Exponents

Length scale exponent  $\nu$ :  $R_N \sim N^\nu$

d	Coil	$\Theta$	Globule
2	$3/4$	$4/7$	$1/2$
3	$0.587 \dots$	$1/2(\log)$	$1/3$
4	$1/2(\log)$	$1/2$	$1/4$

Entropic exponent  $\gamma$ :  $Z_N \sim \mu^N N^{\gamma-1}$

d	Coil	$\Theta$	Globule
2	$43/32$	$8/7$	different scaling form $Z_N \sim \mu^N \mu_s^{N^\sigma} N^{\gamma-1}$ $\sigma = (d-1)/d$ (surface)
3	$1.15 \dots$	$1(\log)$	
4	$1(\log)$	$1$	



# Crossover Scaling at the $\Theta$ -Point

Crossover exponent  $\phi$

$$R_N \sim N^\nu \mathcal{R}(N^\phi \Delta T)$$

$$Z_N \sim \mu^N N^{\gamma-1} \mathcal{Z}(N^\phi \Delta T)$$

Specific heat of  $Z_N$  at  $T = T_c$ :  $C_N \sim N^{\alpha\phi}$

$$2 - \alpha = 1/\phi \quad \text{tri-critical scaling relation}$$

# Crossover Scaling at the $\Theta$ -Point

Crossover exponent  $\phi$

$$R_N \sim N^\nu \mathcal{R}(N^\phi \Delta T)$$

$$Z_N \sim \mu^N N^{\gamma-1} \mathcal{Z}(N^\phi \Delta T)$$

Specific heat of  $Z_N$  at  $T = T_c$ :  $C_N \sim N^{\alpha\phi}$

$$2 - \alpha = 1/\phi \quad \text{tri-critical scaling relation}$$

*Poor man's mean field theory of the  $\Theta$ -Point for  $d \geq 3$*

Balance between “excluded volume” and attractive interaction

⇒ polymer behaves like random walk:  $\nu = 1/2$ ,  $\gamma = 1$

⇒ weak thermodynamic phase transition  $\alpha = 0$ , i.e.  $\phi = 1/2$

# Crossover Scaling at the $\Theta$ -Point

Crossover exponent  $\phi$

$$R_N \sim N^\nu \mathcal{R}(N^\phi \Delta T)$$

$$Z_N \sim \mu^N N^{\gamma-1} \mathcal{Z}(N^\phi \Delta T)$$

Specific heat of  $Z_N$  at  $T = T_c$ :  $C_N \sim N^{\alpha\phi}$

$$2 - \alpha = 1/\phi \quad \text{tri-critical scaling relation}$$

Poor man's mean field theory of the  $\Theta$ -Point for  $d \geq 3$

Balance between “excluded volume” and attractive interaction

⇒ polymer behaves like random walk:  $\nu = 1/2$ ,  $\gamma = 1$


⇒ weak thermodynamic phase transition  $\alpha = 0$ , i.e.  $\phi = 1/2$

d	2	3	4
$\phi$	$3/7$	$1/2(\log)$	$1/2$

# The Canonical Lattice Model

## *Interacting Self-Avoiding Walk (ISAW)*

- Physical space  $\rightarrow$  simple cubic lattice  $\mathbb{Z}^3$
- Polymer  $\rightarrow$  self-avoiding random walk (SAW)
- Quality of solvent  $\rightarrow$  short-range interaction  $\epsilon$



# The Canonical Lattice Model


## Interacting Self-Avoiding Walk (ISAW)

- Physical space  $\rightarrow$  simple cubic lattice  $\mathbb{Z}^3$
- Polymer  $\rightarrow$  self-avoiding random walk (SAW)
- Quality of solvent  $\rightarrow$  short-range interaction  $\epsilon$

Partition function:

$$Z_N(\omega) = \sum_m C_{N,m} \omega^m$$

$C_{N,m}$  is the number of SAWs  
with  $N$  steps and  $m$  interactions



# The Canonical Lattice Model


## Interacting Self-Avoiding Walk (ISAW)

- Physical space  $\rightarrow$  simple cubic lattice  $\mathbb{Z}^3$
- Polymer  $\rightarrow$  self-avoiding random walk (SAW)
- Quality of solvent  $\rightarrow$  short-range interaction  $\epsilon$

Partition function:

$$Z_N(\omega) = \sum_m C_{N,m} \omega^m$$

$C_{N,m}$  is the number of SAWs  
with  $N$  steps and  $m$  interactions



### Thermodynamic Limit for a dilute solution:

$$V = \infty \quad \text{and} \quad N \rightarrow \infty$$

- Theoretical results from e.g.
  - $d = 2$ : Coulomb gas methods, conformal invariance, SLE, ...
  - $d \geq 3$ : self-consistent mean field theory
  - field theory:  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

# Theory and Models

- Theoretical results from e.g.
  - $d = 2$ : Coulomb gas methods, conformal invariance, SLE, ...
  - $d \geq 3$ : self-consistent mean field theory
  - field theory:  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

## A Model of a Polymer in Solution

Random Walk + Excluded Volume + Short Range Attraction




- Theoretical results from e.g.
  - $d = 2$ : Coulomb gas methods, conformal invariance, SLE, ...
  - $d \geq 3$ : self-consistent mean field theory
  - field theory:  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

## A Model of a Polymer in Solution

Random Walk + Excluded Volume + Short Range Attraction


- Canonical model: interacting self-avoiding walks (ISAW)
- Alternative model: interacting self-avoiding trails (ISAT)  
vertex avoidance (walks)  $\Leftrightarrow$  edge avoidance (trails)




nearest-neighbour interaction  $\Leftrightarrow$  contact interaction

# ISAW versus ISAT

ISAW




ISAT




# ISAW versus ISAT

ISAW




ISAT




- simulations of ISAW confirm the theoretical predictions
- simulations of ISAT confound the theoretical predictions

# ISAW versus ISAT

ISAW



ISAT



- simulations of ISAW confirm the theoretical predictions
- simulations of ISAT confound the theoretical predictions

Length scale exponent  $\nu$  for  $\mathbb{Z}^2$ :

Model	Coil	$\Theta$	Globule
ISAW	$3/4$	$4/7$	$1/2$
ISAT	$3/4$	$1/2(\log)$	$1/2$

Entropic exponent  $\gamma$  for  $\mathbb{Z}^2$ :

Model	Coil	$\Theta$
ISAW	$43/32$	$8/7$
ISAT	$43/32$	$1(\log)$

Crossover exponent  $\phi$  for  $\mathbb{Z}^2$ :

Model	
ISAW	$3/7$
ISAT	$0.84(3)$



# Simulations of ISAT

- At critical  $T_c$ , ISAT can be modelled as kinetic growth; simulations up to  $N = 10^6$

AL Owczarek and T Prellberg, J. Stat. Phys. 79 (1995) 951-967

- Pruned Enriched Rosenbluth Method enables simulations for  $T \neq T_c$ ; new simulations up to  $N = 2 \cdot 10^6$

AL Owczarek and T Prellberg, Physica A, in print




# ISAW versus ISAT

- On the square lattice,  
 $\text{SAW} = \text{SAT}$  but  
 $\text{ISAW} \neq \text{ISAT}$

# ISAW versus ISAT

- On the square lattice,  $\text{SAW} = \text{SAT}$  but  $\text{ISAW} \neq \text{ISAT}$
- On the diamond lattice, ISAT shows a bimodal distribution characteristic of a first-order transition, and at  $T_c$  (left peak) one finds purely Gaussian behaviour




T Prellberg and AL Owczarek, Phys. Rev. E 51 (1995) 2142-214

(figure from) P Grassberger and R Hegger, J. Phys. A 29 (1996) 279-288

# ISAW versus ISAT

- On the square lattice,  $\text{SAW} = \text{SAT}$  but  $\text{ISAW} \neq \text{ISAT}$
- On the diamond lattice, ISAT shows a bimodal distribution characteristic of a first-order transition, and at  $T_c$  (left peak) one finds purely Gaussian behaviour



T Prellberg and AL Owczarek, Phys. Rev. E 51 (1995) 2142-214

(figure from) P Grassberger and R Hegger, J. Phys. A 29 (1996) 279-288

*10 years later, this is still not understood!*



# A Proposal of a New Model

- ISAW/ISAT contain on-site and nearest-neighbour interactions
- The field-theory is formulated with purely local interactions
- Field theory is equivalent to Edwards model:
  - Brownian motion + suppression of self-intersections + attractive interactions
  - field theory is  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

# A Proposal of a New Model

- ISAW/ISAT contain on-site and nearest-neighbour interactions
- The field-theory is formulated with purely local interactions
- Field theory is equivalent to Edwards model:
  - Brownian motion + suppression of self-intersections + attractive interactions
  - field theory is  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

*Formulate a lattice model with purely local interactions*

# A Proposal of a New Model

- ISAW/ISAT contain on-site and nearest-neighbour interactions
- The field-theory is formulated with purely local interactions
- Field theory is equivalent to Edwards model:
  - Brownian motion + suppression of self-intersections + attractive interactions
  - field theory is  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

*Formulate a lattice model with purely local interactions*

- Site-weighted random walk:
  - lattice random walk weighted by multiple visits of sites
  - few visits to same site are favoured (attractive interaction)
  - too many visits are disfavoured (excluded volume)

# A Proposal of a New Model

- ISAW/ISAT contain on-site and nearest-neighbour interactions
- The field-theory is formulated with purely local interactions
- Field theory is equivalent to Edwards model:
  - Brownian motion + suppression of self-intersections + attractive interactions
  - field theory is  $\phi^4 - \phi^6$   $O(n)$ -model for  $n \rightarrow 0$

*Formulate a lattice model with purely local interactions*

- Site-weighted random walk:
  - lattice random walk weighted by multiple visits of sites
  - few visits to same site are favoured (attractive interaction)
  - too many visits are disfavoured (excluded volume)

(technically, this is an extension of a Domb-Joyce model)

# Site-Weighted Random Walk

- An  $N$ -step random walk  $\xi = (\vec{\xi}_0, \vec{\xi}_1, \dots, \vec{\xi}_N)$  induces a density-field  $\phi_\xi$  on the lattice sites  $\vec{x}$  via

$$\phi_\xi(\vec{x}) = \sum_{i=0}^N \delta_{\vec{\xi}_i, \vec{x}}$$

# Site-Weighted Random Walk

- An  $N$ -step random walk  $\xi = (\vec{\xi}_0, \vec{\xi}_1, \dots, \vec{\xi}_N)$  induces a density-field  $\phi_\xi$  on the lattice sites  $\vec{x}$  via

$$\phi_\xi(\vec{x}) = \sum_{i=0}^N \delta_{\vec{\xi}_i, \vec{x}}$$

- Define the energy as a functional of the field  $\phi = \phi_\xi$

$$E(\xi) = \sum_{\vec{x}} f(\phi(\vec{x}))$$

# Site-Weighted Random Walk

- An  $N$ -step random walk  $\xi = (\vec{\xi}_0, \vec{\xi}_1, \dots, \vec{\xi}_N)$  induces a density-field  $\phi_\xi$  on the lattice sites  $\vec{x}$  via

$$\phi_\xi(\vec{x}) = \sum_{i=0}^N \delta_{\vec{\xi}_i, \vec{x}}$$

- Define the energy as a functional of the field  $\phi = \phi_\xi$


$$E(\xi) = \sum_{\vec{x}} f(\phi(\vec{x}))$$

- Incorporate self-avoidance and attraction via choice of  $f(t)$ .  
For example,  $f(0) = f(1) = 0$ ,

$$f(2) = \varepsilon_1, \quad f(3) = \varepsilon_2,$$


and  $f(t \geq 4) = \infty$ .

# Site-Weighted Random Walk (ctd)





# Site-Weighted Random Walk (ctd)




- Partition function

$$Z_N(\beta) = \sum_{m_1, m_2} C_{N, m_1, m_2} e^{-\beta(m_1 \varepsilon_1 + m_2 \varepsilon_2)}$$

with density of states  $C_{N, m_1, m_2}$

# Site-Weighted Random Walk (ctd)



- Partition function

$$Z_N(\beta) = \sum_{m_1, m_2} C_{N, m_1, m_2} e^{-\beta(m_1 \varepsilon_1 + m_2 \varepsilon_2)}$$

with density of states  $C_{N, m_1, m_2}$

- Simulate two variants of the model on the square and simple cubic lattice
  - random walks with immediate reversal allowed (RA2, RA3)
  - random walks with immediate reversal forbidden (RF2, RF3)





# The Algorithm and Simulations

# PERM: “Go With The Winners”

PERM = Pruned and Enriched Rosenbluth Method

Grassberger, Phys Rev E 56 (1997) 3682

- Rosenbluth Method: kinetic growth




# PERM: “Go With The Winners”

PERM = Pruned and Enriched Rosenbluth Method

Grassberger, Phys Rev E 56 (1997) 3682

- Rosenbluth Method: kinetic growth




- Enrichment: weight too large  $\rightarrow$  make copies of configuration
- Pruning: weight too small  $\rightarrow$  remove configuration occasionally

# PERM: “Go With The Winners”

PERM = Pruned and Enriched Rosenbluth Method

Grassberger, Phys Rev E 56 (1997) 3682

- Rosenbluth Method: kinetic growth



- Enrichment: weight too large  $\rightarrow$  make copies of configuration
- Pruning: weight too small  $\rightarrow$  remove configuration occasionally


State-of-the-art: flatPERM = flat histogram PERM

M Bachmann and W Janke, PRL 91 (2003) 208105

T Prellberg and J Krawczyk, PRL 92 (2004) 120602


- flatPERM samples a generalised multicanonical ensemble
- Determines the whole density of states in *one* simulation!

# SWRW simulations




- Four simulations: reversal allowed/forbidden, 2d/3d

# SWRW simulations




- Four simulations: reversal allowed/forbidden, 2d/3d
- Density of states  $C_{N,m_1,m_2}$  accessible up to  $N = 256$





- Four simulations: reversal allowed/forbidden, 2d/3d
- Density of states  $C_{N,m_1,m_2}$  accessible up to  $N = 256$
- Perform partial summation, e.g. over  $m_2$

$$\bar{C}_{N,m_1}(\beta_2) = \sum_{m_2} C_{N,m_1,m_2} e^{\beta_2 m_2}$$




- Four simulations: reversal allowed/forbidden, 2d/3d
- Density of states  $C_{N,m_1,m_2}$  accessible up to  $N = 256$
- Perform partial summation, e.g. over  $m_2$

$$\bar{C}_{N,m_1}(\beta_2) = \sum_{m_2} C_{N,m_1,m_2} e^{\beta_2 m_2}$$


- Density of states  $\bar{C}_{N,m_1}(\beta_2)$  accessible up to  $N = 1024$  (for  $\beta_2$  fixed)

# SWRW in 3d, reversal forbidden (RF3)




Phase diagram

# SWRW in 3d, reversal forbidden (RF3)




Phase diagram

$$\beta_2 = -1.0:$$




2nd order transition

$$\beta_1 = -1.0:$$




1st order transition

# SWRW in 3d, reversal forbidden (RF3)




Phase diagram




bimodal distribution

$\beta_2 = -1.0$ :



2nd order transition



$\beta_1 = -1.0$ :



1st order transition

# SWRW in 2d, reversal allowed (RA2)



We find a smooth crossover:



*Both 1st order and 2nd order transitions have disappeared!*

# SWRW in 2d, reversal allowed (RA2)

We find a smooth crossover:



*Both 1st order and 2nd order transitions have disappeared!*

## RA3 and RF2

2nd order transition disappears as in RA2

1st order transition weakens

# SWRW summarised

Model	2d	3d
RA	no transitions	one transition
RF	one transition	two transitions



Model	2d	3d
RA	no transitions	one transition
RF	one transition	two transitions

## Unexpected and intriguing behaviour

Changing the dimension and/or allowing reversals removes the phase transition

Model	2d	3d
RA	no transitions	one transition
RF	one transition	two transitions

## Unexpected and intriguing behaviour

Changing the dimension and/or allowing reversals removes the phase transition

Many open Questions remain ...

- Polymers in solution:
  - Random Walk + Excluded Volume + Attraction?

# Summary

- Polymers in solution:
  - Random Walk + Excluded Volume + Attraction?
- Algorithm:
  - Stochastic growth & flat histogram (PERM/flatPERM)

- Polymers in solution:
  - Random Walk + Excluded Volume + Attraction?
- Algorithm:
  - Stochastic growth & flat histogram (PERM/flatPERM)
- Simulations and results:
  - ISAW versus ISAT
  - Site-weighted random walks (SWRW)

# Summary

- Polymers in solution:
  - Random Walk + Excluded Volume + Attraction?
- Algorithm:
  - Stochastic growth & flat histogram (PERM/flatPERM)
- Simulations and results:
  - ISAW versus ISAT
  - Site-weighted random walks (SWRW)

*An unfinished story!*

Joined work with A.L. Owczarek, A. Rechnitzer, J. Krawczyk

- The algorithm:

- T. Prellberg and J. Krawczyk, "Flat histogram version of the pruned and enriched Rosenbluth method," Phys. Rev. Lett. 92 (2004) 120602; selected for Virt. J. Biol. Phys. Res. 7 (2004)
- T. Prellberg, J. Krawczyk, and A. Rechnitzer, "Polymer simulations with a flat histogram stochastic growth algorithm," Computer Simulation Studies in Condensed Matter Physics XVII, pages 122-135, Springer Verlag, 2006

Joined work with A.L. Owczarek, A. Rechnitzer, J. Krawczyk

- The algorithm:

- T. Prellberg and J. Krawczyk, "Flat histogram version of the pruned and enriched Rosenbluth method," Phys. Rev. Lett. 92 (2004) 120602; selected for Virt. J. Biol. Phys. Res. 7 (2004)
- T. Prellberg, J. Krawczyk, and A. Rechnitzer, "Polymer simulations with a flat histogram stochastic growth algorithm," Computer Simulation Studies in Condensed Matter Physics XVII, pages 122-135, Springer Verlag, 2006

- Current simulations:

- A. L. Owczarek and T. Prellberg, "Collapse transition of self-avoiding trails on the square lattice," Physica A, in print
- J. Krawczyk, T. Prellberg, A. L. Owczarek, and A. Rechnitzer, "Self-avoiding random walk with multiple site weightings and restrictions," Phys. Rev. Lett. 96 (2006) 240603



Joined work with A.L. Owczarek, A. Rechnitzer, J. Krawczyk

- The algorithm:

- T. Prellberg and J. Krawczyk, "Flat histogram version of the pruned and enriched Rosenbluth method," Phys. Rev. Lett. 92 (2004) 120602; selected for Virt. J. Biol. Phys. Res. 7 (2004)
- T. Prellberg, J. Krawczyk, and A. Rechnitzer, "Polymer simulations with a flat histogram stochastic growth algorithm," Computer Simulation Studies in Condensed Matter Physics XVII, pages 122-135, Springer Verlag, 2006

- Current simulations:

- A. L. Owczarek and T. Prellberg, "Collapse transition of self-avoiding trails on the square lattice," Physica A, in print
- J. Krawczyk, T. Prellberg, A. L. Owczarek, and A. Rechnitzer, "Self-avoiding random walk with multiple site weightings and restrictions," Phys. Rev. Lett. 96 (2006) 240603

- Things to come:

- J. Krawczyk, A. L. Owczarek, T. Prellberg, and A. Rechnitzer, "Simulation of Lattice Polymers with Hydrogen-Like Bonding," preprint

# The End