

MTH5105 Differential and Integral Analysis

2010-2011

Solutions 3

1 Exercises for Feedback

- 1) The functions \sinh and \cosh are given by

$$\begin{aligned}\sinh : \mathbb{R} &\rightarrow \mathbb{R}, & x &\mapsto \frac{1}{2}(\exp(x) - \exp(-x)), \\ \cosh : \mathbb{R} &\rightarrow \mathbb{R}, & x &\mapsto \frac{1}{2}(\exp(x) + \exp(-x)).\end{aligned}$$

- (a) Prove that \sinh and \cosh are differentiable and that $\sinh' = \cosh$ and $\cosh' = \sinh$.
(b) Prove that the function

$$f(x) = \cosh^2(x) - \sinh^2(x)$$

is constant by considering $f'(x)$.
What is the value of the constant?

- (c) Prove that \sinh is invertible.
(d) Prove that $\sinh(\mathbb{R}) = \mathbb{R}$. *Hint: show that $\sinh(2x) > x$ for $x > 0$, and mimic the proof of the statement that $\exp(\mathbb{R}) = \mathbb{R}^+$.*
(e) Prove that $\operatorname{arsinh} = \sinh^{-1}$ is differentiable, and that

$$\operatorname{arsinh}'(x) = \frac{1}{\sqrt{1+x^2}}.$$

Solution:

- (a) \exp is differentiable, therefore \sinh and \cosh are differentiable. Using $\exp' = \exp$, the derivatives follow immediately.
(b) f is differentiable, and $f'(x) = 2\cosh(x)\sinh(x) - 2\sinh(x)\cosh(x) = 0$. By Theorem 2.5, f is constant. $f(0) = \cosh^2(0) - \sinh^2(0) = 1$, so $\cosh^2(x) - \sinh^2(x) = 1$.
(c) $\sinh'(x) = \cosh(x) > 0$ for all $x \in \mathbb{R}$, therefore \sinh is strictly increasing by Theorem 2.4, and therefore invertible by the corollary after Theorem 4.2.
(d) Let $x > 0$. We have $\exp(x) > 1 + x$ (see proof of Theorem 3.3) and $\exp(-x) < 1$ (since \exp is strictly increasing, and $\exp(0) = 1$), so that $\sinh(x) > x/2$.
Let $c > 0$. From

$$\sinh(0) = 0 < c < \sinh(2c)$$

it follows by the IVT applied to the interval $[0, 2c]$, that there exists an $x \in (0, 2c)$ such that $\sinh(x) = c$. A similar argument holds for $c < 0$, and for $c = 0$ we have $\sinh(0) = 0 = c$.

- (e) $\sinh'(x) = \cosh(x) > 0$ for all $x \in \mathbb{R}$, therefore by Theorem 4.6, arsinh is differentiable and

$$\operatorname{arsinh}'(x) = \frac{1}{\cosh(\operatorname{arsinh}(x))}.$$

Now $\cosh(x) = \sqrt{1 + \sinh^2(x)}$ (from (b), and the positive square root is taken as $\cosh(x)$ is positive), so that $\operatorname{arsinh}'(x) = 1/\sqrt{1+x^2}$.

2 Extra Exercises

- 2) (a) Find a bijective, continuously differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$ with $f'(0) = 0$ and a continuous inverse.
 (b) Let $f : \mathbb{R}_0^+ \rightarrow \mathbb{R}$ be differentiable and decreasing. Prove or disprove: If $\lim_{x \rightarrow 0} f(x) = 0$, then $\lim_{x \rightarrow 0} f'(x) = 0$.

Solution:

- (a) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = x^3$.
 f is differentiable with continuous derivative $f'(x) = 3x^2$. We have $f'(0) = 0$.
 The inverse is $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$, $x \mapsto x^{1/3}$.
 As f is strictly increasing on \mathbb{R} , f is injective. $f(\mathbb{R}) = \mathbb{R}$ implies that f is surjective as well, so f is bijective.
 As f is differentiable, it is continuous. Therefore f^{-1} is also continuous.
 (b) This can be disproved by a counterexample.
 Let $f : \mathbb{R}_0^+ \rightarrow \mathbb{R}$ be given by $f(x) = -x$.
 f is differentiable and $f'(x) = -1$ for $x \geq 0$.
 $\lim_{x \rightarrow 0} f(x) = 0$, but $\lim_{x \rightarrow 0} f'(x) = -1$.

- 3) Using the Intermediate Value Theorem, prove that a continuous function maps intervals to intervals.

Solution:

We use the following characterisation of an interval: $I \subseteq \mathbb{R}$ is an interval if and only if for all $x_1, x_2 \in I$ with $x_1 < x_2$,

$$x_1 < c < x_2 \Rightarrow c \in I.$$

Let $J = f(I)$. We need to show that J is an interval, i.e. for all $y_1, y_2 \in J$ with $y_1 < y_2$, $y_1 < c < y_2 \Rightarrow c \in J$:

Let $y_1, y_2 \in J$ with $y_1 < y_2$. Then there exist $x_1, x_2 \in I$ such that $y_1 = f(x_1)$ and $y_2 = f(x_2)$.
 As $y_1 \neq y_2$, necessarily $x_1 \neq x_2$ also, so either $x_1 < x_2$ or $x_2 < x_1$.

Consider, without loss of generality, the case $x_1 < x_2$. By assumption, f is a continuous function on I , so it is a continuous function on $[x_1, x_2]$ (or $[x_2, x_1]$, if $x_2 < x_1$).

Hence, by the intermediate value theorem, for all c with $y_1 < c < y_2$ there exists an $a \in [x_1, x_2]$ such that $f(a) = c$.

This implies that $c \in J$.

- 4*) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function, of which it is known that $f'(x)$ exists for all $x \neq 0$ and that $f'(x) \rightarrow 2$ as $x \rightarrow 0$. Does it follow that f is differentiable at 0? If yes, give a rigorous proof; if no, provide a counter-example.

Solution:

Given $x > 0$, f is continuous on $[0, x]$ and differentiable on $(0, x)$. Hence, by the Mean Value Theorem applied to f on $[0, x]$, there exists a $c \in (0, x)$ such that $f'(c) = \frac{f(x) - f(0)}{x - 0}$.

Similarly, if $x < 0$ then there exists a $c \in (x, 0)$ such that $f'(c) = \frac{f(x) - f(0)}{x - 0}$. Clearly in both cases $|c| < |x|$, and $c \rightarrow 0$ as $x \rightarrow 0$. Therefore

$$f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{c \rightarrow 0} f'(c) = 2$$

exists.

Thomas Prellberg, January 2011