


The Mathematics of the Casimir Effect

Thomas Prellberg

School of Mathematical Sciences
Queen Mary, University of London

Sigma Club
LSE Centre for Philosophy of Natural and Social Science
June 28, 2010



A Tiny Force of Nature Is Stronger Than Thought


New York Times, February 2001

Physicists Have 'Solved' Mystery of Gravitation

Telegraph, August 2007

Nanotechnology: Shaping The Void

Nature, August 2008



A Tiny Force of Nature Is Stronger Than Thought


New York Times, February 2001

Physicists Have 'Solved' Mystery of Gravitation

Telegraph, August 2007

Nanotechnology: Shaping The Void

Nature, August 2008



A Tiny Force of Nature Is Stronger Than Thought


New York Times, February 2001

Physicists Have 'Solved' Mystery of Gravitation

Telegraph, August 2007

Nanotechnology: Shaping The Void

Nature, August 2008



Much ado about nothing


The Economist, May 2008

The island possesses a Casimir effect

"Lost", Season Finale, May 2008

*Casimir Effect - a story of
unrequited love set on a backdrop of
temporal paradox*

Indie Film, Release June 2011



Much ado about nothing


The Economist, May 2008

The island possesses a Casimir effect

“Lost”, Season Finale, May 2008

*Casimir Effect - a story of
unrequited love set on a backdrop of
temporal paradox*

Indie Film, Release June 2011



Much ado about nothing

The Economist, May 2008

The island possesses a Casimir effect

“Lost”, Season Finale, May 2008

*Casimir Effect - a story of
unrequited love set on a backdrop of
temporal paradox*

Indie Film, Release June 2011

Topic Outline

- 1 The Casimir Effect
 - History
 - Quantum Electrodynamics
 - Zero-Point Energy Shift
- 2 Making Sense of Infinity - Infinity
 - The Mathematical Setting
 - Divergent Series
 - Euler-Maclaurin Formula
 - Abel-Plana Formula
- 3 Conclusion

Outline

- 1 The Casimir Effect
 - History
 - Quantum Electrodynamics
 - Zero-Point Energy Shift
- 2 Making Sense of Infinity - Infinity
- 3 Conclusion

- student of Ehrenfest, worked with Pauli and Bohr
- *retarded Van-der-Waals-forces*

$$E = \frac{23}{4\pi} \hbar c \frac{\alpha}{R^7}$$

Casimir and Polder, 1948

- *Force between cavity walls*

$$F = -\frac{\pi^2 \hbar c}{240} \frac{A}{d^4}$$

Casimir, 1948



Hendrik Brugt Gerhard
Casimir, 1909 - 2000

- student of Ehrenfest, worked with Pauli and Bohr
- *retarded Van-der-Waals-forces*

$$E = \frac{23}{4\pi} \hbar c \frac{\alpha}{R^7}$$

Casimir and Polder, 1948

- *Force between cavity walls*

$$F = -\frac{\pi^2 \hbar c}{240} \frac{A}{d^4}$$

Casimir, 1948



Hendrik Brugt Gerhard
Casimir, 1909 - 2000

- student of Ehrenfest, worked with Pauli and Bohr
- *retarded Van-der-Waals-forces*

$$E = \frac{23}{4\pi} \hbar c \frac{\alpha}{R^7}$$

Casimir and Polder, 1948

- *Force between cavity walls*

$$F = -\frac{\pi^2 \hbar c}{240} \frac{A}{d^4}$$

Casimir, 1948



Hendrik Brugt Gerhard
Casimir, 1909 - 2000

The Electromagnetic Field

- The electromagnetic field, described by the *Maxwell Equations*, satisfies the *wave equation*

$$\left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \vec{A}(\vec{x}, t) = 0$$

- Fourier-transformation ($\vec{x} \leftrightarrow \vec{k}$) gives

$$\left(\frac{\partial^2}{\partial t^2} + \omega^2 \right) \vec{A}(\vec{k}, t) = 0 \quad \text{with } \omega = c|\vec{k}|$$

which, for each \vec{k} , describes a **harmonic oscillator**

- Quantising harmonic oscillators is easy...

The Electromagnetic Field

- The electromagnetic field, described by the *Maxwell Equations*, satisfies the *wave equation*

$$\left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \vec{A}(\vec{x}, t) = 0$$

- Fourier-transformation ($\vec{x} \leftrightarrow \vec{k}$) gives

$$\left(\frac{\partial^2}{\partial t^2} + \omega^2 \right) \vec{A}(\vec{k}, t) = 0 \quad \text{with } \omega = c|\vec{k}|$$

which, for each \vec{k} , describes a **harmonic oscillator**

- Quantising harmonic oscillators is easy...

The Electromagnetic Field

- The electromagnetic field, described by the *Maxwell Equations*, satisfies the *wave equation*

$$\left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \vec{A}(\vec{x}, t) = 0$$

- Fourier-transformation ($\vec{x} \leftrightarrow \vec{k}$) gives

$$\left(\frac{\partial^2}{\partial t^2} + \omega^2 \right) \vec{A}(\vec{k}, t) = 0 \quad \text{with } \omega = c|\vec{k}|$$

which, for each \vec{k} , describes a **harmonic oscillator**

- Quantising harmonic oscillators is easy...

Quantisation of the Field

- Each harmonic oscillator can be in a discrete state of energy

$$E_m(\vec{k}) = \left(m + \frac{1}{2}\right) \hbar\omega \quad \text{with } \omega = c|\vec{k}|$$

- Interpretation: m photons with energy $\hbar\omega$ and momentum $\hbar\vec{k}$
- In particular, the ground state energy $\frac{1}{2}\hbar\omega$ is non-zero!
- This leads to a **zero-point energy** density of the field

$$\frac{E}{V} = 2 \int E_0(\vec{k}) \frac{d^3k}{(2\pi)^3}$$

(factor 2 due to polarisation of the field)

- **Caveat:** this quantity is infinite...

Quantisation of the Field

- Each harmonic oscillator can be in a discrete state of energy

$$E_m(\vec{k}) = \left(m + \frac{1}{2}\right) \hbar\omega \quad \text{with } \omega = c|\vec{k}|$$

- Interpretation: m photons with energy $\hbar\omega$ and momentum $\hbar\vec{k}$
- In particular, the ground state energy $\frac{1}{2}\hbar\omega$ is non-zero!
- This leads to a **zero-point energy** density of the field

$$\frac{E}{V} = 2 \int E_0(\vec{k}) \frac{d^3k}{(2\pi)^3}$$

(factor 2 due to polarisation of the field)

- **Caveat:** this quantity is infinite...

Quantisation of the Field

- Each harmonic oscillator can be in a discrete state of energy

$$E_m(\vec{k}) = \left(m + \frac{1}{2}\right) \hbar\omega \quad \text{with } \omega = c|\vec{k}|$$

- Interpretation: m photons with energy $\hbar\omega$ and momentum $\hbar\vec{k}$
- In particular, the ground state energy $\frac{1}{2}\hbar\omega$ is non-zero!
- This leads to a **zero-point energy** density of the field

$$\frac{E}{V} = 2 \int E_0(\vec{k}) \frac{d^3k}{(2\pi)^3}$$

(factor 2 due to polarisation of the field)

- **Caveat:** this quantity is infinite...

Quantisation of the Field

- Each harmonic oscillator can be in a discrete state of energy

$$E_m(\vec{k}) = \left(m + \frac{1}{2}\right) \hbar\omega \quad \text{with } \omega = c|\vec{k}|$$

- Interpretation: m photons with energy $\hbar\omega$ and momentum $\hbar\vec{k}$
- In particular, the ground state energy $\frac{1}{2}\hbar\omega$ is non-zero!
- This leads to a **zero-point energy** density of the field

$$\frac{E}{V} = 2 \int E_0(\vec{k}) \frac{d^3k}{(2\pi)^3}$$

(factor 2 due to polarisation of the field)

- **Caveat:** this quantity is infinite...

Quantisation of the Field

- Each harmonic oscillator can be in a discrete state of energy

$$E_m(\vec{k}) = \left(m + \frac{1}{2}\right) \hbar\omega \quad \text{with } \omega = c|\vec{k}|$$

- Interpretation: m photons with energy $\hbar\omega$ and momentum $\hbar\vec{k}$
- In particular, the ground state energy $\frac{1}{2}\hbar\omega$ is non-zero!
- This leads to a **zero-point energy** density of the field


$$\frac{E}{V} = 2 \int E_0(\vec{k}) \frac{d^3k}{(2\pi)^3}$$

(factor 2 due to polarisation of the field)

- **Caveat:** this quantity is infinite...

Making (Physical) Sense of Infinity

The zero-point energy shifts due to a restricted geometry



- In the presence of the boundary

$$E_{discrete} = \sum_n E_{0,n}$$

is a sum over discrete energies $E_{0,n} = \frac{1}{2} \hbar \omega_n$

- In the absence of a boundary

$$E = 2V \int E_0(\vec{k}) \frac{d^3 k}{(2\pi)^3}$$


The difference of the infinite zero-point energies is finite!

$$\Delta E = E_{discrete} - E = -\frac{\pi^2 \hbar c}{720} \frac{L^2}{d^3}$$

for a box of size $L \times L \times d$ with $d \ll L$

Making (Physical) Sense of Infinity

The zero-point energy shifts due to a restricted geometry



- In the presence of the boundary

$$E_{discrete} = \sum_n E_{0,n}$$

is a sum over discrete energies $E_{0,n} = \frac{1}{2} \hbar \omega_n$

- In the absence of a boundary

$$E = 2V \int E_0(\vec{k}) \frac{d^3 k}{(2\pi)^3}$$


The difference of the infinite zero-point energies is finite!

$$\Delta E = E_{discrete} - E = -\frac{\pi^2 \hbar c}{720} \frac{L^2}{d^3}$$

for a box of size $L \times L \times d$ with $d \ll L$

Making (Physical) Sense of Infinity

The zero-point energy shifts due to a restricted geometry



- In the presence of the boundary

$$E_{discrete} = \sum_n E_{0,n}$$

is a sum over discrete energies $E_{0,n} = \frac{1}{2} \hbar \omega_n$

- In the absence of a boundary

$$E = 2V \int E_0(\vec{k}) \frac{d^3 k}{(2\pi)^3}$$


The difference of the infinite zero-point energies is finite!

$$\Delta E = E_{discrete} - E = -\frac{\pi^2 \hbar c}{720} \frac{L^2}{d^3}$$

for a box of size $L \times L \times d$ with $d \ll L$

Making (Physical) Sense of Infinity

The zero-point energy shifts due to a restricted geometry



- In the presence of the boundary

$$E_{discrete} = \sum_n E_{0,n}$$

is a sum over discrete energies $E_{0,n} = \frac{1}{2} \hbar \omega_n$

- In the absence of a boundary

$$E = 2V \int E_0(\vec{k}) \frac{d^3 k}{(2\pi)^3}$$

The difference of the infinite zero-point energies is finite!

$$\Delta E = E_{discrete} - E = -\frac{\pi^2 \hbar c}{720} \frac{L^2}{d^3}$$

for a box of size $L \times L \times d$ with $d \ll L$

- Wolfgang Pauli's initial reaction: 'absolute nonsense'
- Experimental verification
 - Sparnaay (1958): 'not inconsistent with'
 - van Blokland and Overbeek (1978): experimental error of 50%
 - Lamoreaux (1997): experimental error of 5%
 - Chen (2008): experimental error $< 1\%$
- Theoretical extensions
 - Geometry dependence
 - Dynamical Casimir effect
 - Real media: non-zero temperature, finite conductivity, roughness, ...

- Wolfgang Pauli's initial reaction: 'absolute nonsense'
- Experimental verification
 - Sparnaay (1958): 'not inconsistent with'
 - van Blokland and Overbeek (1978): experimental error of 50%
 - Lamoreaux (1997): experimental error of 5%
 - Chen (2008): experimental error $< 1\%$
- Theoretical extensions
 - Geometry dependence
 - Dynamical Casimir effect
 - Real media: non-zero temperature, finite conductivity, roughness, ...

- Wolfgang Pauli's initial reaction: 'absolute nonsense'
- Experimental verification
 - Sparnaay (1958): 'not inconsistent with'
 - van Blokland and Overbeek (1978): experimental error of 50%
 - Lamoreaux (1997): experimental error of 5%
 - Chen (2008): experimental error $< 1\%$
- Theoretical extensions
 - Geometry dependence
 - Dynamical Casimir effect
 - Real media: non-zero temperature, finite conductivity, roughness, ...

Alternative Approaches

- Casimir (1948): QED Calculation
infinite vacuum zero-point energy due to fluctuations of QED field
- Lifshitz (1956): Macroscopic Theory
limiting case of fluctuations in dielectrics, no QED field
- Milloni and Shih (1992): QED with Normal Ordering
formal subtraction of vacuum zero-point energy
- Schwinger (1967): Source Theory
polarisation sources in plates, vacuum is void

We are done with the physics. Let's look at some mathematics!

Alternative Approaches

- Casimir (1948): QED Calculation
infinite vacuum zero-point energy due to fluctuations of QED field
- Lifshitz (1956): Macroscopic Theory
limiting case of fluctuations in dielectrics, no QED field
- Milloni and Shih (1992): QED with Normal Ordering
formal subtraction of vacuum zero-point energy
- Schwinger (1967): Source Theory
polarisation sources in plates, vacuum is void

We are done with the physics. Let's look at some mathematics!

Alternative Approaches

- Casimir (1948): QED Calculation
infinite vacuum zero-point energy due to fluctuations of QED field
- Lifshitz (1956): Macroscopic Theory
limiting case of fluctuations in dielectrics, no QED field
- Milloni and Shih (1992): QED with Normal Ordering
formal subtraction of vacuum zero-point energy
- Schwinger (1967): Source Theory
polarisation sources in plates, vacuum is void

We are done with the physics. Let's look at some mathematics!

Alternative Approaches

- Casimir (1948): QED Calculation
infinite vacuum zero-point energy due to fluctuations of QED field
- Lifshitz (1956): Macroscopic Theory
limiting case of fluctuations in dielectrics, no QED field
- Milloni and Shih (1992): QED with Normal Ordering
formal subtraction of vacuum zero-point energy
- Schwinger (1967): Source Theory
polarisation sources in plates, vacuum is void

We are done with the physics. Let's look at some mathematics!

Alternative Approaches

- Casimir (1948): QED Calculation
infinite vacuum zero-point energy due to fluctuations of QED field
- Lifshitz (1956): Macroscopic Theory
limiting case of fluctuations in dielectrics, no QED field
- Milloni and Shih (1992): QED with Normal Ordering
formal subtraction of vacuum zero-point energy
- Schwinger (1967): Source Theory
polarisation sources in plates, vacuum is void

We are done with the physics. Let's look at some mathematics!

Outline

- 1 The Casimir Effect
- 2 Making Sense of Infinity - Infinity
 - The Mathematical Setting
 - Divergent Series
 - Euler-Maclaurin Formula
 - Abel-Plana Formula
- 3 Conclusion

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally

$$E_{\text{discrete}} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today

- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally

$$E_{\text{discrete}} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today

- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally

$$E_{discrete} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today

- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally

$$E_{discrete} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today

- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally

$$E_{discrete} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today

- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.

Spectral Theory

- Consider $-\Delta$ for a compact manifold Ω with a smooth boundary $\partial\Omega$
- On a suitable function space, this operator is self-adjoint and positive with pure point spectrum
- One finds formally


$$E_{discrete} = \frac{1}{2} \hbar c \text{Trace}(-\Delta)^{1/2}$$

This would be a different talk — let's keep it simple for today


- Choose

$$\Omega = [0, L] \quad \text{and} \quad \Delta = \frac{\partial^2}{\partial x^2}$$

with Dirichlet boundary conditions $f(0) = f(L) = 0$.



Casimir Effect in One Dimension



- The solutions are standing waves with wavelength λ satisfying

$$n \frac{\lambda}{2} = L$$


- We therefore find

$$E_{0,n} = \frac{1}{2} \hbar c \frac{n\pi}{L}$$

- The zero-point energies are given by

$$E_{discrete} = \frac{\pi}{2L} \hbar c \sum_{n=0}^{\infty} n \quad \text{and} \quad E = \frac{\pi}{2L} \hbar c \int_0^{\infty} t \, dt$$

Casimir Effect in One Dimension



- The solutions are standing waves with wavelength λ satisfying

$$n \frac{\lambda}{2} = L$$


- We therefore find

$$E_{0,n} = \frac{1}{2} \hbar c \frac{n\pi}{L}$$

- The zero-point energies are given by

$$E_{discrete} = \frac{\pi}{2L} \hbar c \sum_{n=0}^{\infty} n \quad \text{and} \quad E = \frac{\pi}{2L} \hbar c \int_0^{\infty} t \, dt$$

Casimir Effect in One Dimension



- The solutions are standing waves with wavelength λ satisfying

$$n \frac{\lambda}{2} = L$$

- We therefore find

$$E_{0,n} = \frac{1}{2} \hbar c \frac{n\pi}{L}$$

- The zero-point energies are given by

$$E_{discrete} = \frac{\pi}{2L} \hbar c \sum_{n=0}^{\infty} n \quad \text{and} \quad E = \frac{\pi}{2L} \hbar c \int_0^{\infty} t \, dt$$

The Mathematical Problem

- We need to make sense of

$$\Delta E = E_{discrete} - E = \frac{\pi}{2L} \hbar c \left(\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt \right)$$

- More generally, consider

$$\Delta(f) = \sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt$$

The Mathematical Problem

- We need to make sense of

$$\Delta E = E_{discrete} - E = \frac{\pi}{2L} \hbar c \left(\sum_{n=0}^{\infty} n - \int_0^{\infty} t \, dt \right)$$

- More generally, consider

$$\Delta(f) = \sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) \, dt$$

Divergent Series

On the Whole, Divergent Series are the Works of the Devil and it's a Shame that one dares base any Demonstration upon them. You can get whatever result you want when you use them, and they have given rise to so many Disasters and so many Paradoxes. Can anything more horrible be conceived than to have the following oozing out at you:

$$0 = 1 - 2^n + 3^n - 4^n + \text{etc.}$$

where n is an integer number?

Niels Henrik Abel

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = 1 - 1 + 1 - 1 + 1 - 1 + \dots$$

- Cesaro summation: let $S_N = \sum_{n=0}^N (-1)^n$ and compute

$$S = \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N S_N = \frac{1}{2}$$

- Abel summation:

$$S = \lim_{x \rightarrow 1^-} \sum_{n=0}^{\infty} (-1)^n x^n = \frac{1}{2}$$

- Borel summation, Euler summation, ...: again $S = \frac{1}{2}$

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = 1 - 1 + 1 - 1 + 1 - 1 + \dots$$

- Cesaro summation: let $S_N = \sum_{n=0}^N (-1)^n$ and compute

$$S = \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N S_N = \frac{1}{2}$$

- Abel summation:

$$S = \lim_{x \rightarrow 1^-} \sum_{n=0}^{\infty} (-1)^n x^n = \frac{1}{2}$$

- Borel summation, Euler summation, ...: again $S = \frac{1}{2}$

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = 1 - 1 + 1 - 1 + 1 - 1 + \dots$$

- Cesaro summation: let $S_N = \sum_{n=0}^N (-1)^n$ and compute

$$S = \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N S_N = \frac{1}{2}$$

- Abel summation:

$$S = \lim_{x \rightarrow 1^-} \sum_{n=0}^{\infty} (-1)^n x^n = \frac{1}{2}$$

- Borel summation, Euler summation, ...: again $S = \frac{1}{2}$

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = 1 - 1 + 1 - 1 + 1 - 1 + \dots$$

- Cesaro summation: let $S_N = \sum_{n=0}^N (-1)^n$ and compute

$$S = \lim_{N \rightarrow \infty} \frac{1}{N+1} \sum_{n=0}^N S_N = \frac{1}{2}$$

- Abel summation:

$$S = \lim_{x \rightarrow 1^-} \sum_{n=0}^{\infty} (-1)^n x^n = \frac{1}{2}$$

- Borel summation, Euler summation, ...: again $S = \frac{1}{2}$

An Aside: Madelung's Constant M

- The energy of a single ion in a large NaCl crystal

$$E = -\frac{z^2 e^2}{4\pi\epsilon_0 r_0} M$$

where

$$M = \sum'_{i,j,k} \frac{(-1)^{i+j+k}}{\sqrt{i^2 + j^2 + k^2}}$$

- Evaluate M by summing over spheres with increasing radius

$$M = -6 + 12/\sqrt{2} - 8/\sqrt{3} + 6/2 - 24/\sqrt{5} + \dots$$

Unfortunately, the resulting series *diverges*!

- Evaluate M by summing over cubes with increasing size converges

$$M = -1.74756\dots$$

- Several convergent series for M are known, the simplest being

$$M = -12\pi \sum_{m,n=1,3,5\dots} \operatorname{sech}^2 \left(\frac{1}{2} \pi \sqrt{m^2 + n^2} \right)$$

An Aside: Madelung's Constant M

- The energy of a single ion in a large NaCl crystal

$$E = -\frac{z^2 e^2}{4\pi\epsilon_0 r_0} M$$

where

$$M = \sum'_{i,j,k} \frac{(-1)^{i+j+k}}{\sqrt{i^2 + j^2 + k^2}}$$

- Evaluate M by summing over spheres with increasing radius

$$M = -6 + 12/\sqrt{2} - 8/\sqrt{3} + 6/2 - 24/\sqrt{5} + \dots$$

Unfortunately, the resulting series *diverges*!

- Evaluate M by summing over cubes with increasing size converges

$$M = -1.74756\dots$$

- Several convergent series for M are known, the simplest being

$$M = -12\pi \sum_{m,n=1,3,5\dots} \operatorname{sech}^2 \left(\frac{1}{2}\pi \sqrt{m^2 + n^2} \right)$$

An Aside: Madelung's Constant M

- The energy of a single ion in a large NaCl crystal

$$E = -\frac{z^2 e^2}{4\pi\epsilon_0 r_0} M$$

where

$$M = \sum'_{i,j,k} \frac{(-1)^{i+j+k}}{\sqrt{i^2 + j^2 + k^2}}$$

- Evaluate M by summing over spheres with increasing radius

$$M = -6 + 12/\sqrt{2} - 8/\sqrt{3} + 6/2 - 24/\sqrt{5} + \dots$$

Unfortunately, the resulting series *diverges*!

- Evaluate M by summing over cubes with increasing size converges

$$M = -1.74756\dots$$

- Several convergent series for M are known, the simplest being

$$M = -12\pi \sum_{m,n=1,3,5\dots} \operatorname{sech}^2 \left(\frac{1}{2}\pi \sqrt{m^2 + n^2} \right)$$

An Aside: Madelung's Constant M

- The energy of a single ion in a large NaCl crystal

$$E = -\frac{z^2 e^2}{4\pi\epsilon_0 r_0} M$$

where

$$M = \sum'_{i,j,k} \frac{(-1)^{i+j+k}}{\sqrt{i^2 + j^2 + k^2}}$$

- Evaluate M by summing over spheres with increasing radius

$$M = -6 + 12/\sqrt{2} - 8/\sqrt{3} + 6/2 - 24/\sqrt{5} + \dots$$

Unfortunately, the resulting series *diverges*!

- Evaluate M by summing over cubes with increasing size converges

$$M = -1.74756\dots$$

- Several convergent series for M are known, the simplest being

$$M = -12\pi \sum_{m,n=1,3,5\dots} \operatorname{sech}^2 \left(\frac{1}{2}\pi \sqrt{m^2 + n^2} \right)$$

An Aside: Madelung's Constant M

- The energy of a single ion in a large NaCl crystal

$$E = -\frac{z^2 e^2}{4\pi\epsilon_0 r_0} M$$

where

$$M = \sum'_{i,j,k} \frac{(-1)^{i+j+k}}{\sqrt{i^2 + j^2 + k^2}}$$

- Evaluate M by summing over spheres with increasing radius

$$M = -6 + 12/\sqrt{2} - 8/\sqrt{3} + 6/2 - 24/\sqrt{5} + \dots$$

Unfortunately, the resulting series *diverges*!

- Evaluate M by summing over cubes with increasing size converges

$$M = -1.74756\dots$$

- Several convergent series for M are known, the simplest being

$$M = -12\pi \sum_{m,n=1,3,5\dots} \operatorname{sech}^2 \left(\frac{1}{2} \pi \sqrt{m^2 + n^2} \right)$$

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = \frac{1}{2}$$

- ... but some cannot

$$\sum_{n=0}^{\infty} n = \infty$$

If a divergent series cannot be summed, physicists like to remove infinity

This is known as

- regularization, renormalisation, infrared cutoff, ultraviolet cutoff, ...

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = \frac{1}{2}$$

- ... but some cannot

$$\sum_{n=0}^{\infty} n = \infty$$

If a divergent series cannot be summed, physicists like to remove infinity

This is known as

- regularization, renormalisation, infrared cutoff, ultraviolet cutoff, ...

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = \frac{1}{2}$$

- ... but some cannot

$$\sum_{n=0}^{\infty} n = \infty$$

If a divergent series cannot be summed, physicists like to remove infinity

This is known as

- regularization, renormalisation, infrared cutoff, ultraviolet cutoff, ...

Summing Divergent Series

- Some divergent series can be summed in a sensible way ...

$$S = \sum_{n=0}^{\infty} (-1)^n = \frac{1}{2}$$

- ... but some cannot

$$\sum_{n=0}^{\infty} n = \infty$$

If a divergent series cannot be summed, physicists like to remove infinity

This is known as

- regularization, renormalisation, infrared cutoff, ultraviolet cutoff, ...

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation of $\sum_{n=0}^{\infty} f(n)$ (in particular, $f(n) = n$)

- Heat kernel regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) e^{-sn}$

in particular, $\sum_{n=0}^{\infty} n e^{-sn} = \frac{e^s}{(e^s - 1)^2} = \frac{1}{s^2} - \frac{1}{12} + O(s^2)$

Compare with $\int_0^{\infty} t e^{-st} dt = \frac{1}{s^2}$: divergent terms cancel

- Zeta function regularisation $\tilde{f}(s) = \sum_{n=0}^{\infty} f(n) n^{-s}$

in particular, $\sum_{n=0}^{\infty} n n^{-s} = \zeta(s-1)$, $\zeta(-1) = -\frac{1}{12}$

► Digression

Regularising Divergent Series

- Regularisation result should be independent of the method used
- In particular, for a reasonable class of *cutoff functions*

$$g(t; s) \quad \text{with} \quad \lim_{t \rightarrow \infty} g(t; s) = 0 \quad \text{and} \quad \lim_{s \rightarrow 0^+} g(t; s) = 1$$

replacing $f(t)$ by $f(t)g(t; s)$ should give the same result for $s \rightarrow 0^+$

- We need to study

$$\lim_{s \rightarrow 0^+} \Delta(fg) = \lim_{s \rightarrow 0^+} \left(\sum_{n=0}^{\infty} f(n)g(n; s) - \int_0^{\infty} f(t)g(t; s) dt \right)$$

Two mathematically sound approaches are

- Euler-Maclaurin Formula
- Abel-Plana Formula

Regularising Divergent Series

- Regularisation result should be independent of the method used
- In particular, for a reasonable class of *cutoff functions*

$$g(t; s) \quad \text{with} \quad \lim_{t \rightarrow \infty} g(t; s) = 0 \quad \text{and} \quad \lim_{s \rightarrow 0^+} g(t; s) = 1$$

replacing $f(t)$ by $f(t)g(t; s)$ should give the same result for $s \rightarrow 0^+$

- We need to study

$$\lim_{s \rightarrow 0^+} \Delta(fg) = \lim_{s \rightarrow 0^+} \left(\sum_{n=0}^{\infty} f(n)g(n; s) - \int_0^{\infty} f(t)g(t; s) dt \right)$$

Two mathematically sound approaches are

- Euler-Maclaurin Formula
- Abel-Plana Formula

Regularising Divergent Series

- Regularisation result should be independent of the method used
- In particular, for a reasonable class of *cutoff functions*

$$g(t; s) \quad \text{with} \quad \lim_{t \rightarrow \infty} g(t; s) = 0 \quad \text{and} \quad \lim_{s \rightarrow 0^+} g(t; s) = 1$$

replacing $f(t)$ by $f(t)g(t; s)$ should give the same result for $s \rightarrow 0^+$

- We need to study

$$\lim_{s \rightarrow 0^+} \Delta(fg) = \lim_{s \rightarrow 0^+} \left(\sum_{n=0}^{\infty} f(n)g(n; s) - \int_0^{\infty} f(t)g(t; s) dt \right)$$

Two mathematically sound approaches are

- Euler-Maclaurin Formula
- Abel-Plana Formula

Regularising Divergent Series

- Regularisation result should be independent of the method used
- In particular, for a reasonable class of *cutoff functions*

$$g(t; s) \quad \text{with} \quad \lim_{t \rightarrow \infty} g(t; s) = 0 \quad \text{and} \quad \lim_{s \rightarrow 0^+} g(t; s) = 1$$

replacing $f(t)$ by $f(t)g(t; s)$ should give the same result for $s \rightarrow 0^+$

- We need to study

$$\lim_{s \rightarrow 0^+} \Delta(fg) = \lim_{s \rightarrow 0^+} \left(\sum_{n=0}^{\infty} f(n)g(n; s) - \int_0^{\infty} f(t)g(t; s) dt \right)$$

Two mathematically sound approaches are

- Euler-Maclaurin Formula
- Abel-Plana Formula

The Euler-Maclaurin Formula



Leonhard Euler, 1707 - 1783



Colin Maclaurin, 1698 - 1746

The Euler-Maclaurin Formula

A formal derivation (Hardy, *Divergent Series*, 1949)

- Denoting $Df(x) = f'(x)$, the Taylor series can be written as

$$f(x+n) = e^{nD} f(x)$$

- It follows that

$$\begin{aligned} \sum_{n=0}^{N-1} f(x+n) &= \frac{e^{ND} - 1}{e^D - 1} f(x) = \frac{1}{e^D - 1} (f(x+N) - f(x)) \\ &= \left(D^{-1} + \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} \right) (f(x+N) - f(x)) \end{aligned}$$

$$\sum_{n=0}^{\infty} f(x+n) - \int_0^{\infty} f(x+t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} f(x)$$

The Euler-Maclaurin Formula

A formal derivation (Hardy, *Divergent Series*, 1949)

- Denoting $Df(x) = f'(x)$, the Taylor series can be written as

$$f(x+n) = e^{nD} f(x)$$

- It follows that

$$\begin{aligned} \sum_{n=0}^{N-1} f(x+n) &= \frac{e^{ND} - 1}{e^D - 1} f(x) = \frac{1}{e^D - 1} (f(x+N) - f(x)) \\ &= \left(D^{-1} + \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} \right) (f(x+N) - f(x)) \end{aligned}$$

$$\sum_{n=0}^{\infty} f(x+n) - \int_0^{\infty} f(x+t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} f(x)$$

The Euler-Maclaurin Formula

A formal derivation (Hardy, *Divergent Series*, 1949)

- Denoting $Df(x) = f'(x)$, the Taylor series can be written as

$$f(x+n) = e^{nD} f(x)$$

- It follows that

$$\begin{aligned} \sum_{n=0}^{N-1} f(x+n) &= \frac{e^{ND} - 1}{e^D - 1} f(x) = \frac{1}{e^D - 1} (f(x+N) - f(x)) \\ &= \left(D^{-1} + \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} \right) (f(x+N) - f(x)) \end{aligned}$$

► Skip precise statement

$$\sum_{n=0}^{\infty} f(x+n) - \int_0^{\infty} f(x+t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} f(x)$$

The Euler-Maclaurin Formula

A formal derivation (Hardy, *Divergent Series*, 1949)

- Denoting $Df(x) = f'(x)$, the Taylor series can be written as

$$f(x+n) = e^{nD} f(x)$$

- It follows that

$$\begin{aligned} \sum_{n=0}^{N-1} f(x+n) &= \frac{e^{ND} - 1}{e^D - 1} f(x) = \frac{1}{e^D - 1} (f(x+N) - f(x)) \\ &= \left(D^{-1} + \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} \right) (f(x+N) - f(x)) \end{aligned}$$

► Skip precise statement

$$\sum_{n=0}^{\infty} f(x+n) - \int_0^{\infty} f(x+t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} f(x)$$

The Euler-Maclaurin Formula

A formal derivation (Hardy, *Divergent Series*, 1949)

- Denoting $Df(x) = f'(x)$, the Taylor series can be written as

$$f(x+n) = e^{nD} f(x)$$

- It follows that

$$\begin{aligned} \sum_{n=0}^{N-1} f(x+n) &= \frac{e^{ND} - 1}{e^D - 1} f(x) = \frac{1}{e^D - 1} (f(x+N) - f(x)) \\ &= \left(D^{-1} + \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} \right) (f(x+N) - f(x)) \end{aligned}$$

► Skip precise statement

$$\sum_{n=0}^{\infty} f(x+n) - \int_0^{\infty} f(x+t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} D^{k-1} f(x)$$

Theorem (Euler-Maclaurin Formula)

If $f \in C^{2m}[0, N]$ then

$$\begin{aligned} \sum_{n=0}^N f(n) - \int_0^N f(t) dt &= \frac{1}{2} (f(0) + f(N)) + \\ &+ \sum_{k=1}^{m-1} \frac{B_{2k}}{(2k)!} \left(f^{(2k-1)}(N) - f^{(2k-1)}(0) \right) + R_m \end{aligned}$$

where

$$R_m = \int_0^N \frac{B_{2m} - B_{2m}(t - \lfloor t \rfloor)}{(2m)!} f^{(2m)}(t) dt$$

Here, $B_n(x)$ are Bernoulli polynomials and $B_n = B_n(0)$ are Bernoulli numbers

Applying the Euler-Maclaurin Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} f^{(k-1)}(0)$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ we find $RHS = -\frac{B_1}{1!} f(0) - \frac{B_2}{2!} f'(0)$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Euler-Maclaurin Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} f^{(k-1)}(0)$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ we find $RHS = -\frac{B_1}{1!} f(0) - \frac{B_2}{2!} f'(0)$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Euler-Maclaurin Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} f^{(k-1)}(0)$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ we find $RHS = -\frac{B_1}{1!}f(0) - \frac{B_2}{2!}f'(0)$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Euler-Maclaurin Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = - \sum_{k=1}^{\infty} \frac{B_k}{k!} f^{(k-1)}(0)$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ we find $RHS = -\frac{B_1}{1!}f(0) - \frac{B_2}{2!}f'(0)$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

The Abel-Plana Formula



Niels Henrik Abel, 1802 - 1829



Giovanni Antonio Amedeo Plana,
1781 - 1864

The Abel-Plana Formula

... a remarkable summation formula of Plana ...

Germund Dahlquist, 1997

The only two places I have ever seen this formula are in Hardy's book and in the writings of the "massive photon" people — who also got it from Hardy.

Jonathan P Dowling, 1989

The only other applications I am aware of, albeit for convergent series, are

- q -Gamma function asymptotics (Adri B Olde Daalhuis, 1994)
- uniform asymptotics for $\prod_{k=0}^{\infty} (1 - q^{n+k})^{-1}$ (myself, 1995)

The Abel-Plana Formula

... a remarkable summation formula of Plana ...

Germund Dahlquist, 1997

The only two places I have ever seen this formula are in Hardy's book and in the writings of the "massive photon" people — who also got it from Hardy.

Jonathan P Dowling, 1989

The only other applications I am aware of, albeit for convergent series, are

- q -Gamma function asymptotics (Adri B Olde Daalhuis, 1994)

- uniform asymptotics for $\prod_{k=0}^{\infty} (1 - q^{n+k})^{-1}$ (myself, 1995)

The Abel-Plana Formula

... a remarkable summation formula of Plana ...

Germund Dahlquist, 1997

The only two places I have ever seen this formula are in Hardy's book and in the writings of the "massive photon" people — who also got it from Hardy.

Jonathan P Dowling, 1989

The only other applications I am aware of, albeit for convergent series, are

- q -Gamma function asymptotics (Adri B Olde Daalhuis, 1994)

- uniform asymptotics for $\prod_{k=0}^{\infty} (1 - q^{n+k})^{-1}$ (myself, 1995)


The Abel-Plana Formula

- Use Cauchy's integral formula $f(\zeta) = \frac{1}{2\pi i} \oint_{\Gamma_\zeta} \frac{f(z)}{z - \zeta} dz$ together with

$$\pi \cot(\pi z) = \sum_{n=-\infty}^{\infty} \frac{1}{z - n}$$

to get

$$\sum_{n=0}^{\infty} f(n) = \frac{1}{2\pi i} \sum_{n=0}^{\infty} \oint_{\Gamma_n} \frac{f(z)}{z - n} dz = \frac{1}{2i} \int_{\Gamma} \cot(\pi z) f(z) dz$$



The Abel-Plana Formula

- Rotate the upper and lower arm of Γ by $\pm\pi/2$ to get

$$\sum_{n=0}^{\infty} f(n) = \frac{1}{2}f(0) + \frac{i}{2} \int_0^{\infty} (f(iy) - f(-iy)) \coth(\pi y) dy$$

- A similar trick gives

$$\int_0^\infty f(t) dt = \frac{i}{2} \int_0^\infty (f(iy) - f(-iy)) dy$$

The Abel-Plana Formula

- Rotate the upper and lower arm of Γ by $\pm\pi/2$ to get

$$\sum_{n=0}^{\infty} f(n) = \frac{1}{2}f(0) + \frac{i}{2} \int_0^{\infty} (f(iy) - f(-iy)) \coth(\pi y) dy$$

- A similar trick gives

$$\int_0^\infty f(t) dt = \frac{i}{2} \int_0^\infty (f(iy) - f(-iy)) dy$$

Taking the difference gives the elegant result

- ▶ Skip precise statement

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2}f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

Theorem (Abel-Plana Formula)

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ satisfy the following conditions

- (a) $f(z)$ is analytic for $\Re(z) \geq 0$ (though not necessarily at infinity)
- (b) $\lim_{|y| \rightarrow \infty} |f(x + iy)| e^{-2\pi|y|} = 0$ uniformly in x in every finite interval
- (c) $\int_{-\infty}^{\infty} |f(x + iy) - f(x - iy)| e^{-2\pi|y|} dy$ exists for every $x \geq 0$ and tends to zero for $x \rightarrow \infty$
- (d) $\int_0^{\infty} f(t) dt$ is convergent, and $\lim_{n \rightarrow \infty} f(n) = 0$

Then

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2} f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

Applying the Abel-Plana Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2}f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ one finds $RHS = -2 \int_0^{\infty} \frac{y dy}{e^{2\pi y} - 1}$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Abel-Plana Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2}f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series

- For $f(t) = t$ one finds $RHS = -2 \int_0^{\infty} \frac{y dy}{e^{2\pi y} - 1}$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Abel-Plana Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2}f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ one finds $RHS = -2 \int_0^{\infty} \frac{y dy}{e^{2\pi y} - 1}$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Applying the Abel-Plana Formula

$$\sum_{n=0}^{\infty} f(n) - \int_0^{\infty} f(t) dt = \frac{1}{2}f(0) + i \int_0^{\infty} \frac{f(iy) - f(-iy)}{e^{2\pi y} - 1} dy$$

- Introducing suitable cutoff functions $g(t; s)$ one can justify applying this to divergent series
- For $f(t) = t$ one finds $RHS = -2 \int_0^{\infty} \frac{y dy}{e^{2\pi y} - 1}$

$$\sum_{n=0}^{\infty} n - \int_0^{\infty} t dt = -\frac{1}{12}$$

Outline

- 1 The Casimir Effect
- 2 Making Sense of Infinity - Infinity
- 3 Conclusion

Conclusion

- Mathematical question posed in theoretical physics
- Some really nice, old formulæ from classical analysis
- The result has been verified in the laboratory
- The motivation for this talk:
 - Jonathan P Dowling "The Mathematics of the Casimir Effect"
Math Mag **62** (1989) 324

Doing mathematics and physics together can be more stimulating than doing either one separately, not to mention it's downright fun.

Conclusion

- Mathematical question posed in theoretical physics
- Some really nice, old formulæ from classical analysis
- The result has been verified in the laboratory
- The motivation for this talk:
 - Jonathan P Dowling "The Mathematics of the Casimir Effect"
Math Mag **62** (1989) 324

Doing mathematics and physics together can be more stimulating than doing either one separately, not to mention it's downright fun.

Conclusion

- Mathematical question posed in theoretical physics
- Some really nice, old formulæ from classical analysis
- The result has been verified in the laboratory
- The motivation for this talk:
 - Jonathan P Dowling "The Mathematics of the Casimir Effect"
Math Mag **62** (1989) 324

Doing mathematics and physics together can be more stimulating than doing either one separately, not to mention it's downright fun.

Conclusion

- Mathematical question posed in theoretical physics
- Some really nice, old formulæ from classical analysis
- The result has been verified in the laboratory
- The motivation for this talk:
 - Jonathan P Dowling “The Mathematics of the Casimir Effect”
Math Mag **62** (1989) 324

Doing mathematics and physics together can be more stimulating than doing either one separately, not to mention it's downright fun.

Conclusion

- Mathematical question posed in theoretical physics
- Some really nice, old formulæ from classical analysis
- The result has been verified in the laboratory
- The motivation for this talk:
 - Jonathan P Dowling “The Mathematics of the Casimir Effect”
Math Mag **62** (1989) 324

Doing mathematics and physics together can be more stimulating than doing either one separately, not to mention it's downright fun.

I had a feeling once about Mathematics - that I saw it all. Depth beyond depth was revealed to me - the Byss and Abyss. I saw - as one might see the transit of Venus or even the Lord Mayor's Show - a quantity passing through infinity and changing its sign from plus to minus. I saw exactly why it happened and why the tergiversation was inevitable but it was after dinner and I let it go.

Sir Winston Spencer Churchill, 1874 - 1965

The End

- Define for an increasing sequence $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ the zeta function

$$\zeta_\lambda(s) = \sum_{n=1}^{\infty} \lambda_n^{-s}$$

- If the zeta function has an analytic extension up to 0 then define the regularised infinite sum by

$$\sum_{n=1}^{\infty} \log \lambda_n = -\zeta'_\lambda(0)$$

- Alternatively, the regularised infinite product is given by

$$\prod_{n=1}^{\infty} \lambda_n = e^{-\zeta'_\lambda(0)}$$

- Define for an increasing sequence $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ the zeta function

$$\zeta_\lambda(s) = \sum_{n=1}^{\infty} \lambda_n^{-s}$$

- If the zeta function has an analytic extension up to 0 then define the regularised infinite sum by

$$\sum_{n=1}^{\infty} \log \lambda_n = -\zeta'_\lambda(0)$$

- Alternatively, the regularised infinite product is given by

$$\prod_{n=1}^{\infty} \lambda_n = e^{-\zeta'_\lambda(0)}$$

- Define for an increasing sequence $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ the zeta function

$$\zeta_\lambda(s) = \sum_{n=1}^{\infty} \lambda_n^{-s}$$

- If the zeta function has an analytic extension up to 0 then define the regularised infinite sum by

$$\sum_{n=1}^{\infty} \log \lambda_n = -\zeta'_\lambda(0)$$

- Alternatively, the regularised infinite product is given by

$$\prod_{n=1}^{\infty} \lambda_n = e^{-\zeta'_\lambda(0)}$$

- Let $\lambda_n = p_n$ be the n -th prime so that

$$\prod_p p = e^{-\zeta'_p(0)} \quad \text{where} \quad \zeta_p(s) = \sum_p p^{-s}$$

- A result from Landau and Walfisz (1920) gives

$$e^{\zeta_p(s)} = \prod_{n=1}^{\infty} \zeta(ns)^{\frac{\mu(n)}{n}}$$

- From this it takes a few steps to calculate

$$\zeta'_p(0) = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2 \log(2\pi)$$

- This can be made rigorous, so that

$$\prod_p p = 4\pi^2$$

- Corollary: there are infinitely many primes

- Let $\lambda_n = p_n$ be the n -th prime so that

$$\prod_p p = e^{-\zeta_p'(0)} \quad \text{where} \quad \zeta_p(s) = \sum_p p^{-s}$$

- A result from Landau and Walfisz (1920) gives

$$e^{\zeta_p(s)} = \prod_{n=1}^{\infty} \zeta(ns)^{\frac{\mu(n)}{n}}$$

- From this it takes a few steps to calculate

$$\zeta_p'(0) = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2 \log(2\pi)$$

- This can be made rigorous, so that

$$\prod_p p = 4\pi^2$$

- Corollary: there are infinitely many primes

- Let $\lambda_n = p_n$ be the n -th prime so that

$$\prod_p p = e^{-\zeta_p'(0)} \quad \text{where} \quad \zeta_p(s) = \sum_p p^{-s}$$

- A result from Landau and Walfisz (1920) gives

$$e^{\zeta_p(s)} = \prod_{n=1}^{\infty} \zeta(ns)^{\frac{\mu(n)}{n}}$$

- From this it takes a few steps to calculate

$$\zeta_p'(0) = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2 \log(2\pi)$$

- This can be made rigorous, so that

$$\prod_p p = 4\pi^2$$

- Corollary: there are infinitely many primes

- Let $\lambda_n = p_n$ be the n -th prime so that

$$\prod_p p = e^{-\zeta'_p(0)} \quad \text{where} \quad \zeta_p(s) = \sum_p p^{-s}$$

- A result from Landau and Walfisz (1920) gives

$$e^{\zeta_p(s)} = \prod_{n=1}^{\infty} \zeta(ns)^{\frac{\mu(n)}{n}}$$

- From this it takes a few steps to calculate

$$\zeta'_p(0) = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2 \log(2\pi)$$

- This can be made rigorous, so that

$$\prod_p p = 4\pi^2$$

- Corollary: there are infinitely many primes

- Let $\lambda_n = p_n$ be the n -th prime so that

$$\prod_p p = e^{-\zeta'_p(0)} \quad \text{where} \quad \zeta_p(s) = \sum_p p^{-s}$$

- A result from Landau and Walfisz (1920) gives

$$e^{\zeta_p(s)} = \prod_{n=1}^{\infty} \zeta(ns)^{\frac{\mu(n)}{n}}$$

- From this it takes a few steps to calculate

$$\zeta'_p(0) = \frac{1}{\zeta(0)} \frac{\zeta'(0)}{\zeta(0)} = -2 \log(2\pi)$$

- This can be made rigorous, so that

$$\prod_p p = 4\pi^2$$

- Corollary: there are infinitely many primes

▶ Back to main talk