

From parabolic fixed points to asymptotic enumeration

Thomas Prellberg

`thomas.prellberg@qmul.ac.uk`

School of Mathematical Sciences
Queen Mary, University of London

Combinatorial Enumeration

Problems in Combinatorial Enumeration

Examples of recursively definable structures:

- Number of partitions of a set into subsets (Bell, 1934)
 - Bell Numbers
- Number of partitions of a set into ordered subsets (Cayley, 1859)
 - ordered Bell Numbers
- Partition lattice chains (Lengyel, 1984)
 - Lengyel's Constant
- Analysis of a recursive Program (Knuth, 1991)
 - $t(x, y, z) = \text{if } x \leq y \text{ then } y \text{ else}$
 $t(t(x - 1, y, z), t(y - 1, z, x), t(z - 1, x, y))$

Bell Numbers

- B_n number of partitions of an n -set into subsets

$$B_3 = \binom{2}{0} B_2 + \binom{2}{1} B_1 + \binom{2}{2} B_0$$

$$\begin{array}{ccc} \{\{3\}, \{1, 2\}\} & \{\{3, 1\}, \{2\}\} & \{\{3, 1, 2\}\} \\ \{\{3\}, \{1\}, \{2\}\} & \{\{3, 2\}, \{1\}\} & \end{array}$$

Bell Numbers

- B_n number of partitions of an n -set into subsets

$$B_3 = \binom{2}{0} B_2 + \binom{2}{1} B_1 + \binom{2}{2} B_0$$

$$\begin{array}{ccc} \{\{3\}, \{1, 2\}\} & \{\{3, 1\}, \{2\}\} & \{\{3, 1, 2\}\} \\ \{\{3\}, \{1\}, \{2\}\} & \{\{3, 2\}, \{1\}\} & \end{array}$$

- Recurrence: $B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_{n-k}, \quad B_0 = 1$

Bell Numbers

- B_n number of partitions of an n -set into subsets

$$B_3 = \binom{2}{0} B_2 + \binom{2}{1} B_1 + \binom{2}{2} B_0$$

$$\begin{array}{ccc} \{\{3\}, \{1, 2\}\} & \{\{3, 1\}, \{2\}\} & \{\{3, 1, 2\}\} \\ \{\{3\}, \{1\}, \{2\}\} & \{\{3, 2\}, \{1\}\} & \end{array}$$

- Recurrence: $B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_{n-k}, \quad B_0 = 1$

- Exact expression for B_n from EGF: $b(z) = \sum_{n=0}^{\infty} B_n \frac{z^n}{n!}$

$$b(z) = \exp(e^z - 1) \Rightarrow B_n = \frac{1}{e} \sum_{m=0}^{\infty} \frac{m^n}{m!}$$

Bell Numbers (ctd.)

- Asymptotic growth (from saddle point method):

$$B_n \sim \exp \left(e^w (w^2 - w + 1) - \frac{1}{2} \log(w + 1) - 1 \right)$$

Scale: $w \exp(w) = n$ Lambert W -function:

$$w = W(n)$$

Bell Numbers (ctd.)

- Asymptotic growth (from saddle point method):

$$B_n \sim \exp \left(e^w (w^2 - w + 1) - \frac{1}{2} \log(w + 1) - 1 \right)$$

Scale: $w \exp(w) = n$ Lambert W -function:

$$w = W(n)$$

- Functional equation for OGF: $B(z) = \sum_{n=0}^{\infty} B_n z^n$

$$B(z) = \frac{z}{1-z} B \left(\frac{z}{1-z} \right) + 1$$

Ordered Bell Numbers

- S_n number of partitions of an n -set into ordered subsets
- Recurrence:
$$S_{n+1} = \sum_{k=0}^n \binom{n+1}{k+1} S_{n-k}, \quad S_0 = 1$$
- Exact expression for S_n from EGF: $s(z) = 1/(2 - e^z)$


$$S_n = \sum_{m=1}^n m! S_{n,m}, \quad S_{n,k} \text{ Stirling numbers 2nd kind}$$

- Asymptotic growth:
$$S_n \sim \frac{n!}{2} \frac{1}{(\log 2)^{n+1}}$$
- Functional equation for OGF (Klazar, private comm.):

$$B(z) = \frac{1}{2} \frac{1}{1-z} B\left(\frac{z}{1-z}\right) + \frac{1}{2}$$

Partition Lattice Chains

- Poset of partitions of an n -set



- Z_n number of chains from minimal to maximal element

$$Z_1 = 1, \quad Z_2 = 1, \quad , Z_3 = 4, \quad Z_4 = 32, \quad \dots$$

Partition Lattice Chains (ctd.)

- Recurrence (Lengyel):

$$Z_n = \sum_{k=1}^{n-1} S_{n,k} Z_k, \quad S_{n,k} \text{ Stirling numbers 2nd kind}$$

- Functional equation for EGF (Lengyel):

$$Z(z) = \frac{1}{2} Z(e^z - 1) + \frac{z}{2}$$

- Asymptotic growth (Babai, Lengyel):

$$Z_n \sim C_{\text{Lengyel}} (n!)^2 (2 \log 2)^{-n} n^{-1 - \frac{1}{3} \log 2}$$

- Lengyel's Constant (Flajolet, Salvy): $C_{\text{Lengyel}} = 1.0986858055 \dots$

Takeuchi Numbers

- Recursive function (Takeuchi):

$$t(x, y, z) = \text{if } x \leq y \text{ then } y \text{ else} \\ t(t(x - 1, y, z), t(y - 1, z, x), t(z - 1, x, y))$$

- $T(x, y, z)$ number of times the **else** clause is invoked when evaluating $t(x, y, z)$

- $T_n = T(n, 0, n + 1)$

$$T_1 = 1, \quad T_2 = 4, \quad , T_3 = 14, \quad T_4 = 53, \quad \dots$$

- Actual value of $t(x, y, z)$ is irrelevant

$$t(x, y, z) = \left\{ \begin{array}{ll} y & x \leq y \\ z & y \leq z \\ x & \text{else} \end{array} \right.$$

Takeuchi Numbers (ctd.)

- Recurrence (Knuth):

$$T_{n+1} = \sum_{k=0}^n \left[\binom{n+k}{n} - \binom{n+k}{n+1} \right] T_{n-k} + \sum_{k=1}^{n+1} \binom{2k}{k} \frac{1}{k+1}$$

- Functional equation for OGF (Knuth):

$$T(z) = zC(z)T(zC(z)) + \frac{C(z) - 1}{1 - z}, \quad C(z) = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{z^k}{k+1}$$

- Asymptotic growth (Prellberg):

$$T_n \sim C_{\text{Takeuchi}} B_n \exp \frac{1}{2} W(n)^2, \quad C_{\text{Takeuchi}} = 2.2394331040 \dots$$

General Setting

Common Features

- Linear recurrences:

$$X_n = \sum_{k=1}^n c_{n,k} X_{n-k} + b_n$$

Common Features

- Linear recurrences:

$$X_n = \sum_{k=1}^n c_{n,k} X_{n-k} + b_n$$

- Functional equations for OGF/EGF:

$$X(z) = a(z)X \circ f(z) + b(z)$$

Common Features

- Linear recurrences:

$$X_n = \sum_{k=1}^n c_{n,k} X_{n-k} + b_n$$

- Functional equations for OGF/EGF:

$$X(z) = a(z)X \circ f(z) + b(z)$$

- Parabolic fixed point:

$$f(z) = z + cz^2 + dz^3 + \dots$$

Common Features

- Linear recurrences:

$$X_n = \sum_{k=1}^n c_{n,k} X_{n-k} + b_n$$

- Functional equations for OGF/EGF:

$$X(z) = a(z)X \circ f(z) + b(z)$$

- Parabolic fixed point:

$$f(z) = z + cz^2 + dz^3 + \dots$$

- Caveat: divergence of GF!

Generalization: Recursive Structures

- View combinatorial structures as formed of “atoms”
- Substitution operation $\mathcal{A} = \mathcal{B} \circ \mathcal{C}$:
“substitute elements of \mathcal{C} for atoms of \mathcal{B} ”

$$\mathcal{B} \circ \mathcal{C} = \sum_{k \geq 0} \mathcal{B}_k \times \overbrace{\mathcal{C} \times \mathcal{C} \times \dots \times \mathcal{C}}^k$$

- The associated OGF satisfies $A(z) = B(C(z))$
- A recursively definable structure \mathcal{X} is defined by

$$\mathcal{X} = \mathcal{A} \times \mathcal{X} \circ \mathcal{F} + \mathcal{B}$$

- The associated OGF satisfies $X(z) = A(z)X \circ F(z) + B(z)$

Ingredients for General Theory

- Formal solution of the functional equation
(leads to divergent FPS)
- Cauchy formula
- Analytic iteration theory near parabolic
fixed points (Milnor, Beardon)
- Saddle point analysis

Formal Power Series Solution

Let the FPS $X(z)$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $a(z)$, $f(z)$, and $b(z)$ analytic near $z = 0$ and

$$f(z) = z + cz^2 + dz^3 + \dots, \quad c > 0$$

Formal Power Series Solution

Let the FPS $X(z)$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $a(z)$, $f(z)$, and $b(z)$ analytic near $z = 0$ and

$$f(z) = z + cz^2 + dz^3 + \dots, \quad c > 0$$

Then

$$X(z) = \sum_{m=0}^{\infty} \left(\prod_{k=0}^{m-1} a \circ f^k(z) \right) b \circ f^m(z)$$

as FPS (this is a divergent power series!)

Inversion via Cauchy Formula

From

$$X(z) = \sum_{m=0}^{\infty} \left(\prod_{k=0}^{m-1} a \circ f^k(z) \right) b \circ f^m(z)$$

we compute

$$X_n = [z^n]X(z) = \sum_{m=0}^{\infty} X_{n,m}$$

with

$$X_{n,m} = \frac{1}{2\pi i} \oint \left(\prod_{k=0}^{m-1} a \circ f^k(z) \right) b \circ f^m(z) \frac{dz}{z^{n+1}}$$

Simplification via Homogeneous Eqn.

- Let $Y(z)$ be a solution of the *homogeneous* equation

$$Y(z) = a(z) Y \circ f(z)$$

Then $X_{n,m}$ simplifies to

$$X_{n,m} = \frac{1}{2\pi i} \oint \frac{b \circ f^m(z)}{Y \circ f^m(z)} Y(z) \frac{dz}{z^{n+1}}$$

Simplification via Homogeneous Eqn.

- Let $Y(z)$ be a solution of the *homogeneous* equation

$$Y(z) = a(z) Y \circ f(z)$$

Then $X_{n,m}$ simplifies to

$$X_{n,m} = \frac{1}{2\pi i} \oint \frac{b \circ f^m(z)}{Y \circ f^m(z)} Y(z) \frac{dz}{z^{n+1}}$$

- Needed: existence of $Y(z)$ and analyticity properties
 - Analytic iteration theory (Milnor, Beardon)

Analytic Iteration Theory


Analytic Iteration Theory

“Parabolic Linearization Theorem” \Rightarrow conjugacy of $f(z)$ to a shift

Analytic Iteration Theory

“Parabolic Linearization Theorem” \Rightarrow conjugacy of $f(z)$ to a shift


• $f^{-1}(z)$ exists in cardioid domain and maps contractively into it



Analytic Iteration Theory

“Parabolic Linearization Theorem” \Rightarrow conjugacy of $f(z)$ to a shift

- $f^{-1}(z)$ exists in cardioid domain and maps contractively into it




- $f^{-1}(z) = f^{-1} \circ \mu(s) = \mu(s + 1)$ for $s \in \mathcal{D}(\mu)$

Analytic Iteration Theory

“Parabolic Linearization Theorem” \Rightarrow conjugacy of $f(z)$ to a shift

- $f^{-1}(z)$ exists in cardioid domain and maps contractively into it



- $f^{-1}(z) = f^{-1} \circ \mu(s) = \mu(s + 1)$ for $s \in \mathcal{D}(\mu)$
- $f^k(z) = \mu(\mu^{-1}(z) - k)$ for z sufficiently small

Analytic Iteration Theory (ctd.)

- $\mu(s)$ admits a complete asymptotic expansion for $\Re(s) \rightarrow \infty$:

$$\mu(s) \sim \frac{1}{cs} \left(1 + \left(1 - \frac{d}{c^2} \right) \frac{\log s}{s} + \sum_{k=2}^{\infty} \sum_{j=0}^k \mu_{j,k} \frac{(\log s)^j}{s^k} \right)$$

- $f^{-m} \circ \mu(s) = \mu(s+m)$ admits a complete asymptotic expansion for $m \rightarrow \infty$:

$$\mu(s+m) \sim \frac{1}{cm} \left(1 + \left(1 - \frac{d}{c^2} - s \right) \frac{\log m}{m} + \sum_{k=2}^{\infty} \sum_{j=0}^k \nu_{j,k}(s) \frac{(\log m)^j}{m^k} \right)$$

Solution of the Homogeneous Eqn.

- Substitute $z = \mu(s)$:

$$Y(z) = a(z)Y \circ f(z) \quad \implies \quad Y \circ \mu(s) = a \circ \mu(s)Y \circ \mu(s-1)$$

- Solution is given by

$$Y \circ \mu(s) = \lim_{n \rightarrow \infty} \frac{a \circ \mu(1)a \circ \mu(2) \dots a \circ \mu(n) (a \circ \mu(n))^s}{a \circ \mu(s+1)a \circ \mu(s+2) \dots a \circ \mu(s+n)}$$

which defines an analytic function in $\mathcal{D}(\mu)$

- Asymptotics as $n \rightarrow \infty$:

$$\frac{Y \circ \mu(s+n)}{Y \circ \mu(n)} \sim (a \circ \mu(n))^s$$

The Saddle Point Condition

$$X_n = \sum_{m=0}^{\infty} X_{n,m}, \quad X_{n,m} = \frac{1}{2\pi i} \oint \frac{b \circ f^m(z)}{Y \circ f^m(z)} Y(z) \frac{dz}{z^{n+1}}$$

$$\sim \dots \int_C \frac{b \circ \mu(s)}{Y \circ \mu(s)} \left(a \circ \mu(m) e^{\frac{n}{m}} \right)^s ds$$

- Saddle point at $a \circ \mu(m) e^{\frac{n}{m}} = 1$
- $a(z) = a_k z^k + \dots, \mu(s) \sim (cs)^{-1} \Rightarrow a \circ \mu(m) \sim a_k (cm)^{-k}$
- Different behavior according to

● $k = 0$:

$$m = -\frac{n}{\log a_0} \quad 0 < a_0 < 1$$

● $k \geq 1$:

$$m = \frac{n}{kW(cn/ka_k^{1/k})} \quad a_k > 0$$

More tedious calculations...

Main Results

RESULT 1: Let the FPS $X(z) = \sum_{n=0}^{\infty} X_n z^n$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $f(z) = z + cz^2 + dz^3 + \dots$, $a(z) = a_0 + a_1 z + \dots$, and $b(z)$ analytic near zero.

Main Results

RESULT 1: Let the FPS $X(z) = \sum_{n=0}^{\infty} X_n z^n$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $f(z) = z + cz^2 + dz^3 + \dots$, $a(z) = a_0 + a_1 z + \dots$, and $b(z)$ analytic near zero.

If $c > 0$ and $0 < a_0 < 1$ then

$$X_n \sim Dn! (-c/\log a_0)^n n^{(1-\frac{d}{c^2})\log a_0 - 1 + \frac{a_1}{a_0}}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2\pi i} \int_c \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds (-\log a_0)^{-(1-\frac{d}{c^2})\log a_0}$$

Main Results (ctd.)

RESULT 2: Let the FPS $X(z) = \sum_{n=0}^{\infty} X_n z^n$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $f(z) = z + cz^2 + dz^3 + \dots$, $a(z) = a_1 z + \dots$, and $b(z)$ analytic near zero.

Main Results (ctd.)

RESULT 2: Let the FPS $X(z) = \sum_{n=0}^{\infty} X_n z^n$ satisfy

$$X(z) = a(z)X \circ f(z) + b(z)$$

with $f(z) = z + cz^2 + dz^3 + \dots$, $a(z) = a_1 z + \dots$, and $b(z)$ analytic near zero.

If $c > 0$ and $a_1 > 0$ then

$$X_n \sim D c^n e^{-\frac{1}{2}(1-\frac{d}{c^2})W(\frac{c}{a_1}n)^2} \sum_{m=0}^{\infty} \frac{m^n}{m!} \left(\frac{a_1}{c}\right)^m$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds e^{\frac{1}{2}(1-\frac{d}{c^2})(\log \frac{a_1}{c})^2}$$

Applications

Application: Bell Numbers

- Functional equation for OGF:

$$B(z) = \frac{z}{1-z} B\left(\frac{z}{1-z}\right) + 1$$

- $a(z) = \frac{z}{1-z}, f(z) = \frac{z}{1-z}, b(z) = 1$
- $\mu(s) = 1/s, Y \circ \mu(s) = 1/\Gamma(s)$

Application: Bell Numbers

- Functional equation for OGF:

$$B(z) = \frac{z}{1-z} B\left(\frac{z}{1-z}\right) + 1$$

- $a(z) = \frac{z}{1-z}, f(z) = \frac{z}{1-z}, b(z) = 1$

- $\mu(s) = 1/s, Y \circ \mu(s) = 1/\Gamma(s)$

insert $c = 1, d = 1, a_1 = 1$ into

$$B_n \sim D c^n e^{-\frac{1}{2}(1-\frac{d}{c^2})W(\frac{c}{a_1}n)^2} \sum_{m=0}^{\infty} \frac{m^n}{m!} \left(\frac{a_1}{c}\right)^m$$

with

$$D = \frac{1}{2\pi i} \int_c \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds e^{\frac{1}{2}(1-\frac{d}{c^2})(\log \frac{a_1}{c})^2}$$

Application: Bell Numbers

- Functional equation for OGF:

$$B(z) = \frac{z}{1-z} B\left(\frac{z}{1-z}\right) + 1$$

- $a(z) = \frac{z}{1-z}, f(z) = \frac{z}{1-z}, b(z) = 1$

- $\mu(s) = 1/s, Y \circ \mu(s) = 1/\Gamma(s)$

- Asymptotics:

$$B_n \sim D \sum_{m=0}^{\infty} \frac{m^n}{m!}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2\pi i} \int_C \Gamma(s) ds = \frac{1}{e} \quad (\text{sum of residues})$$

Application: Ordered Bell Numbers

- Functional equation for OGF:

$$S(z) = \frac{1}{2} \frac{1}{1-z} S\left(\frac{z}{1-z}\right) + \frac{1}{2}$$

- $a(z) = \frac{1}{2} \frac{1}{1-z}, f(z) = \frac{z}{1-z}, b(z) = \frac{1}{2}$
- $\mu(s) = 1/s, Y \circ \mu(s) = 1/\Gamma(s)$

Application: Ordered Bell Numbers

- Functional equation for OGF:

$$S(z) = \frac{1}{2} \frac{1}{1-z} S\left(\frac{z}{1-z}\right) + \frac{1}{2}$$

- $a(z) = \frac{1}{2} \frac{1}{1-z}, f(z) = \frac{z}{1-z}, b(z) = \frac{1}{2}$
- $\mu(s) = 1/s, Y \circ \mu(s) = 1/\Gamma(s)$

insert $c = 1, d = 1, a_0 = \frac{1}{2}, a_1 = \frac{1}{2}$ into

$$S_n \sim Dn! (-c/\log a_0)^n n^{(1-\frac{d}{c^2})\log a_0 - 1 + \frac{a_1}{a_0}}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2\pi i} \int_C \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds (-\log a_0)^{-(1-\frac{d}{c^2})\log a_0}$$

Application: Ordered Bell Numbers

- Functional equation for OGF:

$$S(z) = \frac{1}{2} \frac{1}{1-z} S\left(\frac{z}{1-z}\right) + \frac{1}{2}$$

- $a(z) = \frac{1}{2} \frac{1}{1-z}$, $f(z) = \frac{z}{1-z}$, $b(z) = \frac{1}{2}$
- $\mu(s) = 1/s$, $Y \circ \mu(s) = 1/\Gamma(s)$

- Asymptotics:

$$S_n \sim D n! \frac{1}{(\log 2)^n}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2 \log 2}$$

Application: Partition Lattice Chains

- Functional equation for EGF:

$$Z(z) = \frac{1}{2}Z(e^z - 1) + \frac{z}{2}$$

- $a(z) = \frac{1}{2}, f(z) = e^z - 1, b(z) = \frac{z}{2}$

- $\mu(s) \sim \frac{2}{s}(1 - \frac{\log s}{3s} + \dots), Y \circ \mu(s) = 2^s$

Application: Partition Lattice Chains

- Functional equation for EGF:

$$Z(z) = \frac{1}{2}Z(e^z - 1) + \frac{z}{2}$$

- $a(z) = \frac{1}{2}, f(z) = e^z - 1, b(z) = \frac{z}{2}$

- $\mu(s) \sim \frac{2}{s}(1 - \frac{\log s}{3s} + \dots), Y \circ \mu(s) = 2^s$

insert $c = \frac{1}{2}, d = \frac{1}{6}, a_0 = \frac{1}{2}, a_1 = 0$ into

$$\frac{Z_n}{n!} \sim Dn! (-c/\log a_0)^n n^{(1 - \frac{d}{c^2}) \log a_0 - 1 + \frac{a_1}{a_0}}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2\pi i} \int_C \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds (-\log a_0)^{-(1 - \frac{d}{c^2}) \log a_0}$$

Application: Partition Lattice Chains

- Functional equation for EGF:

$$Z(z) = \frac{1}{2}Z(e^z - 1) + \frac{z}{2}$$

- $a(z) = \frac{1}{2}$, $f(z) = e^z - 1$, $b(z) = \frac{z}{2}$

- $\mu(s) \sim \frac{2}{s}(1 - \frac{\log s}{3s} + \dots)$, $Y \circ \mu(s) = 2^s$

- Asymptotics:

$$Z_n \sim D(n!)^2 (2 \log 2)^{-n} n^{-1 - \frac{1}{3} \log 2}$$

as $n \rightarrow \infty$, where

$$D = \frac{1}{2} (\log 2)^{\frac{1}{3} \log 2} \frac{1}{2\pi i} \int_C 2^s \mu(s) ds = 1.0986858055 \dots$$

Application: Takeuchi Numbers

- Functional equation for OGF:

$$T(z) = zC(z)T(zC(z)) + \frac{C(z) - 1}{1 - z}, \quad C(z) = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{z^k}{k+1}$$

- $a(z) = zC(z), f(z) = zC(z), b(z) = \frac{C(z)-1}{1-z}$

- $\mu(s) \sim \frac{1}{s}(1 - \frac{\log s}{s} + \dots), Y \circ \mu(s) \sim e^{-\frac{1}{2}(\log s)^2} / \Gamma(s)$

Application: Takeuchi Numbers

- Functional equation for OGF:

$$T(z) = zC(z)T(zC(z)) + \frac{C(z) - 1}{1 - z}, \quad C(z) = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{z^k}{k+1}$$

- $a(z) = zC(z), f(z) = zC(z), b(z) = \frac{C(z)-1}{1-z}$

- $\mu(s) \sim \frac{1}{s}(1 - \frac{\log s}{s} + \dots), Y \circ \mu(s) \sim e^{-\frac{1}{2}(\log s)^2} / \Gamma(s)$

insert $c = 1, d = 2, a_1 = 1$ into

$$T_n \sim Dc^n e^{-\frac{1}{2}(1-\frac{d}{c^2})W(\frac{c}{a_1}n)^2} \sum_{m=0}^{\infty} \frac{m^n}{m!} \left(\frac{a_1}{c}\right)^m$$

with

$$D = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds e^{\frac{1}{2}(1-\frac{d}{c^2})(\log \frac{a_1}{c})^2}$$

Application: Takeuchi Numbers

- Functional equation for OGF:

$$T(z) = zC(z)T(zC(z)) + \frac{C(z) - 1}{1 - z}, \quad C(z) = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{z^k}{k+1}$$

- $a(z) = zC(z), f(z) = zC(z), b(z) = \frac{C(z)-1}{1-z}$

- $\mu(s) \sim \frac{1}{s}(1 - \frac{\log s}{s} + \dots), Y \circ \mu(s) \sim e^{-\frac{1}{2}(\log s)^2} / \Gamma(s)$

- Asymptotics:

$$T_n \sim D \sum_{m=0}^{\infty} \frac{m^n}{m!} e^{\frac{1}{2}W(n)^2} = D' B_n e^{\frac{1}{2}W(n)^2}$$

as $n \rightarrow \infty$, where

$$D' = \frac{e}{2\pi i} \int_{\mathcal{C}} \frac{b \circ \mu(s)}{Y \circ \mu(s)} ds = 2.2394331040 \dots$$

Summary

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration
- A large class of linear recurrences corresponding to recursively definable structures can be treated asymptotically

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration
- A large class of linear recurrences corresponding to recursively definable structures can be treated asymptotically
- Further applications?

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration
- A large class of linear recurrences corresponding to recursively definable structures can be treated asymptotically
- Further applications?

Other results:

- Numerical evaluation of the constants to about 50 decimal places

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration
- A large class of linear recurrences corresponding to recursively definable structures can be treated asymptotically
- Further applications?

Other results:

- Numerical evaluation of the constants to about 50 decimal places
- Computation of the next terms in the asymptotic expansion (by a different, non-rigorous method)

Summary

- Interesting application of analytic iteration theory and classical complex analysis in the services of asymptotic enumeration
- A large class of linear recurrences corresponding to recursively definable structures can be treated asymptotically
- Further applications?

Other results:

- Numerical evaluation of the constants to about 50 decimal places
- Computation of the next terms in the asymptotic expansion (by a different, non-rigorous method)

To be done:

- Computation of the contour integrals determining the constants

The End