

# Computing scaling functions for two-dimensional vesicle models

from generating functions to coalescing saddle points

Thomas Prellberg

School of Mathematical Sciences  
Queen Mary, University of London

November 8, 2010

# Topic Outline


- 1 Motivation
  - Vesicle Generating Function
  - Singularity Diagram
  - Scaling Function
- 2 From Lattice Walks to Basic Hypergeometric Series
  - $q$ -Deformed Algebraic Equations
  - $q$ -Difference Equations
  - Basic Hypergeometric Series
- 3 Asymptotic Analysis
  - Contour Integral Representation
  - Saddle Point Analysis
  - Generalisation
- 4 Outlook

# Outline

- 1 Motivation
  - Vesicle Generating Function
  - Singularity Diagram
  - Scaling Function
- 2 From Lattice Walks to Basic Hypergeometric Series
  - $q$ -Deformed Algebraic Equations
  - $q$ -Difference Equations
  - Basic Hypergeometric Series
- 3 Asymptotic Analysis
  - Contour Integral Representation
  - Saddle Point Analysis
  - Generalisation
- 4 Outlook

# Vesicle Generating Function


- 3-dim vesicle (bubble) with surface and volume
- 2-dim lattice model: polygons on the square lattice



$c_{m,n}$  number of polygons with area  $m$  and perimeter  $2n$

# Vesicle Generating Function

- 3-dim vesicle (bubble) with surface and volume
- 2-dim lattice model: polygons on the square lattice



$c_{m,n}$  number of polygons with area  $m$  and perimeter  $2n$


$$G(x, q) = \sum_{n,m} c_{m,n} x^n q^m \quad \text{generating function}$$

Wanted:

- an explicit formula for  $G(x, q)$
- singularity structure, e.g.  $q_c(x)$


# Singularity Diagram

Folklore: universal behaviour near a “critical point”



# Singularity Diagram

Folklore: universal behaviour near a “critical point”



- scaling function  $f$  with crossover exponent  $\phi$ :

$$G^{\text{sing}}(x, q) \sim (1 - q)^{-\gamma_t} f([1 - q]^{-\phi} [x_t - x])$$

as  $q \rightarrow 1$  and  $x \rightarrow x_t$  with  $z = [1 - q]^{-\phi} [x_t - x]$  fixed

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$



# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)
- Staircase polygons (skew-Ferrer diagrams)
  - Rigorous derivation (Prellberg)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)
- Staircase polygons (skew-Ferrer diagrams)
  - Rigorous derivation (Prellberg)
- Brownian excursion (BE) in half-plane
  - Probabilistic analysis (Louchard)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)
- Staircase polygons (skew-Ferrer diagrams)
  - Rigorous derivation (Prellberg)
- Brownian excursion (BE) in half-plane
  - Probabilistic analysis (Louchard)
  - Langevin equation (Kearney and Majumdar)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)
- Staircase polygons (skew-Ferrer diagrams)
  - Rigorous derivation (Prellberg)
- Brownian excursion (BE) in half-plane
  - Probabilistic analysis (Louchard)
  - Langevin equation (Kearney and Majumdar)
- Area statistics of outer boundary of random loops
  - Monte-Carlo simulation (Richard)

# Scaling Function

*Surprisingly often*  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$


- Square lattice vesicle generating function
  - Enumeration of  $c_{m,n}$ , numerical analysis of moments (Richard, Guttmann, Jensen)
- Directed models with  $q$ -algebraic equations
  - Heuristic scaling Ansatz (Richard)
- Staircase polygons (skew-Ferrer diagrams)
  - Rigorous derivation (Prellberg)
- Brownian excursion (BE) in half-plane
  - Probabilistic analysis (Louchard)
  - Langevin equation (Kearney and Majumdar)
- Area statistics of outer boundary of random loops
  - Monte-Carlo simulation (Richard)
- $q$ -Analogue of the Painlevé II equation (Witte)

# Outline

- 1 Motivation
  - Vesicle Generating Function
  - Singularity Diagram
  - Scaling Function
- 2 From Lattice Walks to Basic Hypergeometric Series
  - $q$ -Deformed Algebraic Equations
  - $q$ -Difference Equations
  - Basic Hypergeometric Series
- 3 Asymptotic Analysis
  - Contour Integral Representation
  - Saddle Point Analysis
  - Generalisation
- 4 Outlook




## Example 1: Dyck Paths



$2n = 14$  steps enclosing an area of size  $m = 9$

# Example 1: Dyck Paths




$2n = 14$  steps enclosing an area of size  $m = 9$

$$G(t, q) = \sum_{m, n} c_{m, n} t^n q^m$$

$t$  counts pairs of up/down steps,  $q$  counts enclosed area

# Example 1: Dyck Paths


- A functional equation



$$G(t, q) = 1 + tG(qt, q)G(t, q)$$

# Example 1: Dyck Paths

- A functional equation




$$G(t, q) = 1 + tG(qt, q)G(t, q)$$

- $C(t) = G(t, 1)$  satisfies  $C(t) = 1 + tC(t)^2$

$$C(t) = \frac{1 - \sqrt{1 - 4t}}{2t} = \sum_{n=0}^{\infty} \frac{t^n}{n+1} \binom{2n}{n}$$


Generating function of Catalan numbers

## Example 2: A Pair of Directed Walks



Two directed walks not allowed to cross

## Example 2: A Pair of Directed Walks




Two directed walks not allowed to cross

$$G(x, y, q) = \sum_{m, n_x, n_y} c_{m, n_x, n_y} x^{n_x} y^{n_y} q^m$$

$x$  and  $y$  count pairs of east and north steps,  $q$  counts enclosed area

## Example 2: A Pair of Directed Walks


- A functional equation



$$G(x, y, q) = 1 + yG(qx, y, q) + xG(x, y, q) + yG(qx, y, q)xG(x, y, q)$$

## Example 2: A Pair of Directed Walks

- A functional equation




$$G(x, y, q) = 1 + yG(qx, y, q) + xG(x, y, q) + yG(qx, y, q)xG(x, y, q)$$

- $G(t, t, 1) = 1 + tC(t)$  Catalan generating function



## Example 3: Partially Directed Walks Above $y = x$


- A functional equation



$$G(x, y, q) = 1 + yG(qx, y, q)xG(x, y, q) + y(G(qx, y, q) - 1)y$$

## Example 3: Partially Directed Walks Above $y = x$

- A functional equation




$$G(x, y, q) = 1 + yG(qx, y, q)xG(x, y, q) + y(G(qx, y, q) - 1)y$$

- $G(x, y, 1) = C\left(\frac{xy}{1-y^2}\right)$  Catalan generating function


## Example 4: Directed Column-Convex Polygons

- Lattice polygon
  - partially directed upper perimeter
  - fully directed lower perimeter



## Example 4: Directed Column-Convex Polygons


- Lattice polygon
  - partially directed upper perimeter
  - fully directed lower perimeter
- A functional equation



$$\begin{aligned}
 0 &= G(q^2x)G(qx)G(x) \\
 &+ yG(q^2x)G(qx) + yG(q^2x)G(x) - (1+q)G(qx)G(x) \\
 &+ y^2G(q^2x) - y(1+q)G(qx) + q(1+qx(y-1))G(x) \\
 &+ yq^2x(y-1)
 \end{aligned}$$

# Example 4: Directed Column-Convex Polygons

- Lattice polygon
  - partially directed upper perimeter
  - fully directed lower perimeter



- A functional equation

$$\begin{aligned}
 0 &= G(q^2x)G(qx)G(x) \\
 &+ yG(q^2x)G(qx) + yG(q^2x)G(x) - (1+q)G(qx)G(x) \\
 &+ y^2G(q^2x) - y(1+q)G(qx) + q(1+qx(y-1))G(x) \\
 &+ yq^2x(y-1)
 \end{aligned}$$

- $q = 1$  gives a cubic equation for  $G(x, y, 1)$

# Summary of the Examples

Different  $q$ -deformations of Catalan-type generating functions:

- Dyck paths

$$G(t) = 1 + tG(t)G(qt)$$

- Pair of directed walks

$$G(x) = (1 + xG(x))(1 + yG(qx))$$

- Partially directed walks above the diagonal

$$G(x) = 1 + xyG(x)G(qx) + y^2(G(qx) - 1)$$

and also of higher-order algebraic generating functions:

- e.g. directed column-convex polygons

# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

An aside:

- $G(t)$  admits a nice continued fraction expansion

$$G(t) = \frac{1}{1 - \frac{t}{1 - \frac{qt}{1 - \frac{q^2 t}{1 - \dots}}}}$$

- Connections with orthogonal polynomials, combinatorics of weighted lattice paths, ...

# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

An aside:

- $G(t)$  admits a nice continued fraction expansion

$$G(t) = \frac{1}{1 - \frac{t}{1 - \frac{qt}{1 - \frac{q^2 t}{1 - \dots}}}}$$

- Connections with orthogonal polynomials, combinatorics of weighted lattice paths, ...
- However, useless for finer asymptotic analysis of  $q \rightarrow 1$ .



# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

Better:

- Linearise the functional equation using

$$G(t) = \frac{H(qt)}{H(t)}$$

# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

Better:

- Linearise the functional equation using

$$G(t) = \frac{H(qt)}{H(t)}$$

- Obtain a linear  $q$ -difference equation

$$H(qt) = H(t) + tH(q^2t)$$

# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

Better:

- Linearise the functional equation using

$$G(t) = \frac{H(qt)}{H(t)}$$

- Obtain a linear  $q$ -difference equation

$$H(qt) = H(t) + tH(q^2t)$$

- Explicit solution

$$H(t) = \sum_{n=0}^{\infty} \frac{q^{n^2-n}(-t)^n}{(q; q)_n} = {}_0\phi_1(-; 0; q, -t)$$

# Example 1: Solving $G(t) = 1 + tG(t)G(qt)$

Better:

- Linearise the functional equation using

$$G(t) = \frac{H(qt)}{H(t)}$$

- Obtain a linear  $q$ -difference equation

$$H(qt) = H(t) + tH(q^2t)$$

- Explicit solution

$$H(t) = \sum_{n=0}^{\infty} \frac{q^{n^2-n}(-t)^n}{(q; q)_n} = {}_0\phi_1(-; 0; q, -t)$$

${}_0\phi_1(-; 0; q, -qt)$  a  $q$ -Airy function (Ismail)

## Example 2: Solving $G(x) = (1 + xG(x))(1 + yG(qx))$

Again:

- Linearise the functional equation using

$$G(x) = \frac{1}{x} \left( \frac{H(qx)}{H(x)} - 1 \right)$$

- Obtain a linear  $q$ -difference equation

$$q(H(qx) - H(x)) = qxH(qx) + y(H(q^2x) - H(qx))$$

- Explicit solution

$$H(t) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-x)^n}{(y; q)_n (q; q)_n} = {}_1\phi_1(0; y; q, x)$$

# Example 3: $G(x) = 1 + xyG(x)G(qx) + y^2(G(qx) - 1)$

One more time:

- Linearise the functional equation using

$$G(x) = \frac{y}{x} \left( \frac{H(qx)}{H(x)} - 1 \right)$$

- Obtain a linear  $q$ -difference equation

$$q(H(qx) - H(x)) = qx(1/y - y)H(qx) + y^2(H(q^2x) - H(qx))$$

- Explicit solution

$$H(t) = \sum_{n=0}^{\infty} \frac{(-x(1-y^2)/y)^n}{(y^2; q)_n (q; q)_n} = {}_2\phi_1(0, 0; y^2; q, -x(1-y^2)/y)$$

## Example 4: Directed Column-Convex Polygons

Surprisingly, this trick works also here:

- Linearise the functional equation using

$$G(x) = y \left( \frac{H(qx)}{H(x)} - 1 \right)$$

- Obtain a linear  $q$ -difference equation

$$y^2 H(q^3 x) - y(q + y + 1) H(q^2 x) + (y + q + qy + q^2 x(y - 1)) H(qx) - qH(x) = 0$$

- Explicit solution

$$H(t) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (-qx(1-y))^n}{(y; q)_n (qy; q)_n (q; q)_n} = {}_2\phi_2(0, 0; y, qy; q, qx(1-y))$$

# Summary:

Different  $q$ -deformations of Catalan-type generating functions:

- Dyck paths

$$G(t, q) = \frac{{}_0\phi_1(-; 0; q, -qt)}{{}_0\phi_1(-; 0; q, -t)}$$

- Pair of directed walks

$$G(x, y, q) = \frac{1}{x} \left( \frac{{}_1\phi_1(0; y; q, qx)}{{}_1\phi_1(0; y; q, x)} - 1 \right)$$

- Partially directed walks above the diagonal

$$G(x, y, q) = \frac{y}{x} \left( \frac{{}_2\phi_1(0, 0; y^2; q, qx(y - 1/y))}{{}_2\phi_1(0, 0; y^2; q, x(y - 1/y))} - 1 \right)$$

- Directed column-convex polygons

$$G(x, y, q) = y \left( \frac{{}_2\phi_2(0, 0; y, qy; q, qqx(1 - y))}{{}_2\phi_2(0, 0; y, qy; q, qx(1 - y))} - 1 \right)$$



# Outline

- 1 Motivation
  - Vesicle Generating Function
  - Singularity Diagram
  - Scaling Function
- 2 From Lattice Walks to Basic Hypergeometric Series
  - $q$ -Deformed Algebraic Equations
  - $q$ -Difference Equations
  - Basic Hypergeometric Series
- 3 **Asymptotic Analysis**
  - Contour Integral Representation
  - Saddle Point Analysis
  - Generalisation
- 4 Outlook

# A Puzzle

- The full generating function is a quotient of  $q$ -series, e.g.

$$G(t, q) = \frac{\sum_{n=0}^{\infty} \frac{q^{n^2} (-t)^n}{(q; q)_n}}{\sum_{n=0}^{\infty} \frac{q^{n^2 - n} (-t)^n}{(q; q)_n}}$$

# A Puzzle

- The full generating function is a quotient of  $q$ -series, e.g.

$$G(t, q) = \frac{\sum_{n=0}^{\infty} \frac{q^{n^2} (-t)^n}{(q; q)_n}}{\sum_{n=0}^{\infty} \frac{q^{n^2 - n} (-t)^n}{(q; q)_n}}$$

- However, for  $q = 1$  we have a simple algebraic generating function

$$G(t, 1) = \frac{1 - \sqrt{1 - 4t}}{2t}$$

# A Puzzle

- The full generating function is a quotient of  $q$ -series, e.g.

$$G(t, q) = \frac{\sum_{n=0}^{\infty} \frac{q^{n^2} (-t)^n}{(q; q)_n}}{\sum_{n=0}^{\infty} \frac{q^{n^2 - n} (-t)^n}{(q; q)_n}}$$

- However, for  $q = 1$  we have a simple algebraic generating function

$$G(t, 1) = \frac{1 - \sqrt{1 - 4t}}{2t}$$

*How can one understand the limit  $q \rightarrow 1$ ?*

# A Standard Trick For Evaluating Alternating Series

- Write an alternating series as a contour integral

$$\sum_{n=0}^{\infty} (-x)^n c_n = \frac{1}{2\pi i} \oint_{\mathcal{C}} x^s c(s) \frac{\pi}{\sin(\pi s)} ds$$

$\mathcal{C}$  runs counterclockwise around the zeros of  $\sin(\pi s)$

# A Standard Trick For Evaluating Alternating Series

- Write an alternating series as a contour integral

$$\sum_{n=0}^{\infty} (-x)^n c_n = \frac{1}{2\pi i} \oint_{\mathcal{C}} x^s c(s) \frac{\pi}{\sin(\pi s)} ds$$

$\mathcal{C}$  runs counterclockwise around the zeros of  $\sin(\pi s)$

- For example,

$$\exp(-x) = \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} = \frac{1}{2\pi i} \int_{-c-i\infty}^{-c+i\infty} x^s \Gamma(-s) ds$$

where  $c > 0$  (here, we have used  $\Gamma(s)\Gamma(1-s) = \pi/\sin(\pi s)$ )

# A Standard Trick For Evaluating Alternating Series

- Write an alternating series as a contour integral

$$\sum_{n=0}^{\infty} (-x)^n c_n = \frac{1}{2\pi i} \oint_{\mathcal{C}} x^s c(s) \frac{\pi}{\sin(\pi s)} ds$$

$\mathcal{C}$  runs counterclockwise around the zeros of  $\sin(\pi s)$

- For example,

$$\exp(-x) = \sum_{n=0}^{\infty} \frac{(-x)^n}{n!} = \frac{1}{2\pi i} \int_{-c-i\infty}^{-c+i\infty} x^s \Gamma(-s) ds$$

where  $c > 0$  (here, we have used  $\Gamma(s)\Gamma(1-s) = \pi/\sin(\pi s)$ )

*Find suitable  $q$ -version for this trick*

# Contour Integral Representation

Use that

$$\operatorname{Res} [(z; q)_{\infty}^{-1}; z = q^{-n}] = -\frac{(-1)^n q^{\binom{n}{2}}}{(q; q)_n (q; q)_{\infty}} \quad n = 0, 1, 2, \dots$$



# Contour Integral Representation

Use that

$$\text{Res} [(z; q)_{\infty}^{-1}; z = q^{-n}] = -\frac{(-1)^n q^{\binom{n}{2}}}{(q; q)_n (q; q)_{\infty}} \quad n = 0, 1, 2, \dots$$

to prove that

## Lemma

For complex  $t$  with  $|\arg(t)| < \pi$  and  $0 < q < 1$  we have for  $0 < \rho < 1$

$$\sum_{n=0}^{\infty} \frac{q^{n^2-n} (-t)^n}{(q; q)_n} = \frac{(q; q)_{\infty}}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} \frac{z^{\frac{1}{2} \log_q z - \log_q t}}{(z; q)_{\infty}} \sqrt{z} \, dz$$

# Some Asymptotics

Approximate  $\log(z; q)_\infty \sim \frac{1}{\log q} \text{Li}_2(z) + \frac{1}{2} \log(1 - z)$  to get

## Lemma

For  $0 < t < 1$  and with  $\varepsilon = -\log q$

$$\sum_{n=0}^{\infty} \frac{q^{n^2-n}(-t)^n}{(q; q)_n} = \frac{(q; q)_\infty}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{\frac{1}{\varepsilon} \left[ -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \right]} \sqrt{\frac{z}{1-z}} dz [1 + O(\varepsilon)]$$

where  $t < \rho < 1$

# Some Asymptotics

Approximate  $\log(z; q)_\infty \sim \frac{1}{\log q} \text{Li}_2(z) + \frac{1}{2} \log(1 - z)$  to get

## Lemma

For  $0 < t < 1$  and with  $\varepsilon = -\log q$

$$\sum_{n=0}^{\infty} \frac{q^{n^2-n}(-t)^n}{(q; q)_n} = \frac{(q; q)_\infty}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{\frac{1}{\varepsilon} \left[ -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \right]} \sqrt{\frac{z}{1-z}} dz [1 + O(\varepsilon)]$$

where  $t < \rho < 1$

We find a Laplace-type integral, where the saddles are given by

$$0 = \frac{d}{dz} \left[ -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \right]$$

# Saddle Point Analysis

- The asymptotics of

$$\int_C e^{g(z)/\varepsilon} f(z) dz$$

is dominated by the saddles with  $g'(z) = 0$ .

# Saddle Point Analysis

- The asymptotics of

$$\int_C e^{g(z)/\varepsilon} f(z) dz$$

is dominated by the saddles with  $g'(z) = 0$ .

- For  $g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z)$  we find two saddles given by the zeros of

$$z(1-z) = t \quad \Rightarrow \quad z = \frac{1}{2} \pm \frac{1}{2} \sqrt{1-4t}$$

# Saddle Point Analysis

- The asymptotics of

$$\int_C e^{g(z)/\varepsilon} f(z) dz$$

is dominated by the saddles with  $g'(z) = 0$ .

- For  $g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z)$  we find two saddles given by the zeros of

$$z(1-z) = t \quad \Rightarrow \quad z = \frac{1}{2} \pm \frac{1}{2} \sqrt{1-4t}$$

*As  $t$  approaches  $t_t = 1/4$ , the saddles coalesce*

# Coalescing Saddle Points

Standard procedure: reparametrise locally by a cubic and compute a uniform asymptotic expansion

# Coalescing Saddle Points

Standard procedure: reparametrise locally by a cubic and compute a uniform asymptotic expansion

- Reparametrize locally by a cubic  $\Rightarrow$  normal form.



# Coalescing Saddle Points

Standard procedure: reparametrise locally by a cubic and compute a uniform asymptotic expansion

- Reparametrize locally by a cubic  $\Rightarrow$  normal form.  
Write

$$g(z) = \frac{1}{3}u^3 - \alpha u + \beta \text{ with saddles } u_{1,2} = \pm\alpha^{1/2}$$

# Coalescing Saddle Points

Standard procedure: reparametrise locally by a cubic and compute a uniform asymptotic expansion

- Reparametrize locally by a cubic  $\Rightarrow$  normal form.  
Write

$$g(z) = \frac{1}{3}u^3 - \alpha u + \beta \text{ with saddles } u_{1,2} = \pm\alpha^{1/2}$$

$g(z_i) = u_i$  determines  $\alpha$  and  $\beta$ :

$$g(z_1) = -\frac{2}{3}\alpha^{3/2} + \beta \quad g(z_2) = \frac{2}{3}\alpha^{3/2} + \beta$$

# Coalescing Saddle Points

Standard procedure: reparametrise locally by a cubic and compute a uniform asymptotic expansion

- Reparametrize locally by a cubic  $\Rightarrow$  normal form.  
Write

$$g(z) = \frac{1}{3}u^3 - \alpha u + \beta \text{ with saddles } u_{1,2} = \pm\alpha^{1/2}$$

$g(z_i) = u_i$  determines  $\alpha$  and  $\beta$ :

$$g(z_1) = -\frac{2}{3}\alpha^{3/2} + \beta \quad g(z_2) = \frac{2}{3}\alpha^{3/2} + \beta$$

The transformation is one-to-one and analytic in a neighbourhood of  $t = 1/4$ .

# Finally: $\text{Ai}(x)$

- Substitute  $z = z(u)$  into

$$I(\epsilon) = \int_{\mathcal{C}} e^{g(z)/\epsilon} f(z) dz = \int_{\mathcal{C}'} e^{g(z(u))/\epsilon} f(z(u)) \frac{dz}{du} du$$

# Finally: $\text{Ai}(x)$

- Substitute  $z = z(u)$  into

$$I(\epsilon) = \int_C e^{g(z)/\epsilon} f(z) dz = \int_{C'} e^{g(z(u))/\epsilon} f(z(u)) \frac{dz}{du} du$$

and expand

$$f(z) \frac{dz}{du} = \sum_{n=0}^{\infty} (p_n + q_n u) (u^2 - d)^n$$

# Finally: $\text{Ai}(x)$

- Substitute  $z = z(u)$  into

$$I(\epsilon) = \int_C e^{g(z)/\epsilon} f(z) dz = \int_{C'} e^{g(z(u))/\epsilon} f(z(u)) \frac{dz}{du} du$$

and expand

$$f(z) \frac{dz}{du} = \sum_{n=0}^{\infty} (p_n + q_n u) (u^2 - d)^n$$

- Defining

$$V(\lambda) = \frac{1}{2\pi i} \int_{C'} e^{u^3/3 - \lambda u} du$$

arrive at a complete uniform asymptotic expansion for  $I(\epsilon)$

# Finally: $\text{Ai}(x)$

- Substitute  $z = z(u)$  into

$$I(\epsilon) = \int_C e^{g(z)/\epsilon} f(z) dz = \int_{C'} e^{g(z(u))/\epsilon} f(z(u)) \frac{dz}{du} du$$

and expand

$$f(z) \frac{dz}{du} = \sum_{n=0}^{\infty} (p_n + q_n u) (u^2 - d)^n$$

- Defining

$$V(\lambda) = \frac{1}{2\pi i} \int_{C'} e^{u^3/3 - \lambda u} du$$

arrive at a complete uniform asymptotic expansion for  $I(\epsilon)$

- Depending on the contour  $C'$ ,  $V(\lambda)$  is expressible using  $\text{Ai}(\lambda)$  and  $\text{Ai}'(\lambda)$

# Uniform Asymptotics

## Theorem

Let  $0 < t < 1$  and  $\varepsilon = -\log q$ . Then, as  $\varepsilon \rightarrow 0^+$ ,

$$G(t, q) \sim \frac{1}{2} \left( 1 - \sqrt{1 - 4t} \left[ -\frac{\text{Ai}'(\alpha \varepsilon^{-2/3})}{\alpha^{1/2} \varepsilon^{-1/3} \text{Ai}(\alpha \varepsilon^{-2/3})} \right] \right)$$

where  $\alpha = \alpha(t)$  is an explicitly given function of  $t$ . In particular,

$$\alpha(t) \sim 1 - 4t \quad \text{as } t \rightarrow 1/4$$



# Uniform Asymptotics

## Theorem

Let  $0 < t < 1$  and  $\varepsilon = -\log q$ . Then, as  $\varepsilon \rightarrow 0^+$ ,

$$G(t, q) \sim \frac{1}{2} \left( 1 - \sqrt{1 - 4t} \left[ -\frac{\text{Ai}'(\alpha \varepsilon^{-2/3})}{\alpha^{1/2} \varepsilon^{-1/3} \text{Ai}(\alpha \varepsilon^{-2/3})} \right] \right)$$

where  $\alpha = \alpha(t)$  is an explicitly given function of  $t$ . In particular,

$$\alpha(t) \sim 1 - 4t \quad \text{as } t \rightarrow 1/4$$

Some remarks:

- Uniform convergence to  $G(t, 1) = \frac{1}{2} (1 - \sqrt{1 - 4t})$

# Uniform Asymptotics

## Theorem

Let  $0 < t < 1$  and  $\varepsilon = -\log q$ . Then, as  $\varepsilon \rightarrow 0^+$ ,

$$G(t, q) \sim \frac{1}{2} \left( 1 - \sqrt{1 - 4t} \left[ -\frac{\text{Ai}'(\alpha \varepsilon^{-2/3})}{\alpha^{1/2} \varepsilon^{-1/3} \text{Ai}(\alpha \varepsilon^{-2/3})} \right] \right)$$

where  $\alpha = \alpha(t)$  is an explicitly given function of  $t$ . In particular,

$$\alpha(t) \sim 1 - 4t \quad \text{as } t \rightarrow 1/4$$

Some remarks:

- Uniform convergence to  $G(t, 1) = \frac{1}{2} (1 - \sqrt{1 - 4t})$
- Scaling function  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$

# Uniform Asymptotics

## Theorem

Let  $0 < t < 1$  and  $\varepsilon = -\log q$ . Then, as  $\varepsilon \rightarrow 0^+$ ,

$$G(t, q) \sim \frac{1}{2} \left( 1 - \sqrt{1-4t} \left[ -\frac{\text{Ai}'(\alpha\varepsilon^{-2/3})}{\alpha^{1/2}\varepsilon^{-1/3}\text{Ai}(\alpha\varepsilon^{-2/3})} \right] \right)$$

where  $\alpha = \alpha(t)$  is an explicitly given function of  $t$ . In particular,

$$\alpha(t) \sim 1 - 4t \quad \text{as } t \rightarrow 1/4$$

Some remarks:

- Uniform convergence to  $G(t, 1) = \frac{1}{2} (1 - \sqrt{1-4t})$
- Scaling function  $f(z) = -\text{Ai}'(z)/\text{Ai}(z)$
- Stronger than scaling limit which keeps  $z = (1-4t)\varepsilon^{-2/3}$  fixed

# More Saddle Points

Saddle point coalescence occurs in all four cases:

- Dyck paths,  ${}_0\phi_1(-; 0; q, -t)$ :

$$g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \quad \Rightarrow \quad (z-1)z + t = 0$$

# More Saddle Points

Saddle point coalescence occurs in all four cases:

- Dyck paths,  ${}_0\phi_1(-; 0; q, -t)$ :

$$g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \Rightarrow (z-1)z + t = 0$$

- Pair of directed walks,  ${}_1\phi_1(0; y; q, x)$ :

$$g(z) = -\text{Li}_2(y/z) + \log(z) \log(x) + \text{Li}_2(z) \Rightarrow (z-1)(z-y) + zx = 0$$

# More Saddle Points

Saddle point coalescence occurs in all four cases:

- Dyck paths,  ${}_0\phi_1(-; 0; q, -t)$ :

$$g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \Rightarrow (z-1)z + t = 0$$

- Pair of directed walks,  ${}_1\phi_1(0; y; q, x)$ :

$$g(z) = -\text{Li}_2(y/z) + \log(z) \log(x) + \text{Li}_2(z) \Rightarrow (z-1)(z-y) + zx = 0$$

- Part. directed walks above the diagonal,  ${}_2\phi_1(0, 0; y^2; q, x(y-1/y))$ :

$$g(z) = \dots \Rightarrow (z-1)(z-y^2) + z^2x(1/y - y) = 0$$

# More Saddle Points

Saddle point coalescence occurs in all four cases:

- Dyck paths,  ${}_0\phi_1(-; 0; q, -t)$ :

$$g(z) = -\frac{1}{2}(\log z)^2 + \log(z) \log(t) + \text{Li}_2(z) \Rightarrow (z-1)z + t = 0$$

- Pair of directed walks,  ${}_1\phi_1(0; y; q, x)$ :

$$g(z) = -\text{Li}_2(y/z) + \log(z) \log(x) + \text{Li}_2(z) \Rightarrow (z-1)(z-y) + zx = 0$$

- Part. directed walks above the diagonal,  ${}_2\phi_1(0, 0; y^2; q, x(y-1/y))$ :

$$g(z) = \dots \Rightarrow (z-1)(z-y^2) + z^2x(1/y - y) = 0$$

- Directed column-convex polygons,  ${}_2\phi_2(0, 0; y, qy; q, qx(1-y))$ :

$$g(z) = \dots \Rightarrow (z-1)(z-y)^2 + z^2x(1-y) = 0$$

# Contour Integral for Basic Hypergeometric Series

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, t \right) = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q, b_1, \dots, b_s; q)_n} \left[ (-1)^n q^{\binom{n}{2}} \right]^{1+s-r} t^n$$



# Contour Integral for Basic Hypergeometric Series

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, t \right) = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q, b_1, \dots, b_s; q)_n} \left[ (-1)^n q^{\binom{n}{2}} \right]^{1+s-r} t^n$$

## Lemma

For complex  $t$  with  $|\arg(t)| < \pi$ , and  $0 < q < 1$  we have for  $0 < \rho < 1$

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, (-1)^{s-r} t \right) = \frac{(a_1, \dots, a_r; q)_{\infty}}{(b_1, \dots, b_s; q)_{\infty}} \times \\ \frac{(q; q)_{\infty}}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} \frac{(b_1/z, \dots, b_s/z; q)_{\infty}}{(a_1/z, \dots, a_r/z; q)_{\infty}} \frac{z^{\frac{s-r}{2} \log_q z - \log_q t}}{(z; q)_{\infty}} z^{\frac{s-r}{2}} dz$$

# Contour Integral for Basic Hypergeometric Series

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, t \right) = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_r; q)_n}{(q, b_1, \dots, b_s; q)_n} \left[ (-1)^n q^{\binom{n}{2}} \right]^{1+s-r} t^n$$

## Lemma

For complex  $t$  with  $|\arg(t)| < \pi$ , and  $0 < q < 1$  we have for  $0 < \rho < 1$

$${}_r\phi_s \left( \begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, (-1)^{s-r} t \right) = \frac{(a_1, \dots, a_r; q)_{\infty}}{(b_1, \dots, b_s; q)_{\infty}} \times \\ \frac{(q; q)_{\infty}}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} \frac{(b_1/z, \dots, b_s/z; q)_{\infty}}{(a_1/z, \dots, a_r/z; q)_{\infty}} \frac{z^{\frac{s-r}{2} \log_q z - \log_q t}}{(z; q)_{\infty}} z^{\frac{s-r}{2}} dz$$

**Proof:** Compute residues at  $z = q^{-n}$  for  $n \in \mathbb{N}_0$  (and consider a sequence of contours)

# General Saddle Point Equation

- As  $\varepsilon = -\log q \rightarrow 0$ , we again obtain a Laplace-type integral

$$\int e^{g(z)/\varepsilon} f(z) dz$$

where

$$\begin{aligned} g(z) = & \operatorname{Li}_2(a_1/z) + \dots + \operatorname{Li}_2(a_r/z) + \operatorname{Li}_2(z) + \log z \log t \\ & - \operatorname{Li}_2(b_1/z) - \dots - \operatorname{Li}_2(b_s/z) - \frac{s-r}{2} (\log z)^2 \end{aligned}$$

# General Saddle Point Equation

- As  $\varepsilon = -\log q \rightarrow 0$ , we again obtain a Laplace-type integral

$$\int e^{g(z)/\varepsilon} f(z) dz$$

where

$$g(z) = \text{Li}_2(a_1/z) + \dots + \text{Li}_2(a_r/z) + \text{Li}_2(z) + \log z \log t \\ - \text{Li}_2(b_1/z) - \dots - \text{Li}_2(b_s/z) - \frac{s-r}{2} (\log z)^2$$

- $g'(z) = 0$  yields a polynomial saddle point equation

$$t(a_1 - z) \dots (a_r - z) + (1 - z)(b_1 - z) \dots (b_s - z) = 0$$

# General Saddle Point Equation

- As  $\varepsilon = -\log q \rightarrow 0$ , we again obtain a Laplace-type integral

$$\int e^{g(z)/\varepsilon} f(z) dz$$

where

$$g(z) = \text{Li}_2(a_1/z) + \dots + \text{Li}_2(a_r/z) + \text{Li}_2(z) + \log z \log t \\ - \text{Li}_2(b_1/z) - \dots - \text{Li}_2(b_s/z) - \frac{s-r}{2} (\log z)^2$$

- $g'(z) = 0$  yields a polynomial saddle point equation

$$t(a_1 - z) \dots (a_r - z) + (1 - z)(b_1 - z) \dots (b_s - z) = 0$$

Work in progress ...

# Outline

- 1 Motivation
  - Vesicle Generating Function
  - Singularity Diagram
  - Scaling Function
- 2 From Lattice Walks to Basic Hypergeometric Series
  - $q$ -Deformed Algebraic Equations
  - $q$ -Difference Equations
  - Basic Hypergeometric Series
- 3 Asymptotic Analysis
  - Contour Integral Representation
  - Saddle Point Analysis
  - Generalisation
- 4 Outlook

# Outlook

So far:

- simple  $q$ -algebraic equation
- simple  $q$ -series solution
- contour integral
- saddle-point analysis

# Outlook

So far:

- simple  $q$ -algebraic equation
- simple  $q$ -series solution
- contour integral
- saddle-point analysis

Can be generalised to

- asymptotics for general  ${}_m\phi_n$



# Outlook

So far:

- simple  $q$ -algebraic equation
- simple  $q$ -series solution
- contour integral
- saddle-point analysis

Can be generalised to

- asymptotics for general  ${}_m\phi_n$

More ambitiously

- get the asymptotics directly from the functional equation  
(without solving it first!)

# Outlook

So far:

- simple  $q$ -algebraic equation
- simple  $q$ -series solution
- contour integral
- saddle-point analysis

Can be generalised to

- asymptotics for general  ${}_m\phi_n$

More ambitiously

- get the asymptotics directly from the functional equation  
(without solving it first!)

*The End*