The pressure of surface-attached polymers and vesicles

Thomas Prellberg¹ and Aleks Owczarek²

¹Queen Mary University of London ²The University of Melbourne

Open Statistical Physics 2015



Topic Outline

- Surface-grafted polymers
- Exactly Solvable Models
- 3 Conclusion

Outline

- Surface-grafted polymers
- 2 Exactly Solvable Models
- Conclusion

Surface-grafted polymers



A polymer grafted to a surface exerts pressure on it

Bickel et al, PRE 62 (2000) 1124

Surface-grafted polymers



A polymer grafted to a surface exerts pressure on it


Bickel et al, PRE 62 (2000) 1124

Related to change of entropy




From Polymer to Lattice Model

Coarse-grained "beads on a necklace"





From Polymer to Lattice Model

Coarse-grained "beads on a necklace"




Self-avoiding walk on a regular lattice





 C_n : number of *n*-step polymers

 $C_n(r)$: number of *n*-step polymers avoiding site at distance r




Pressure of polymer on surface

$$P_n(r) = -\log \frac{C_n(r)}{C_n}$$

 C_n : number of *n*-step polymers

 $C_n(r)$: number of *n*-step polymers avoiding site at distance r




Pressure of polymer on surface

$$P_n(r) = -\log \frac{C_n(r)}{C_n}$$

$$P_n(r) \to P(r) \text{ as } n \to \infty$$

 C_n : number of *n*-step polymers

 $C_n(r)$: number of *n*-step polymers avoiding site at distance r



 C_n : number of *n*-step polymers

 $C_n(r)$: number of *n*-step polymers avoiding site at distance r

Pressure of polymer on surface

$$P_n(r) = -\log \frac{C_n(r)}{C_n}$$


$$P_n(r) \to P(r) \text{ as } n \to \infty$$

Gaussian chains (Bickel, 2000)

$$P_G(r) = \frac{\Gamma(d/2)}{\pi^{d/2}} \frac{1}{(r^2+1)^{d/2}}$$



SAW versus Gaussian chain



extrapolated from exact enumeration data up to n=59, r=10

Jensen et al., J. Phys. 46 (2013) 115004 \sim \sim

Outline

- Surface-grafted polymers
- Exactly Solvable Models
 - Directed Walk Models
 - Dyck paths
 - Adsorbing Dyck paths
 - Area-weighted Dyck paths
- 3 Conclusion

Directed walk models

Directed walk models

Directed walks (both/one ends grafted)



Directed walk models

Directed walks (both/one ends grafted)



Pairs of directed walks



Directed walk models


Directed walks (both/one ends grafted)



Pairs of directed walks



Partially directed walks



Directed walk models


Directed walks (both/one ends grafted)



Pairs of directed walks



Partially directed walks



Moreover, add contact weights, area weights, nearest-neighbour interactions, . . .

Much activity ensued ...

So far

- Directed walks with contacts
 - E J Janse van Rensburg and TP, J. Phys. A **46** (2013) 115202
- Pairs of walks with contacts
 - E J Janse van Rensburg and TP, J. Phys. A 46 (2013) 115202
- Directed walks with area
- A Owczarek and TP, J. Phys. A 47 (2014) 215001
- Partially directed walks with contacts
 - E J Janse van Rensburg, and TP, in preparation

Much activity ensued ...

So far

- Directed walks with contacts
 - E J Janse van Rensburg and TP, J. Phys. A 46 (2013) 115202
- Pairs of walks with contacts
 - E J Janse van Rensburg and TP, J. Phys. A 46 (2013) 115202
- Directed walks with area
- A Owczarek and TP, J. Phys. A 47 (2014) 215001
- Partially directed walks with contacts
 - E J Janse van Rensburg, and TP, in preparation

The next model to be done

Interacting partially directed walks

Much activity ensued . . .

So far

Directed walks with contacts

E J Janse van Rensburg and TP, J. Phys. A **46** (2013) 115202

Pairs of walks with contacts

E J Janse van Rensburg and TP, J. Phys. A 46 (2013) 115202

Directed walks with area

A Owczarek and TP, J. Phys. A 47 (2014) 215001

Partially directed walks with contacts

E J Janse van Rensburg, and TP, in preparation

The next model to be done

Interacting partially directed walks

Also

Rooted self-avoiding polygons

F Gassoumov and E J Janse van Rensburg, J. Stat. Mech. (2013) P10005

Directed walks with both ends tethered \Leftrightarrow Dyck paths



Directed walks with both ends tethered ⇔ Dyck paths



• 2n-step Dyck paths are counted by the n-th Catalan number C_n

$$D_{2n} = C_n \equiv \frac{1}{n+1} \binom{2n}{n} \sim \frac{4^n}{\sqrt{\pi} n^{3/2}}$$

Directed walks with both ends tethered ⇔ Dyck paths



• 2n-step Dyck paths are counted by the n-th Catalan number C_n

$$D_{2n} = C_n \equiv \frac{1}{n+1} \binom{2n}{n} \sim \frac{4^n}{\sqrt{\pi} n^{3/2}}$$

• Exclude paths touching the surface at distance 2r

$$D_{2n}(r) = D_{2n} - D_{2r}D_{2n-2r}$$





Compute pressure

$$P_{2n}(r) = \log \frac{D_{2n}(r)}{D_{2n}} = -\log \left(1 - \frac{C_r C_{n-r}}{C_n}\right)$$



Compute pressure


$$P_{2n}(r) = \log \frac{D_{2n}(r)}{D_{2n}} = -\log \left(1 - \frac{C_r C_{n-r}}{C_n}\right)$$

Limiting pressure

$$P_{2n}(2r) \sim rac{1}{\sqrt{\pi}(na(1-a))^{3/2}} \qquad \qquad r=na, \; n o \infty$$
 $P(r) \sim rac{1}{\sqrt{\pi}r^{3/2}} \qquad \qquad ext{first } n o \infty ext{, then } r o \infty$

Adsorbing Dyck paths

Surface contact weight z



Adsorbing Dyck paths

Surface contact weight z



Exact partition function

$$Z_n(z) = \sum_{m=0}^{\lfloor n/2 \rfloor} \frac{4m+2}{n+2(m+1)} \binom{n}{\lfloor n/2 \rfloor + m} (z-1)^m$$

Adsorbing Dyck paths

Surface contact weight z



Exact partition function

$$Z_n(z) = \sum_{m=0}^{\lfloor n/2 \rfloor} \frac{4m+2}{n+2(m+1)} \binom{n}{\lfloor n/2 \rfloor + m} (z-1)^m$$

Limiting pressure

$$P_n^D(z; 2 \lfloor an/2 \rfloor) \simeq \begin{cases} \frac{8}{\sqrt{2\pi n^3 a^3 (1-a)^3} \log^2(z-1)}, & \text{if } z < 2; \\ \frac{\sqrt{2}}{\sqrt{\pi n a (1-a)}} & \text{if } z = 2; \\ \log(z-1) + \frac{A}{12na(1-a)(z-1)(z-2)}, & \text{if } z > 2; \end{cases}$$

for *n* even, with
$$A = (a^2 - a + 1)(z^4 + 8z^3 + 30z^2 - 32z + 16)$$


Area-weighted Dyck paths

Dyck paths weighted by enclosed area, area weight \boldsymbol{q}



Area-weighted Dyck paths

Dyck paths weighted by enclosed area, area weight q




• Replace the *n*-th Catalan number C_n by $C_n(q)$

$$C_0(q) = 1 \; , \quad C_{n+1}(q) = \sum_{k=0}^n q^k C_k(q) C_{n-k}(q) \quad n \geq 0$$

Area-weighted Dyck paths

Dyck paths weighted by enclosed area, area weight q



• Replace the *n*-th Catalan number C_n by $C_n(q)$

$$C_0(q) = 1 \; , \quad C_{n+1}(q) = \sum_{k=0}^n q^k C_k(q) C_{n-k}(q) \quad n \geq 0$$

• Pressure $P_{2n}(q,r)$ is now given by

$$P_{2n}(q,r) = -\log\left(\frac{C_n(q) - C_r(q)C_{n-r}(q)}{qC_n(q)}\right)$$



Are-weighted Dyck paths

G(t,q) is fairly well understood, $C_n(q)$ much less

Are-weighted Dyck paths

G(t,q) is fairly well understood, $C_n(q)$ much less

Continued fraction representation

$$G(t,q) = rac{1}{1-rac{t}{1-rac{qt}{1-rac{q^2t}{1-\dots}}}}$$

Are-weighted Dyck paths

G(t,q) is fairly well understood, $C_n(q)$ much less

Continued fraction representation

$$G(t,q)=rac{1}{1-rac{t}{1-rac{qt}{1-rac{q^2t}{1-\dots}}}$$

Alternatively

$$G(t,q) = rac{A_q(t)}{A_q(t/q)}$$

where

$$A_q(t) = \sum_{n=0}^{\infty} \frac{q^{n^2}(-t)^n}{\prod_{k=1}^n (1-q^k)}$$

is Ramanujan's Airy function



Scaling for q o 1


Scaling function near q=1 and $t=t_c(1)=1/4$

$$f(s) = \lim_{q \to 1^{-}} \left(2 - (1 - q)^{-1/3} G(1/4 - s(1 - q)^{2/3}, q) \right) = -2 \frac{\operatorname{Ai}'(4s)}{\operatorname{Ai}(4s)}$$

Scaling for q o 1

Scaling function near q=1 and $t=t_c(1)=1/4$

$$f(s) = \lim_{q \to 1^{-}} \left(2 - (1 - q)^{-1/3} G(1/4 - s(1 - q)^{2/3}, q) \right) = -2 \frac{\operatorname{Ai}'(4s)}{\operatorname{Ai}(4s)}$$



see talk by Nils Haug today, N Haug and TP, arXiv:1412.5108



Compute pressure in the bulk using surface contacts

Weight surface contacts by z

$$G(t,q;\mathbf{z}) = \frac{1}{1 - \frac{t\mathbf{z}}{1 - \frac{qt}{1 - \frac{q^2t}{1 - \dots}}}}$$

Compute pressure in the bulk using surface contacts

Weight surface contacts by z

$$G(t,q;\mathbf{z}) = rac{1}{1-rac{t\mathbf{z}}{1-rac{qt}{1-rac{q^2t}{1-\dots}}}}$$

Equivalently

$$G(t,q;z) = \frac{1}{1 - tzG(qt,q)}$$

• Density of contacts $\rho(q)$ is related to singularity $t_c(q;z)$

$$\rho(q) = -\left. \frac{\partial \log t_c(q; z)}{\partial \log z} \right|_{z=1}$$


• Density of contacts $\rho(q)$ is related to singularity $t_c(q;z)$


$$\rho(q) = -\left. \frac{\partial \log t_c(q; z)}{\partial \log z} \right|_{z=1}$$

• We find for the pressure P(q)

$$P(q) = -\log(1 - \rho(q)) + \log q$$


Bulk pressure for n = 10 to 80 (left) and in the TDL (right)






Pressure Profiles

Pressure profiles for q=0.98 (left), 1.00 (center), and 1.02 (right)



Pressure Profiles

Pressure profiles for q=0.98 (left), 1.00 (center), and 1.02 (right)



Rate of convergence: ρ^n (left), $n^{-1/2}$ (center), and ρ^{n^2} (right)

Outline

- Surface-grafted polymers
- Exactly Solvable Models
- Conclusion

• Entropy-driven pressure of polymers

- Entropy-driven pressure of polymers
- Gaussian Chain and Self-Avoiding Walk

- Entropy-driven pressure of polymers
- Gaussian Chain and Self-Avoiding Walk
- Exactly solvable directed walk model with
 - contacts
 - area

- Entropy-driven pressure of polymers
- Gaussian Chain and Self-Avoiding Walk
- Exactly solvable directed walk model with
 - contacts
 - area

Simple lattice models still manage to give useful physical insight!