

−3
0
3
Proline
Cystathionine
Homocysteine
Methionine
SAM
Isoleucine
Valine
Leucine
D−Ribulose 5−ph\
5−Phospho−alpha)
Glycerone phosp\
(2R)−2−Hydroxy−3−(phosphonooxy)−propanal
D−Xylulose 5−ph
D−Fructose 1,6−bisphosphate
D−Fructose 6−ph
D−Erythrose 4−ph
2−Dehydro−3−deo\
3−Dehydroshikim\
Shikimate
5−O−(1−Carboxyv\
Shikimate 3−phos
Chorismic acid
Prephenic acid
Phenylpyruvate
Phenylalanine
Arogenate
Tyrosine
Phosphoenolpyru\
3−Phospho−D−glyceroyl phosphate
Alanine
(S)−2−Acetolact\
3−Phospho−D−glycerate
S−Sulfo−L−cyste\
Serine
O−Acetyl−L−serine
Cysteine
Glycine
O−Phospho−L−serine
Threonine
(S)−2−Hydroxy−2−ethyl−3−oxobutanoate
L−Homoserine
D−erythro−3−Methylaspartate 4−s\
L−4−Aspartyl
meso−2,6−Diamin\
Lysine
C20258
C03972
N−Acetyl−L−glut\
N−Acetyl−L−glutamate 5−semialdehyde
N2−Acetyl−L−orn\
L−Ornithine
C15532
Carbamoyl phosph\
L−Citrulline
L−Argininosucci\
Arginine
2−Hydroxytricarballylic acid
2−Hydroxysuccinic acid
1−Hydroxypropan\
1,2,3−tricarb\
oxylic acid
alpha−Ketogluta\
Glutamic acid
Glutamine
Asparagine
L−Glutamyl 5−phosphate
L−Glutamate 5−s\
−Data with KEGG pathway−
−Rendered by Pathview−

KEGG diagram legend

Edge Types

compound

hidden compound

activation

inhibition

expression

repression

indirect effect

state change

binding/association

dissociation

phosphorylation +p

dephosphorylation −p

glycosylation +g

ubiquitination +u

methylation +m

others/unknown ?

Node Types

gene (protein/enzyme)

group (complex)

compound (metabolite/glycan)

map (pathway) Pathway name