# Discrete Spacetime with a Cosmological Cutoff

How the finiteness of the universe could tame the non-locality of Lorentz-invariant discreteness

Draft v2 — internal notebook checkpoint

25 October 2026

Not for circulation. Claims are tagged established, literature or ours.

**Abstract**

We ask whether spacetime can be a discrete set of events that looks like a smooth Lorentzian manifold from far away, under three requirements: discreteness comes first, there is no background structure (only relations between events), and local Lorentz invariance holds, although a *cosmological* preferred frame is allowed because one is observed. Defining neighbours as events with nothing causally between them, an elementary calculation shows that Lorentz invariance gives every event one neighbour per unit of rapidity, and therefore infinitely many in an infinite universe. This is a known result of causal set theory. Our contribution is to take the finite size of the universe seriously as the physical cutoff. Neighbours then extend only up to a maximum rapidity $\eta_{\max} \simeq \ln(2L/\ell_{\mathrm{P}}) \approx 142$, set by the ratio of the size of the universe to the Planck length. This is far beyond the largest rapidities at which Lorentz kinematics has been tested ($\eta \approx 25$–$27$), and the frame it singles out is the cosmological one. A background-free toy model, a causal diamond analysed through its order relation alone, confirms the logarithmic neighbour count numerically and shows that proper time measured by counting carries an intrinsic fuzziness scaling as $n^{1/6}$. Finally, a classical growth model in which events are created one at a time defines cosmic time and a cosmological frame purely by counting, and with a suitable rule produces an expanding universe from a single first event. We discuss what this does and does not achieve, notably that in $3+1$ dimensions the cutoff leaves $\sim 10^{124}$ neighbours per event, that gravity has not yet been recovered, and that the expanding growth rule secretly uses the birth order as a clock.

## 1 The question and the rules

In 1900 Planck assumed that energy comes in discrete quanta. We ask the analogous question for spacetime: can it be a discrete set of events that only *looks* continuous from far away, with general relativity (GR) as its large-scale description? The working constraints, refined over two sessions, are:

**C1 Discrete spacetime.** Events and the causal order between them; the continuum is a large-scale limit.

**C2 No background.** No absolute space (Newton) and no fixed aether; every statement is about relations between events.

**C3 Non-locality regulated by the finite size of the universe** (Sec. 3).

**C4 Local Lorentz invariance** for all boosts well below the cutoff.

**C5 A cosmological frame produced by the universe itself.** It should coincide with the rest frame of matter, i.e. the cosmic microwave background (CMB) frame established. This is a test, not an input.

**C6 Newton and Einstein from far away.**

**C7 Creation (and perhaps destruction) of events drives expansion**, possibly through quantum creation operators.

**C8 The links between events are quantum.**

Missing so far is *matter*: nothing in the model yet plays the role of what gravitates.

C2 and C5 rest on a distinction worth stating explicitly. A *fixed background* is structure the laws refer to but which nothing affects; it is forbidden. A *dynamical preferred-frame field*, such as the Einstein-aether of [1], is not a background but breaks local Lorentz invariance. A *preferred frame of the state*, such as the CMB frame, comes from the contents of the universe while the laws stay invariant. GR is consistent with such a frame but does not explain it. A discrete model that *produced* it while leaving local physics invariant would be interesting.

# 2 Neighbours in a Lorentz-invariant discrete spacetime

## 2.1 Two failures that fix the vocabulary

**Regular lattices.** literature If neighbours are lattice sites one hop apart, the set of events reachable in $N$ ticks is a polygon (a diamond on a square lattice, a hexagon on a hexagonal one), never a circle. This follows from Burago's theorem on periodic metrics [2] and is the rigorous form of Weyl's tile argument [3]. A lattice predicts a direction-dependent speed of light and has a rest frame of its own. Random structures avoid both problems [4].

**Neighbours at a fixed proper time.** Requiring neighbours at *exactly* one tick of proper time picks out a curve, which random points never land on. Smearing the points ("quantum blur") gives the curve a thickness but does not change the conclusion below.

## 2.2 Links

literature We adopt the causal-set notion [5, 6]: $y$ is a *neighbour* (a *link*) of $x$ if $y$ is in the causal future of $x$ and no other event lies causally between them. This uses only the order relation, so it complies with C2.

## 2.3 One neighbour per unit of rapidity

Work in $1+1$ dimensions with $c = 1$, with events distributed with density $\rho$ (one per Planck area, say). Use light-cone coordinates $u = t - x$, $v = t + x$, and write a future event as $u = \tau e^{-\eta}$, $v = \tau e^{\eta}$. Here $\tau$ is its proper time from the origin and $\eta$ the rapidity of the straight worldline joining them, with velocity $\tanh \eta$. The area element is $dt\,dx = \tau\, d\tau\, d\eta$. The causal interval between the origin and that event has area $\tau^2/2$, so the probability that it is empty, i.e. that the event is a link, is $e^{-\rho\tau^2/2}$. Hence

$$\langle N_{\text{links}} \rangle = \int d\eta \int_0^\infty \rho\, \tau\, e^{-\rho\tau^2/2}\, d\tau = \int d\eta \cdot 1. \tag{1}$$

ours (elementary form of a known result.) *There is exactly one expected neighbour per unit of rapidity.* A boost shifts $\eta$, so it maps the neighbours in one rapidity band onto those in the next (Fig. 1). If all rapidities are allowed, every event has infinitely many neighbours. Most of them sit close to the light cone, far away in space but only a Planck time away in proper time.





Figure 1: (a) Where the neighbours of an event live: near the light cone, coloured by rapidity band. (b) The same region in coordinates $(\eta, \tau)$: an infinite strip of equivalent boxes, each containing one neighbour on average. A boost slides the strip sideways like a conveyor belt.

**The dilemma.** literature This is the elementary core of a theorem of Bombelli, Henson and Sorkin [7]: no discrete structure can give each event finitely many neighbours without singling out a frame. The reason is that a normalisable distribution cannot be uniform in rapidity, because rapidity, unlike angle, has no end. Lorentz-invariant discreteness is therefore non-local [8]. Attempts to escape by making the points "quantum blurry" run into the same obstruction: a single neighbour in superposition over all rapidity bands, with equal weights as boost invariance demands, cannot be normalised.

# 3 The cosmological cutoff

## 3.1 What rapidity is

established Rapidity $\eta$ is the natural measure of a boost. Velocity is $v = \tanh\eta$ and the Lorentz factor is $\gamma = \cosh\eta$. Unlike velocities, rapidities simply *add* under successive boosts along a line. A boost by $\eta$ stretches one light-cone direction by $e^{\eta}$ and squeezes the other by $e^{-\eta}$. Some reference points:

| | $\gamma$ | $\eta \approx \ln 2\gamma$ |
|---|---|---|
| Earth relative to the CMB frame (370 km/s) | $1 + 10^{-6}$ | 0.0012 |
| LHC proton (6.8 TeV) | $7 \times 10^{3}$ | 9.6 |
| GZK threshold proton ($5 \times 10^{19}$ eV) | $5 \times 10^{10}$ | 25 |
| Highest-energy cosmic ray ($3.2 \times 10^{20}$ eV, if a proton) | $3.4 \times 10^{11}$ | 27 |
| **Cutoff $\eta_{\max}$ (this note)** | $\sim 10^{61}$ | **142** |

The GZK suppression of cosmic rays [9] is a genuine kinematic test at $\eta \approx 25$. It requires that a CMB photon, seen from the rest frame of a $\gamma \sim 10^{11}$ proton, has the boosted energy predicted by Lorentz transformations, and the observed suppression is consistent with that.

## 3.2 A finite universe caps the rapidity

ours A neighbour at proper time $\tau$ and rapidity $\eta$ lies at spatial distance $r = \tau \sinh\eta \approx (\tau/2)\, e^{\eta}$ from the event. In a universe of size $L$ this cannot exceed $L$, so

$$\eta_{\max} \simeq \ln \frac{2L}{\tau} \approx \ln \frac{2L}{\ell_{\mathrm{P}}} \approx 142, \tag{2}$$

using $L \approx 4.4 \times 10^{26}\,\mathrm{m}$ (radius of the observable universe) and $\tau \sim \ell_{\mathrm{P}} = 1.6 \times 10^{-35}\,\mathrm{m}$. Physically, $\eta_{\max}$ is the most extreme boost under which a Planck-sized neighbourhood still

fits inside the universe: $e^{142} \approx 10^{61.7} \approx 2L/\ell_{\mathrm{P}}$. Because the dependence on the inputs is logarithmic, the number is robust. Changing $L$ or the discreteness scale by a factor $10^{10}$ shifts it by only $\pm 23$.

By Eq. (1), an event in $1+1$ dimensions has about $2\eta_{\max} \approx 280$ future neighbours. That is finite and small. In $3+1$ dimensions the space of boost directions is a hyperboloid whose area grows like $\sinh^2\eta$. With $d^4x = \tau^3 \sinh^2\eta \, d\tau \, d\eta \, d\Omega$ and causal-interval volume $\pi\tau^4/24$,

$$N_{\mathrm{links}}^{(3+1)} \approx 2\pi\rho L^2 \int_0^\infty \tau \, e^{-\rho\pi\tau^4/24} d\tau = \frac{\pi\sqrt{24}}{2}\left(\frac{L}{\ell_{\mathrm{P}}}\right)^2 \approx 10^{124}. \tag{3}$$

### 3.3 What the cutoff buys

- **A finite neighbour count, set by cosmology.** It is small in $1+1$ dimensions and huge in $3+1$.
- **A preferred frame of the right kind.** The only frame singled out is that of the universe as a whole, cf. the compact-space twin paradox [10]. That is a frame of the *state*, compatible with C5. In relational terms it is the frame of the universe's own longest causal chain.
- **Consistency with experiment.** Local Lorentz invariance holds for $\eta \ll 142$. Tests reach $\eta \approx 25$–$27$.
- **Causality intact.** All neighbours are inside the light cone. The non-locality is "light-cone shaped": the emission and absorption of a photon from a distant galaxy are almost neighbours.

## 4 Relations only: a background-free toy model

To comply with C2 we analyse a finite $1+1$ "universe", a causal diamond, using only its order relation. As in causal set theory, we generate test configurations by Poisson sprinkling, a stand-in for unknown dynamics, and then discard all coordinates. In light-cone coordinates a sprinkled diamond is equivalent to a random permutation, which makes two exact statements available.

**Proper time by counting.** literature The proper time between the tips of the diamond is estimated by the length $\mathcal{L}$ of the longest causal chain [11]. In $1+1$ dimensions this is the longest increasing subsequence of a random permutation (Ulam's problem), for which $\langle\mathcal{L}\rangle \simeq 2\sqrt{n} - 1.77\, n^{1/6}$ with Tracy–Widom fluctuations $\delta\mathcal{L} \simeq 0.90\, n^{1/6}$ [12]. With $n = \rho\tau^2/2$ this gives

$$\frac{\delta\tau}{\tau} \simeq 0.45\, n^{-1/3}. \tag{4}$$

ours (interpretation) This is the background-free version of our earlier "quantum blur" intuition: geometry measured by counting is *intrinsically fuzzy*. The fuzziness is statistical, not yet quantum, and it is a relation between events, not a smearing in some background space.

**Neighbours by counting.** established The links of an event are the minimal elements of its future. Ordered along one light-cone direction, they are the running minima ("records") along the other. For $m$ events in the future the expected number of records is the harmonic number $H_m = 1 + \frac{1}{2} + \cdots + \frac{1}{m} \simeq \ln m + 0.577$. For an event at the centre of a diamond of $n$ events, $m \approx n/4$ and $\langle N_{\mathrm{links}}\rangle \simeq \ln n - 0.81$. Since $\ln n \simeq 2\ln(L/\ell_{\mathrm{P}})$, this is the $2\eta_{\max}$ of Sec. 3, now derived without any background.

| $n$ | $\langle \mathcal{L} \rangle$ | $2\sqrt{n}$ | $\delta\mathcal{L}/n^{1/6}$ | $\langle N_{\text{links}} \rangle$ | $\langle H_m \rangle$ |
|---|---|---|---|---|---|
| $10^2$ | 16.7 | 20.0 | 0.75 | 3.78 | 3.80 |
| $10^3$ | 58.4 | 63.2 | 0.92 | 6.09 | 6.10 |
| $10^4$ | 192.1 | 200.0 | 0.89 | 8.43 | 8.40 |
| $10^5$ | 620.6 | 632.5 | 0.92 | 10.76 | 10.70 |
| $10^6$ | — | — | — | 13.04 | 13.01 |

Table 1: Numerics for sprinkled causal diamonds (`sims/causal_diamond.py`; 100–2000 samples per row). The longest chain approaches $2\sqrt{n}$ with fluctuations $\approx 0.9\, n^{1/6}$ (Tracy–Widom value 0.90). The neighbour count of the central event matches the harmonic number, i.e. grows like $\ln n$.





Figure 2: Size of a grown universe against cosmic time, both measured by counting ($N = 4000$ events, `sims/growth_percolation.py`; first 60 layers of the fixed-$p$ run shown). The last few layers of the expanding run are still being filled: that is "the present". Interactive version: `viz/growing-universe.html`.

## 5 Growth: a universe that creates its events

literature If the number of events measures spacetime volume ("Number = Volume" [6]), an expanding universe must keep producing new events. In causal set theory this is literal: in sequential growth dynamics [13], the passage of time *is* the birth of new elements.

### 5.1 A classical growth toy

ours (implementation of a known model family). Events are born one at a time. The newborn chooses each existing event as a direct cause with probability $p$, with at least one cause ("originary" growth), and inherits their entire past. Only the order relation is used afterwards. We measure:

- **Cosmic time** $h(x)$: the length of the longest chain of causes from the first event to $x$. This is the discrete analogue of the GR *cosmological time function*, the supremum of proper times back to the initial singularity [16]. Its level sets (layers) define a cosmological frame by counting alone, as C5 requires.
- **Size of the universe** at cosmic time $h$: the number of events in layer $h$.
- **Valency**: the number of past links per event.

Two rules give two universes (Fig. 2). With *fixed* $p$ the universe quickly reaches a fixed width and stays there, a steady state. With $p_n = c/n$, about $c$ causes per newborn, a single first event is followed by layers of $1, 2, 3, 3, 5, 7, \ldots, \sim 280$ events: *an expanding universe from a "bang", produced only by creating events.* The mean valency stays small, about 2.8.

**Caveat.** A rule in which $p$ depends on $n$ uses the *birth order* as a hidden clock. The covariant growth rules of [13] forbid exactly this, because birth order is not observable (C2). The covariant route to the same behaviour is cosmological: cycles of expansion and contraction renormalise the effective $p$ downward [17], and de Sitter-like expansion has been reported after an origin element [18]. An open hypothesis (Roberto) is that quantum creation operators remove the hidden clock.

### 5.2 Quantum creation and quantum links: existing homes

literature C7 and C8 match active programmes. In *group field theory cosmology* [19, 20], quanta of geometry are created by field operators, the universe is a condensate of them, and Friedmann equations with a bounce emerge. In *quantum graphity* [21], each link is a qubit (on/off), and geometry appears as the low-energy connectivity. In *quantum sequential growth* [22], probabilities are replaced by amplitudes, so different growth histories interfere. Combining these with the cosmological cutoff of Sec. 3 is the direction of stage S3 below.

**Cutoff and growth.** In this picture the cutoff grows with cosmic time, $\eta_{\max}(t) \simeq \ln[2L(t)/\ell_{\mathrm{P}}]$. Sorkin's heuristic $\Lambda \sim 1/\sqrt{N}$ [14] uses the same counting, and whether random creation reproduces it is open.

## 6 Discussion

**Where this sits.** The construction lies inside causal set theory: links, infinite valency, the Bombelli–Henson–Sorkin dilemma, Number = Volume and growth are all theirs. Our only additions are (i) treating the finite size of the universe as the physical regulator of the nonlocality, (ii) accepting the resulting cosmological frame as an explanation rather than a defect, and (iii) a pedagogical, fully elementary route to these statements. A brief search did not find (i) stated in this form, but it may exist.

**Limitations.**

1. $3+1$ *is not tamed.* $10^{124}$ neighbours per event is finite but hardly local. The $1+1$ result (about 280) is misleadingly mild.
2. *No prediction yet.* The cutoff is unreachable by construction. Effects leaking down to observable rapidities would need dynamics we do not have. A naive guess $e^{-(\eta_{\max}-\eta)}$ gives about $10^{-50}$ at $\eta = 27$.
3. *No dynamics.* Nothing here yields Einstein's equations. Sprinkling stands in for unknown dynamics, and most random orders do not look like spacetime at all (the Kleitman–Rothschild problem [6]).
4. *The fuzziness is classical.* The $n^{1/6}$ fluctuations are statistical. A quantum version is not attempted.
5. *"Size of the universe" is ambiguous.* The Hubble radius, the particle horizon and the event horizon differ by $O(1)$ factors. Thanks to the logarithm this changes $\eta_{\max}$ by $O(1)$, but it must be defined relationally.

**Next steps.**

- **S2, gravity by counting (C6).** Compute the Benincasa–Dowker action [23], the causal-set analogue of the Einstein–Hilbert action built from counts of small intervals, on grown universes. Compare with an expanding GR universe.

- **S3, quantum growth (C7, C8).** Replace probabilities by amplitudes. Test whether this removes the hidden birth-order clock.
- Define $L$, and hence $\eta_{\max}$, relationally in a growing causal set. Test whether random growth reproduces $\Lambda \sim 1/\sqrt{N}$. Repeat in $2+1$ and $3+1$ dimensions.

**Open questions raised in session 2 (to discuss).**

1. Why do the geometric cosmological frame and the matter (CMB) frame coincide in homogeneous GR cosmologies, and what would it take to show this in the discrete model?
2. Can quantum creation operators cure the hidden birth-order clock?
3. How can a newborn event land at an *earlier* cosmic time than events already born? (Birth order versus cosmic time.)
4. Are events the same thing as spacetime points $(x^0, x^1, x^2, x^3)$ in relativity?
5. Roberto's picture: space as discrete "bubbles", each filled (matter) or empty (void). Creating space is a bubble appearing between two others, so light must travel further; destroying space is a bubble bursting, e.g. at the centre of a black hole. How does this spatial picture relate to the spacetime-event picture? Bubble occupancy could be the missing *matter*.

# A Paths explored and dropped

- **A regular crystal of events.** Polygonal light cones and a lattice rest frame (Sec. 2.1).
- **Quantum blur of positions.** Blur in *what*? It presupposes a background. It also cures the zero-measure problem but not the divergence, and an equal superposition over rapidity cannot be normalised.
- **A quantum chain as the source of blur.** Positional zero-point fluctuations grow as $\ln N$ in one dimension. The model is background-dependent, and its claimed analogy with atomic linewidths was overstated.
- **Exactly-one-tick neighbours on lattices with discrete boost symmetry** (Schild [15]). These also give infinitely many neighbours. This confirms the dilemma but adds nothing beyond it.
- **Expansion as a source and gravity as a sink of points** (river model [24]). It needs a preferred family of observers, i.e. an aether, which C2 forbids.
- **A link to Bell non-locality.** Links join only timelike-separated events, while Bell correlations are spacelike. Wrong kind of non-locality.
- $(L/\ell_{\mathrm{P}})^2 \sim 10^{122}$ **and the cosmological constant.** Numerology: every combination of $L$ and $\ell_{\mathrm{P}}$ is a power of $10^{61}$.

# References


[1] T. Jacobson, D. Mattingly, "Gravity with a dynamical preferred frame", Phys. Rev. D 64, 024028 (2001), gr-qc/0007031.

[2] D. Burago, "Periodic metrics"; statement in arXiv:1104.4460.

[3] H. Weyl, *Philosophy of Mathematics and Natural Science* (1949).

[4] M. Q. Vahidi-Asl, J. C. Wierman, in *Random Graphs 87* (1990) and *Random Graphs 89* (1992).

[5] L. Bombelli, J. Lee, D. Meyer, R. D. Sorkin, "Space-time as a causal set", Phys. Rev. Lett. 59, 521 (1987).

[6] S. Surya, "The causal set approach to quantum gravity", Living Rev. Rel. (2019), arXiv:1903.11544.

[7] L. Bombelli, J. Henson, R. D. Sorkin, "Discreteness without symmetry breaking: a theorem", gr-qc/0605006.

[8] R. D. Sorkin, "Does locality fail at intermediate length-scales?", gr-qc/0703099.

[9] S. Coleman, S. L. Glashow, "High-energy tests of Lorentz invariance", Phys. Rev. D 59, 116008 (1999), hep-ph/9812418.

[10] J. D. Barrow, J. Levin, "The twin paradox in compact spaces", Phys. Rev. A 63, 044104 (2001).

[11] G. Brightwell, R. Gregory, "Structure of random discrete spacetime", Phys. Rev. Lett. 66, 260 (1991).

[12] J. Baik, P. Deift, K. Johansson, "On the distribution of the length of the longest increasing subsequence of random permutations", J. Amer. Math. Soc. 12, 1119 (1999).

[13] D. P. Rideout, R. D. Sorkin, "A classical sequential growth dynamics for causal sets", Phys. Rev. D 61, 024002 (2000).

[14] M. Ahmed, S. Dodelson, P. Greene, R. D. Sorkin, "Everpresent $\Lambda$", Phys. Rev. D 69, 103523 (2004).

[15] A. Schild, "Discrete space-time and integral Lorentz transformations", Phys. Rev. 73, 414 (1948).

[16] L. Andersson, G. J. Galloway, R. Howard, "The cosmological time function", Class. Quant. Grav. 15, 309 (1998), gr-qc/9709084.

[17] X. Martín, D. O'Connor, D. P. Rideout, R. D. Sorkin, "On the renormalization transformations induced by cycles of expansion and contraction in causal set cosmology", Phys. Rev. D 63, 084026 (2001).

[18] M. Ahmed, D. Rideout, "Indications of de Sitter spacetime from classical sequential growth dynamics of causal sets", Phys. Rev. D 81, 083528 (2010).

[19] S. Gielen, D. Oriti, L. Sindoni, "Cosmology from group field theory formalism for quantum gravity", arXiv:1303.3576.

[20] D. Oriti, L. Sindoni, E. Wilson-Ewing, "Bouncing cosmologies from quantum gravity condensates", arXiv:1602.08271.

[21] T. Konopka, F. Markopoulou, L. Smolin, "Quantum graphity", hep-th/0611197.

[22] S. Gudder, "Causal set approach to discrete quantum gravity", arXiv:1204.5767.

[23] D. M. T. Benincasa, F. Dowker, "The scalar curvature of a causal set", Phys. Rev. Lett. 104, 181301 (2010).

[24] A. J. S. Hamilton, J. P. Lisle, "The river model of black holes", Am. J. Phys. 76, 519 (2008).