Counting identities in two-dimensional causal sets: spatial distance, depth-drift curvature, and a fixed-size propagator target

Roberto Bada[1, 2, *]

[1] *SapereIA, Barcelona, Spain*
[2] *Institute of Cosmos Sciences, University of Barcelona (ICCUB), Barcelona, Spain*

(Dated: October 29, 2026)

In causal set theory, geometry must be recovered from the causal order and from counting. In two dimensions two exact counting identities are available, and we use their violations as diagnostics. The first is geometric: for two spacelike elements and *any* element c in their common past, the volumes of the causal-overlap regions of Boguñá and Krioukov satisfy $d^2 = 2V_AV_B/V_C$ in flat spacetime. With a Poisson bias correction it gives a spatial-distance estimator that is unbiased for d^2 above two discreteness lengths and more accurate than the equal-depth and past–future constructions on the same pairs. In curved spacetime the identity fails in a way that depends on the depth of c: in de Sitter space $d^2_{\rm est}/d^2 = \tanh^2(H\tau/2)/[2\ln\cosh(H\tau/2)] = 1 - R\tau^2/16 + O(\tau^4)$, which we check against exact volumes in power-law FRW spacetimes and turn into a curvature estimator; on sprinklings the estimator distinguishes flat from curved but is noisy and reads somewhat low. The second identity is combinatorial: the mean of Johnston's chain-sum propagator over intervals with exactly n elements of a random two-dimensional order is $\frac{1}{2}L_n(m^2/2\rho)$, a Laguerre polynomial whose Poisson average is the continuum propagator; it allows a density-free, pair-by-pair check of the propagator on any two-dimensional causal set, although as a manifoldlikeness diagnostic it is not stronger than the ordering fraction. Complementing these, we calibrate the Lorentzian-embedding test of Clough and Evans against size and curvature, show that it misses orders with about ten layers, and add the fraction of null-separated pairs as the missing statistic. Together the diagnostics separate the continuum-like and layered phases of the $e^{-\beta S}$ model at $N = 200$ and 400, including on held-out samples.

I. INTRODUCTION

A causal set is a locally finite partially ordered set: every interval $I(x, y) = \{z : x \prec z \prec y\}$ is finite [1, 2]. Causal set theory proposes that spacetime is such a set, with the continuum emerging at large scales. The number of elements in a region then measures its spacetime volume and the order fixes the light cones; the kinematic programme is to recover all other geometric information from these two ingredients.

Part of this programme is well established. The longest chain between two elements estimates their proper time [3]; interval and chain abundances give the dimension [4, 5] and the curvature [6, 7]; the Benincasa–Dowker operator gives the scalar curvature and the action [8, 9]. Spatial distance is harder, because spacelike elements bound no interval. Boguñá and Krioukov measured it through the overlap of the pasts of two elements as seen from an element c at equal proper time below both [10]; Brightwell and Gregory and Rideout and Wallden used the smallest interval containing both elements [3, 11]; Hu, Meyer and Meyer gave an exact reconstruction in $1 + 1$ dimensions from a past and a future element [12]. Whether a causal set resembles a manifold at all is tested by dimension estimators, interval abundances [7] and Lorentzian embeddings [13, 15, 16]. Matter can be introduced through Johnston's sum over chains [17, 18], whose average over sprinklings reproduces the Klein–Gordon propagator.

Two-dimensional causal sets are a natural laboratory: every 1+1 metric is conformally flat, intervals are rectangles in light-cone coordinates, and a sprinkled diamond is a uniformly random two-dimensional order, whose combinatorics is known [19, 20]; the $e^{-\beta S}$ model of two-dimensional orders also has a well-studied phase structure [19–21]. This paper is built on two exact identities that follow from these facts, and on the use of their violations as diagnostics.

1. A *rectangle identity* expresses the spatial distance of two elements through the overlap volumes seen from any element of their common past (Sec. III). It removes the equal-depth requirement of Ref. [10] and yields an accurate distance estimator. Its violation in curved spacetime, as a function of the depth of c, measures curvature (Sec. IV).

2. A *Laguerre identity* gives the exact mean of Johnston's propagator at fixed interval size (Sec. V). It is a density-free check of the propagator; deviations from it can flag orders that are not continuum-like, although not better than the ordering fraction that is its first term.

To these we add an empirical diagnostic, a calibration of the Lorentzian-embedding test of Clough and Evans with a statistic that closes a blind spot for layered orders (Sec. VI), and apply the diagnostics together to sequential growth and to the two phases of the $e^{-\beta S}$

* roberto@sapereia.com

model (Sec. VII). Section VIII discusses novelty and limitations; other approaches we tested are summarized in Appendix A. Spatial distances in $3+1$ dimensions, where the rectangle identity has no analogue, are treated in a companion paper [22].

II. SETTING

Test causal sets are generated by sprinkling: a Poisson process of density ρ in a region of a given spacetime, with the order induced by the light cones. The coordinates are then discarded and used only for comparison. We write $\ell = \rho^{-1/2}$ for the discreteness length: a region of volume ℓ^2 contains one element on average.

We use light-cone coordinates $u = t - x$, $v = t + x$, in which $dt\,dx = du\,dv/2$ and $x \prec y$ if and only if $u_x < u_y$ and $v_x < v_y$. The interval between $x \prec y$ is the rectangle $[u_x, u_y] \times [v_x, v_y]$, with volume $V = \Delta u\,\Delta v/2 = \tau^2/2$. A sprinkled causal diamond is therefore a uniformly random two-dimensional order, the intersection of two random linear orders [19, 20]. For two spacelike elements, $d^2 = \Delta x^2 - \Delta t^2 = -\Delta u\,\Delta v$.

The proper time from the longest chain has mean $\simeq 2\sqrt{n}$ in an interval of n elements, with Tracy–Widom fluctuations of order $n^{1/6}$ [23]; wherever a volume is available we use the count instead, $\tau^2 = 2n/\rho$. Pairs of elements in one sprinkling share elements and their errors are correlated; quoted uncertainties are therefore computed from the spread between independent sprinklings.

III. THE RECTANGLE IDENTITY AND A DISTANCE ESTIMATOR

Let a, b be spacelike and c an element in their common past. Following Ref. [10], we split the two intervals into

$$A = I(c,a) \setminus I(c,b), \qquad B = I(c,b) \setminus I(c,a), \qquad C = I(c,a) \cap I(c,b). \tag{1}$$

Boguñá and Krioukov choose c at equal proper time below a and b and derive d from the overlap fraction. In $1+1$ dimensions the choice of c is immaterial, because of the following identity (Fig. 1).

Derivation. Place c at the origin and take $u_a < u_b$, $v_a > v_b$ (a and b spacelike). Then $I(c,a) = [0,u_a] \times [0,v_a]$ and $I(c,b) = [0,u_b] \times [0,v_b]$. The rectangle from c to the join (u_b, v_a) splits into four tiles,

$$C = [0,u_a] \times [0,v_b], \quad A = [0,u_a] \times [v_b,v_a], \quad B = [u_a,u_b] \times [0,v_b], \quad D = [u_a,u_b] \times [v_b,v_a]. \tag{2}$$

With $\Delta u = u_b - u_a$ and $\Delta v = v_a - v_b$, the volumes are $V_C = u_a v_b/2$, $V_A = u_a \Delta v/2$, $V_B = \Delta u\, v_b/2$, $V_D = \Delta u\,\Delta v/2$, so that $V_C V_D = V_A V_B$. The tile D has a and b at opposite corners, so $V_D = d^2/2$. Hence, for every c in the common past,




FIG. 1. The rectangle identity in light-cone coordinates (rotated by 45° with respect to the usual picture). The rectangle from c to the join of a and b splits into the overlap C, the strips A and B seen by only one of a, b, and the tile D with a and b at opposite corners. Opposite tiles have equal products, $V_C V_D = V_A V_B$, and $V_D = d^2/2$.

$$d^2 = \frac{2 V_A V_B}{V_C}. \tag{3}$$

The tile D is the interval between the meet and the join of a and b, i.e., the region whose volume defines the $1+1$ spatial distance of Brightwell and Gregory and of Rideout and Wallden [3, 11]. The identity shows that this volume can be read from the past alone, through a single element c. The equal-time choice of Ref. [10] is $V_A = V_B$, in which case their $1+1$ formula reduces to Eq. (3); for general c the search for equal-time elements, which requires the density and the dimension, is unnecessary. The identity relies on the overlap of two $1+1$ intervals being again an interval; in higher dimensions it is not, and Eq. (3) has no direct analogue.

Estimator. On a causal set, V_X is replaced by N_X/ρ, the number of elements in the region. The three regions are disjoint, so N_A, N_B, N_C are independent Poisson variables with means $\lambda_X = \rho V_X$. The product $N_A N_B$ is unbiased for $\lambda_A \lambda_B$, but $1/N_C$ is biased and undefined at $N_C = 0$. Since $E[1/(N+1)] = (1 - e^{-\lambda})/\lambda$ for a Poisson variable,

$$\widehat{d^2} = \frac{2 N_A N_B}{\rho\,(N_C + 1)} \tag{4}$$

is unbiased for d^2 up to a factor $1 - e^{-\lambda_C}$. The relative variance is approximately $1/\lambda_A + 1/\lambda_B + 1/\lambda_C$, so the deepest available c is optimal; we take the element with the largest N_C. The strips A and B are long and thin and contain many elements even when $d \sim \ell$, which is why the estimator works down to the discreteness scale; by contrast, a direct count of the tile D has relative variance $1/\lambda_D = 2\ell^2/d^2$, which is $\frac{1}{2}$ at $d = 2\ell$.

Tests. Table I and Fig. 2 use flat sprinklings of a unit diamond, random spacelike pairs among late elements, and the deepest c for each pair. The median error falls approximately as $N^{-1/2}$ at large d, and remains at 15–25% below 2ℓ, the discreteness scale itself, where $\widehat{d^2}$ is

TABLE I. Median relative error of Eq. (4) with the deepest c; flat $1+1$ sprinklings, 300 pairs among late elements in each of 4 sprinklings per N. (Table II uses pairs in the middle of the diamond, with smaller common pasts, and therefore larger errors.)

N	$d < 2\ell$	2ℓ–10ℓ	$> 10\ell$
1000	23%	6.5%	5.2%
2000	15%	5.0%	3.4%
4000	19%	5.1%	2.5%

TABLE II. Rectangle estimator, Eq. (4), against the equal-depth filter of Ref. [10] and the smallest past–future interval [3, 11], on the same spacelike pairs in the middle of flat diamonds (12 sprinklings × 100 pairs per N). Bias: mean relative error of d^2, in percent, with the standard error from the spread between sprinklings in parentheses. Error: median absolute relative error of d.

		2ℓ–10ℓ		$> 10\ell$	
N	method	bias	error	bias	error
1000	rectangle	+1(5)	0.095	−4(4)	0.082
	equal depth	+10(4)	0.131	+13(3)	0.126
	past–future	+86(3)	0.272	+25(3)	0.095
4000	rectangle	0(2)	0.059	−2(1)	0.039
	equal depth	+6(2)	0.106	+2(1)	0.075
	past–future	+79(5)	0.282	+14(1)	0.049

also biased high [by $(37 \pm 7)\%$ at $N = 4000$ on the pairs of Table II]. Table II compares the estimator with two existing constructions on the same pairs, chosen in the middle of the diamond so that both a common past and a common future exist: the equal-depth filter of Boguñá and Krioukov [their Eq. (34)], and the smallest interval $I(c, f)$ with $c \prec a, b \prec f$ (Brightwell–Gregory, Rideout–Wallden), $d^2 = 2(N_{\min} - 2)/\rho$. Above 2ℓ the rectangle estimator is unbiased for d^2 within the uncertainties, and its median error is the smallest of the three. For d itself it reads 1–3.5% low, as expected from the square root of an unbiased estimate of d^2. The past–future count is biased high, because the smallest interval between elements cannot coincide with the tile D. The equal-depth estimator coincides with Eq. (3) in the continuum; its positive bias on the causal set comes from the overlap fraction, a ratio of small counts used without a Poisson correction. For the baseline tests we add a bottom element that precedes all others; the deepest c is then always that element, selected without reference to its own count, so the selection introduces no bias (in Table I, without such an element, it is one of the lowest elements); the freedom in the choice of c is exercised by the curvature tests of Sec. IV.




FIG. 2. Eq. (4) with the deepest c on flat $1+1$ sprinklings, from the causal order only. (a) Estimate against true distance, $N = 4000$. (b) Median relative error per distance bin; the dotted line marks 2ℓ.

IV. CURVATURE FROM DEPTH DRIFT

In flat spacetime every c in the common past gives the same estimate. In curved spacetime Eq. (3) no longer holds exactly, and the estimate depends on the depth of c. This dependence measures curvature.

de Sitter volumes. Two-dimensional de Sitter space in conformal coordinates,

$$ds^2 = \frac{-d\eta^2 + dx^2}{(H\eta)^2}, \qquad \eta < 0, \tag{5}$$

has the light cones of flat space. The tiles A, B, C, D are the same rectangles in $(u, v) = (\eta - x, \eta + x)$, but volumes carry the conformal weight,

$$dV = \frac{d\eta\, dx}{(H\eta)^2} = \frac{2\, du\, dv}{H^2 (u+v)^2}. \tag{6}$$

Integrating over $[u_1, u_2] \times [v_1, v_2]$, the inner integral gives $1/(u+v_1) - 1/(u+v_2)$ and the outer one a logarithm,

$$V_\square = \frac{2}{H^2} \ln \frac{\eta_L\, \eta_R}{\eta_{\rm bot}\, \eta_{\rm top}}, \tag{7}$$

with $\eta = (u+v)/2$ evaluated at the left, right, bottom and top corners (all $\eta < 0$, so the argument is positive).

Closed form. Place a, b on one slice at $(\eta_0, \mp h)$ and c directly below their midpoint at $(\eta_0 - T, 0)$, and set $H = 1$. By symmetry $V_A = V_B$. Writing $\bar\eta = |\eta_0|$ and $q = \bar\eta + T/2$ (so that $|\eta|$ at the corners of A is $q \mp h/2$, $\bar\eta + h$ and $\bar\eta$), Eq. (7) gives, to first order in h,

$$V_A = 2 \ln \frac{(q - \frac{h}{2})(\bar\eta + h)}{(q + \frac{h}{2})\, \bar\eta} \simeq 2h \Big(\frac{1}{\bar\eta} - \frac{1}{q} \Big) = \frac{hT}{\bar\eta\, q}, \tag{8}$$

and at $h = 0$

$$V_C = 2 \ln \frac{q^2}{(\bar\eta + T)\, \bar\eta} = -2 \ln(1 - x), \qquad x = \frac{T^2}{4q^2}. \tag{9}$$

The true distance is $d \simeq 2h/\bar\eta$, so

$$\frac{d_{\rm est}^2}{d^2} = \frac{2V_A^2/V_C}{4h^2/\bar\eta^2} = \frac{x}{-\ln(1-x)}. \tag{10}$$

The proper time from c to the midpoint of a, b is $\tau = \ln[(\bar{\eta}+T)/\bar{\eta}]$, so that $T/\bar{\eta} = e^\tau - 1$ and $x = [(e^\tau - 1)/(e^\tau+1)]^2 = \tanh^2(\tau/2)$. Restoring H,

$$\frac{d^2_{\rm est}}{d^2} = \frac{\tanh^2(H\tau/2)}{2\ln\cosh(H\tau/2)} = 1 - \frac{(H\tau)^2}{8} + O(\tau^4) \\ = 1 - \frac{R\,\tau^2}{16} + \ldots, \qquad R = 2H^2, \tag{11}$$

exact as $d \to 0$ at any depth, for c directly below the midpoint of a and b. The denominator is $H^2/2$ times the de Sitter volume of an interval of height τ, $V = (4/H^2)\ln\cosh(H\tau/2)$, and the result is independent of η_0, as required by the de Sitter symmetry. We verified Eq. (11) against the exact volumes of Eq. (7) to 10^{-5} for $H\tau = 0.1$–3. A sideways offset of c changes it little, and a finite d adds an approximately depth-independent shift. We call $1 - d^2_{\rm est}/d^2$ the depth drift.

Other spacetimes. Every $1+1$ metric is conformally flat, so the leading term should hold locally in any $1+1$ spacetime. We checked this in FRW spacetimes $ds^2 = a(\eta)^2(-d\eta^2 + dx^2)$ with $a = \eta^p$, where $R = 2(\ln a)''/a^2 = -2p\,\eta^{-2-2p}$ varies in time and the rectangle volumes have closed forms. For $p = 1$, -2 and $\frac{1}{2}$ (negative, positive and negative curvature), $16(1 - d^2_{\rm est}/d^2)/\tau^2$ tends to R at the location of a, b as $\tau \to 0$ (to 1% at $\tau = 0.01$), with the correct sign: for $R < 0$ the ratio exceeds 1. At finite depth it measures a curvature between the values at a, b and at the proper-time midpoint: the drift averages R over the region, with more weight near the top. We have not derived the weighting kernel in closed form.

Test on sprinklings. We sprinkle de Sitter space ($H = 1$, $\rho = 10^4$) and, as a control, flat space in the same coordinates, insert a, b by hand $\approx 14\ell$ apart, and evaluate Eq. (4) for every sprinkled c in their common past whose rectangle lies inside the region. Figure 3 shows the mean ratio per depth bin over 400 sprinklings against the continuum value computed for the same elements c, which includes the effects of finite d and of sideways offsets and therefore lies slightly below Eq. (11). The flat control remains at 1 and de Sitter follows the continuum. Neighboring depth bins share elements and are correlated at the 0.9 level, so a run of deviations in adjacent bins counts as a single fluctuation. In the shallowest bin the tile C contains few elements, the factor $1 - e^{-\lambda C}$ of Eq. (4) is not negligible, and the estimator reads low in both geometries.

A curvature estimator. Using counts only, we fit $\widehat{d^2} = \alpha + \beta\tau^2$ over the elements c of one pair with $H\tau \in [0.2, 0.7]$ and $\tau^2 = (N_{I(c,a)} + N_{I(c,b)})/\rho$, and set $R_{\rm est} = -16\beta/\alpha$. Table III shows the result over 100 sprinklings with one pair each. The value expected from the same fit applied to the continuum depends on whether τ is the true proper time ($R^{\rm cont}_{\rm est} = 1.88$) or is read from interval volumes as the counts do (2.2). In the $p = 1$ FRW spacetime ($\rho = 10^4$, 200 sprinklings, a, b at $\eta_0 = 1.5$), the binned ratio follows the exact continuum within 1σ for $\tau \geq 0.15$, and the fit gives $R_{\rm est} = -1.25 \pm 0.30$ against $R^{\rm cont}_{\rm est} = -0.81$, while R ranges from -0.4 at a, b to -2.8 at the bottom of the window. The estimator therefore measures a depth-weighted average of R, not its value at a, b. All sprinkled values lie within 1.5σ of the continuum fit and distinguish flat from curved, but they lie systematically on the low side, so an offset of order -0.5 cannot be excluded with this statistics; a single pair is not informative, and about 100 pairs are needed for ± 0.4. Other counting routes to curvature use the Benincasa–Dowker operator [8], chain abundances [6] or Ollivier–Ricci curvature [24]; we have not compared their noise with this one at equal density. The present estimator is geometrically transparent: the curvature appears as the failure of an exact flat-space identity, as a function of depth.




FIG. 3. Depth drift of the rectangle estimate: mean over 400 sprinklings ($\rho = 10^4$, $H = 1$) per bin of the proper time τ from c to a, b, with errors from the spread between sprinklings. Solid: continuum ratio from the exact volumes, averaged over the same elements c (finite d and offsets included). Dashed: small-τ term of Eq. (11).

TABLE III. Curvature estimator $R_{\rm est} = -16\beta/\alpha$ over 100 sprinklings, one pair each ($H = 1$). The same fit on the continuum gives 1.88–2.2 for de Sitter (see text). Last column: spread of single-pair values.

ρ	flat ($R = 0$)	de Sitter ($R = 2$)	spread per pair
4000	-0.7 ± 0.6	1.3 ± 0.7	≈ 6–7
10^4	0.3 ± 0.4	1.6 ± 0.4	≈ 4

V. A DENSITY-FREE TARGET FOR JOHNSTON'S PROPAGATOR

The propagator. In Johnston's model a particle hops along the order with amplitude a per hop and b per intermediate stop [17]. With C the causal matrix ($C_{xy} = 1$ if $x \prec y$) and $\Phi = aC$,

$$K = \Phi + b\,\Phi^2 + b^2\Phi^3 + \cdots = \Phi\,(I - b\Phi)^{-1}, \tag{12}$$

with $a = \frac{1}{2}$ and $b = -m^2/\rho$ in $1+1$ dimensions. $K(x,y)$ is a weighted count of chains, $K(x,y) = \frac{1}{2}\sum_k c_k(x,y)\,(-\mu/2)^k$, where c_k is the number of k-element chains strictly inside $I(x,y)$ and $\mu = m^2/\rho$. In a sprinkled interval of volume $V = \tau^2/2$ the expected number of k-chains is $(\rho V)^k/(k!)^2$, so that $\langle K\rangle = \frac{1}{2}\sum_k[-(m\tau/2)^2]^k/(k!)^2 = \frac{1}{2}J_0(m\tau)$, the retarded Klein–Gordon propagator [17].

Fixed interval size. K depends only on the order and on μ. In a uniformly random two-dimensional order the n elements inside an interval again form a uniformly random two-dimensional order, and any k of them form a chain with probability $1/k!$ (the two linear orders must agree on them). Hence $E[c_k \mid n] = \binom{n}{k}/k!$ [25], and

$$E[K \mid n] = \frac{1}{2}\sum_{k=0}^{n}\binom{n}{k}\frac{(-\mu/2)^k}{k!} = \frac{1}{2}L_n\Big(\frac{\mu}{2}\Big) \approx \frac{1}{2}e^{\mu/4}J_0\Big(\sqrt{2\mu(n+\tfrac{1}{2})}\Big), \tag{13}$$

a Laguerre polynomial in the interval size; the Bessel form is the Hilb-type asymptotic for large n at fixed μ [26]. Averaging over Poisson-distributed n with mean $\lambda = \rho\tau^2/2$, using $\sum_n e^{-\lambda}\frac{\lambda^n}{n!}\binom{n}{k} = \frac{\lambda^k}{k!}$, recovers $\frac{1}{2}\sum_k(-\mu\lambda/2)^k/(k!)^2 = \frac{1}{2}J_0(m\tau)$: Johnston's result is the Poisson average of Eq. (13). The fixed-n form requires neither the density nor coordinates: every related pair of a causal set supplies its n and its K, and Eq. (13) is the target. The identity itself is combinatorial: at $\mu = -2$ and summed over permutations it counts all increasing subsequences, $n!\,L_n(-1)$ [27, 28]. A fixed-n test also exists for maximal chains, whose length distribution in a two-dimensional interval of n elements measures dimension and embeddability [29]; Eq. (13) instead sums all chains with the propagator's weights. Path-sum studies of the propagator average over sprinklings [30, 31].

Tests. Figure 4 bins all related pairs of an $N = 400$ order by n and compares $\langle K\rangle$ with Eq. (13) at $\mu = 0.05$. As a statistic we use the rms deviation S_K of K from the target over all related pairs with $n \geq 2$, and compare it with its distribution over 40 random orders ($S_K = 0.0158 \pm 0.0018$). We apply it to 60 final states of $N = 400$ chains of the $e^{-\beta S}$ model (Sec. VII; the 44 states used there and 16 from chains completed later): 28 continuum-like states (30–39 layers), 14 states with 4–11 layers, and 18 with 2–3 layers, which contain too few large intervals for the statistic. Continuum-like states lie within 3 standard deviations of the random orders. States with 9–11 layers deviate by 39–62 standard deviations. In 4- to 7-layer crystals no interval contains more than a few elements, so only small n, where the target is close to $\frac{1}{2}$, is probed: their deviations range from -9 to $+25$ standard deviations, and one 4-layer state lies within 1.2 standard deviations of the random orders. S_K alone therefore does not flag every crystal. Sequentially grown universes ($N = 1500$) deviate strongly, one ($p = 0.02$) reaching $|K| \approx 10$ for individual pairs, whereas the target is bounded by $\frac{1}{2}$; since such orders are not two-dimensional orders, this only confirms that the target is not met by arbitrary orders.




FIG. 4. Johnston's propagator averaged over related pairs with interval size n, against $\frac{1}{2}L_n(\mu/2)$, at $\mu = 0.05$ and $N = 400$. At large n few pairs remain and the bins are noisy. In the 4-layer crystal every interval has fewer than 10 elements, so it contributes a single point near $n = 0$.

Relation to the ordering fraction. Expanding Eq. (13) in μ, the first nontrivial term is c_2, the number of related pairs inside the interval, whose fixed-n mean $\binom{n}{2}/2$ is the ordering fraction behind the Myrheim–Meyer dimension [4, 5]. We therefore compared S_K with the analogous statistic S_f, the rms deviation of $c_2/\binom{n}{2}$ from $\frac{1}{2}$, on the same states. The layered states deviate from random orders by 50–58 standard deviations in S_f, comparable to S_K, and all six 4- to 7-layer crystals by more than 100, including the one that S_K misses. As a diagnostic of manifoldlikeness, the propagator target is therefore not stronger than its first term. Its use lies elsewhere: it is an exact, density-free statement about the propagator itself, against which an implementation of the chain sum can be checked pair by pair on any two-dimensional order, without knowledge of the density.

VI. AN EMBEDDING TEST AND ITS BLIND SPOT

Embedding. Given the squared intervals s_{ij} between M sampled elements, classical multidimensional scaling (MDS) with a Lorentzian Gram matrix reconstructs coordinates. Clough and Evans applied it to causal sets and proposed its eigenvalue spectrum as a test of dimension and manifoldlikeness [13, 14]. With $s_{ij} = (x_i - x_j)^\top \eta\,(x_i - x_j)$, double centering, $G = -\frac{1}{2}JSJ$ with $J = I - \mathbf{1}\mathbf{1}^\top/M$, gives $G = X\eta X^\top$ for the centered coordinates X. A $1+1$ Lorentzian set shows one large negative eigenvalue (time), one large positive eigenvalue (space), and the others near zero; the two eigenvectors give coordinates up to a Lorentz transformation.

We supply MDS with counts only: $s = -2N(I)/\rho$ for related pairs and Eq. (4) with the deepest c for spacelike pairs. On $M = 150$ elements of a flat sprinkling with $N = 2000$, positions are recovered to $0.43 \pm 0.07\,\ell$ across a region $\approx 38\ell$ wide, and the map to the true coordinates

is a Lorentz transformation to within 3–10%. The two large eigenvalues are $\approx \mp 2.2$ and all others are below 0.1, in the coordinate units of the sprinkled region (a unit square in u, v).

Statistics and calibration. As test statistics we use intrinsic numbers only: r_+, the ratio of the second to the first positive eigenvalue; the share of the spectrum in one time and one space direction, $(|\lambda_-| + \lambda_+)/\sum_i |\lambda_i|$; and the *stress*, the rms misfit of the squared intervals by the best $1+1$ Minkowski embedding relative to their rms. The sample consists of the $M = 40$ elements with the largest pasts, after removing elements with no common past with the rest. Figure 5(a) shows the calibration on flat universes (random two-dimensional orders): stress and r_+ settle only for $N \gtrsim 100$ (stress 0.16, 0.10, 0.08 at $N = 100$, 200, 400), so the test is used only above this size. Curvature raises the stress: for de Sitter diamonds with fixed $N = 400$ it stays in the flat band up to $H\tau \approx 1$ and reaches 0.25 at $H\tau = 4$ [Fig. 5(c)]. The stress thus tests whether the order is a *flat* sheet, while Sec. IV measures its curvature.

Blind spot. Applied to Monte Carlo samples of the $e^{-\beta S}$ model at $N = 400$ (Sec. VII), this test passes every continuum-like state and fails orders with 4 and 7 layers, but it does not reliably flag orders with about 10 layers: at $M = 40$ one of seven lies in the flat band and two more just above it, and at $M = 80$ two others have stress 0.03, lower than flat orders. Two effects are responsible. The sample contains only the top one or two layers. And in a layered order most sampled pairs have zero squared interval: linked pairs have $N(I) = 0$, and spacelike pairs often have an empty strip, $N_A = 0$ or $N_B = 0$. MDS fits an all-zero matrix perfectly; a bilayer produces exactly that.

Null-pair fraction. We add the fraction z of sampled pairs with zero squared interval, taken as the larger of its values on the top sample and on a sample spread evenly in height (longest chain below the element) over the upper 70% of the order. On sprinklings z decreases with N (median 0.18, 0.12, 0.09 at $N = 100$, 200, 400) and does not respond to curvature: de Sitter diamonds at $H\tau = 4$ give 0.08–0.11. Layered orders give 0.4–1 [Fig. 5(b)]. We flag an order as layered when z exceeds 1.5 times the flat median at the same N. Orders with only 2–3 layers cannot be processed (too few elements share a common past), and the rule counts them as flagged. The fraction z is related to the interval-abundance tests of Glaser and Surya [7], and layered orders are also separated from manifoldlike ones by graph observables [32]; the point here is that the embedding test requires it, together with its calibration.

VII. APPLICATIONS

Sequential growth. In transitive percolation, the simplest classical sequential growth model [33], elements are born one at a time and each newborn element chooses each existing element as an ancestor with probability p, and the order is the transitive closure. As a sanity check of the diagnostics we grew universes of $N = 100$ and 200 elements with $p = 0.05$, and with the non-covariant variant $p_n = 3/n$. Neither produces a $1+1$ spacetime: the ordering-fraction dimension drifts with scale, and the embedding test fails on every statistic (share of the spectrum in one time and one space direction ≈ 0.33 against 0.65 for flat orders at $N = 100$; stress 0.64–0.78 against 0.10–0.12 at $N = 200$). Their null-pair fraction is not what flags them; grown universes with $N = 1500$ also fail the propagator target (Sec. V). Covariant approaches to cosmology in growth models are discussed in Refs. [34, 35].

TABLE IV. Embedding diagnostics on $e^{-\beta S}$ states at $N = 200$ ($\epsilon = 0.21$). f: ordering fraction; L: number of layers. The last column gives the range of the null-pair fraction z on fresh chains, for $\beta \le 0.3$ (second row) and $\beta \ge 0.6$ with 8–13 layers (third row).

$N = 200$	f	L	r_+	stress	z
random orders	0.50	24	0.07	0.10	0.10–0.15
$\beta = 0.05$	0.51	24	0.10	0.11	0.12–0.13
$\beta = 0.6$	0.59	9	0.39	0.28	0.39–0.48

The two phases of the $e^{-\beta S}$ model. Weighting two-dimensional orders by $e^{-\beta S}$, with S the Benincasa–Dowker action, gives a first-order transition between a continuum-like and a layered phase [19–21]. We sampled it with Metropolis swap moves at nonlocality parameter $\epsilon = 0.21$ (an $O(N^2)$ update, since a swap changes two rows and columns of the causal matrix). Table IV shows the diagnostics at $N = 200$. The states at $\beta \ge 0.6$ are metastable intermediates; the ground state at large β is a bilayer [36].

We then applied the flag to 44 final states of $N = 400$ chains run from random and from layered initial orders at $0.01 \le \beta \le 0.7$. All 22 continuum-like states (30–39 layers) pass; all 9 states with 4–11 layers are flagged ($z = 0.55$–1.0 against a threshold of 0.13); the 13 states with 2–3 layers cannot be processed and are counted as flagged. The fixed-n ordering fraction (Sec. V) separates the same groups; the propagator statistic misses one 4-layer state. Because the flag was designed after these states had been seen, we tested it on held-out samples. On fresh $N = 200$ chains, 12 of 12 continuum-like states pass and 18 of 18 layered states ($\beta = 0.6$–1.5) are flagged. At $N = 800$, with the threshold fixed in advance from flat orders ($z > 0.120$; none of 50 fresh flat orders and 20 de Sitter diamonds is flagged), [r27: results of 28 chains, $\beta \in \{0.01, \ldots, 0.7\} \times 4$ seeds, random initial orders, 2000 sweeps]. These states are test inputs, not equilibrium samples: chains started from layered orders accept $\le 0.1\%$ of the moves. They bracket the transition only loosely: chains from layered starts become continuum-like at $\beta \le 0.05$, and chains from random starts become layered at $\beta \ge 0.45$. The bracket




FIG. 5. Calibration of the embedding test. (a) Flat universes (random two-dimensional orders, 10 per size, $M = 40$): stress and r_+ against the number of elements N. (b) Null-pair fraction z at $N = 400$ ($M = 80$) for flat orders, de Sitter diamonds at $H\tau = 4$, and final states of $e^{-\beta S}$ Monte Carlo chains (continuum-like, 30–39 layers; layered, 4–11 layers). Dashed: the flag threshold, 1.5 times the flat median. The 13 states with 2–3 layers are not shown. (c) Stress of de Sitter diamonds at fixed $N = 400$ (8 per height) against $H\tau$; shaded: the flat band (± 1 standard deviation).

$0.05 < \beta_c(400) < 0.45$ is consistent with the finite-size scaling $\beta_c \approx 1.66/(\epsilon^2 N) = 0.094$ of Ref. [21].

VIII. DISCUSSION

What is new. The overlap regions are those of Ref. [10], the embedding test is that of Ref. [13], the null-pair fraction is related to Ref. [7], and the Laguerre identity is classical combinatorics. To our knowledge the following are new: the rectangle identity for general c, Eq. (3), with the bias-corrected estimator and its comparison with existing constructions; the depth-drift curvature estimator and its closed form, Eq. (11); the application of the Laguerre identity to Johnston's propagator, Eq. (13); and the calibration of the embedding test against size, curvature and layering. Our literature search (October 2026) covered the citing papers of Refs. [10, 13, 17, 18]; the rectangle identity is elementary enough that it may appear in passing in work we did not reach.

Limitations. The exact results are specific to $1+1$ dimensions; the rectangle identity has no direct analogue in higher dimensions, where the overlap of two intervals is not an interval. The curvature estimator is noisy (about 100 pairs for $\pm 20\%$), was tested on de Sitter and one FRW spacetime, measures a depth-weighted average of R, and may carry a small negative offset at the densities used. The embedding thresholds are empirical and must be recalibrated at each N; the $e^{-\beta S}$ states used are test inputs, not equilibrium samples. The Laguerre target is exact only for two-dimensional orders, and as a manifoldlikeness diagnostic it is not stronger than the ordering fraction.

Outlook. A version of the depth drift built on the past-and-future construction of Hu, Meyer and Meyer [12] may be less noisy. A fixed-n propagator target in $3+1$ dimensions would not be exact: Johnston's $3+1$ particle hops along links only [17], so a path requires every skipped sub-interval to be empty, and $E[K \mid n]$ becomes a sum of $(k-1)$-point integrals with factors $(1 - W/V)^{n-k+1}$. Its first term can be evaluated at large n: the expected number of two-hop link paths tends to 3π, which is what Johnston's amplitudes require to reproduce the short-distance limit $m^2/8\pi$ of the continuum mass term [17], so this is a consistency check rather than a new result; a Monte Carlo evaluation approaches it as $3\pi(1 - 1.6/\sqrt{n})$ at fixed n. A $3+1$ target would therefore be a computed table rather than a formula.

DATA AVAILABILITY

The code that produces every number and figure in this paper, and the research notebook in which the work was carried out, are openly available at `https://github.com/presindev/causet-counting`. Figures are produced by `sims/fig_part1.py`; the baseline comparison by `sims/r31_rect_baseline.py`; the null-pair fraction by `sims/r8_detector_spread.py`; the curvature fit and the FRW check by `sims/r9_drift_fit_frw.py`; the $e^{-\beta S}$ states by `sims/n400_crystal.py`; the propagator statistics by `sims/b_johnston_n400.py` and `sims/r32_laguerre_vs_fraction.py`; the $N = 800$ test by `sims/r27_n800_detector.py`.

ACKNOWLEDGMENTS

This work was funded by SapereIA. Simulations were run on the Nyx cluster of the Institute of Cosmos Sciences of the University of Barcelona (ICCUB). An AI system (Claude Opus 5.5, Anthropic) was used for liter-

ature search, derivations, code and drafting, under the author's direction and checks; the procedure is described in Ref. [38]. The author takes full responsibility for the content.

Appendix A: Other approaches tested

- *Fixed-n propagator as a curvature estimator.* In curved $1+1$ spacetime a term $(R\tau^2/48)J_2(m\tau)$ appears in the propagator [37]. In a de Sitter diamond conditioned on n elements the order depends only on $(n, H\tau)$, so a fixed-n curvature test exists. We find that the fraction of related pairs rises by $\approx 0.027\,(H\tau)^2$ for $n = 10$–60, and that the fixed-n propagator is no more sensitive than that fraction alone: a 3σ detection at $n = 30$, $H\tau = 1$ requires 175 intervals in either case. The fixed-n propagator therefore detects curvature only through the known chain-abundance route [6].

- *Embedding test on small orders.* We first applied the test to exhaustive ensembles of small orders ($N \leq 12$). Its statistics do not separate flat from layered orders at these sizes, and only settle for $N \gtrsim 100$ [Fig. 5(a)], so we moved to Monte Carlo samples at $N \geq 200$.

- *Hysteresis in the $e^{-\beta S}$ chains.* At $N = 400$, chains started from random and from layered orders at the same β end in different phases. We first read this as hysteresis of the transition; the chains from layered starts accept $\leq 0.1\%$ of the moves and are effectively frozen, so the difference reflects the sampler rather than the model.

- *Two-dimensional estimator on $2+1$ sprinklings.* Fed into the embedding, Eq. (4) applied to a $2+1$ sprinkling (with the $2+1$ density) produces two large positive eigenvalues. The formula is not valid there, so this is at most a qualitative indication that the spectrum counts spatial dimensions.

[1] L. Bombelli, J. Lee, D. Meyer, and R. D. Sorkin, Spacetime as a causal set, Phys. Rev. Lett. **59**, 521 (1987).
[2] S. Surya, The causal set approach to quantum gravity, Living Rev. Relativ. **22**, 5 (2019), arXiv:1903.11544.
[3] G. Brightwell and R. Gregory, Structure of random discrete spacetime, Phys. Rev. Lett. **66**, 260 (1991).
[4] J. Myrheim, Statistical geometry, CERN Report No. TH-2538, 1978.
[5] D. A. Meyer, The dimension of causal sets, Ph.D. thesis, Massachusetts Institute of Technology, 1988.
[6] M. Roy, D. Sinha, and S. Surya, Discrete geometry of a small causal diamond, Phys. Rev. D **87**, 044046 (2013), arXiv:1212.0631.
[7] L. Glaser and S. Surya, Towards a definition of locality in a manifoldlike causal set, Phys. Rev. D **88**, 124026 (2013), arXiv:1309.3403.
[8] D. M. T. Benincasa and F. Dowker, The scalar curvature of a causal set, Phys. Rev. Lett. **104**, 181301 (2010), arXiv:1001.2725.
[9] F. Dowker and L. Glaser, Causal set d'Alembertians for various dimensions, Classical Quantum Gravity **30**, 195016 (2013), arXiv:1305.2588.
[10] M. Boguñá and D. Krioukov, Measuring spatial distances in causal sets via causal overlaps, Phys. Rev. D **110**, 024008 (2024), arXiv:2401.17376.
[11] D. Rideout and P. Wallden, Spacelike distance from discrete causal order, Classical Quantum Gravity **26**, 155013 (2009), arXiv:0810.1768.
[12] C. A. Hu, D. A. Meyer, and E. J. Q. Meyer, Reconstructing Minkowski geometry from causal separations, J. Math. Phys. **66**, 122502 (2025), arXiv:2601.03280.
[13] J. R. Clough and T. S. Evans, Embedding graphs in Lorentzian spacetime, PLoS One **12**, e0187301 (2017), arXiv:1602.03103.
[14] J. R. Clough, Causal structure in networks, Ph.D. thesis, Imperial College London, 2017, hdl:10044/1/50162.
[15] J. Henson, Constructing an interval of Minkowski space from a causal set, Classical Quantum Gravity **23**, L29 (2006), arXiv:gr-qc/0601069.
[16] S. Johnston, Simpler embeddings of causal sets into Minkowski spacetime, Phys. Rev. D **111**, 106020 (2025), arXiv:2502.09701.
[17] S. Johnston, Particle propagators on discrete spacetime, Classical Quantum Gravity **25**, 202001 (2008), arXiv:0806.3083.
[18] S. Johnston, Feynman propagator for a free scalar field on a causal set, Phys. Rev. Lett. **103**, 180401 (2009), arXiv:0909.0944.
[19] G. Brightwell, J. Henson, and S. Surya, A 2D model of causal set quantum gravity: the emergence of the continuum, Classical Quantum Gravity **25**, 105025 (2008), arXiv:0706.0375.
[20] S. Surya, Evidence for the continuum in 2D causal set quantum gravity, Classical Quantum Gravity **29**, 132001 (2012), arXiv:1110.6244.
[21] L. Glaser, D. O'Connor, and S. Surya, Finite size scaling in 2d causal set quantum gravity, Classical Quantum Gravity **35**, 045006 (2018), arXiv:1706.06432.
[22] R. Bada, Spatial embedding of $3+1$ causal sets: low-count spacelike pairs and a likelihood fit to interval counts (companion paper, 2026), to be submitted.
[23] J. Baik, P. Deift, and K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations, J. Am. Math. Soc. **12**, 1119 (1999).
[24] J. Barton, S. Borza, and J. Röhrig, Ollivier–Ricci curvature for causal sets, arXiv:2606.04910 (2026).
[25] R. G. Pinsky, Law of large numbers for increasing subsequences of random permutations, Random Struct. Algorithms **29**, 277 (2006), arXiv:math/0407353, Eq. (1.1).
[26] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society Colloquium Publications Vol. 23 (AMS, New

York, 1939).

[27] V. Lifschitz and B. Pittel, The number of increasing subsequences of the random permutation, J. Comb. Theory Ser. A **31**, 1 (1981).

[28] OEIS Foundation, The On-Line Encyclopedia of Integer Sequences, sequence A002720, https://oeis.org/A002720.

[29] M. E. Aghili, L. Bombelli, and B. B. Pilgrim, Path length distribution in two-dimensional causal sets, Eur. Phys. J. C **78**, 744 (2018), arXiv:1805.07312.

[30] S. Shuman, Path sums for propagators in causal sets, Phys. Rev. D **109**, 046008 (2024), arXiv:2307.08864.

[31] A. Kastrati and H. Hinrichsen, Numerical evaluation of the causal set propagator in 2D anti-de Sitter spacetime, Phys. Rev. D **113**, 065003 (2026), arXiv:2504.12919.

[32] A. Eichhorn, H. Mack, K. T. Le, and F. Wagner, Charting causal set configuration space with graph observables, arXiv:2605.27514 (2026).

[33] D. P. Rideout and R. D. Sorkin, Classical sequential growth dynamics for causal sets, Phys. Rev. D **61**, 024002 (2000).

[34] X. Martín, D. O'Connor, D. P. Rideout, and R. D. Sorkin, On the renormalization transformations induced by cycles of expansion and contraction in causal set cosmology, Phys. Rev. D **63**, 084026 (2001).

[35] M. Ahmed and D. Rideout, Indications of de Sitter spacetime from classical sequential growth dynamics of causal sets, Phys. Rev. D **81**, 083528 (2010).

[36] W. J. Cunningham and S. Surya, Dimensionally restricted causal set quantum gravity: examples in two and three dimensions, Classical Quantum Gravity **37**, 054002 (2020), arXiv:1908.11647.

[37] Nomaan X, F. Dowker, and S. Surya, Scalar field Green functions on causal sets, Classical Quantum Gravity **34**, 124002 (2017), arXiv:1701.07212.

[38] R. Bada, Two causal-set papers written with an AI research partner: methodology note (2026), https://github.com/presindev/causet-counting.