Spatial embedding of $3+1$ causal sets: low-count spacelike pairs and a likelihood fit to interval counts

Roberto Bada[1, 2, *]

[1] *SapereIA, Barcelona, Spain*
[2] *Institute of Cosmos Sciences, University of Barcelona (ICCUB), Barcelona, Spain*

(Dated: November 1, 2026)

Recovering spatial distances from the causal order alone is a central open problem in the kinematics of causal sets. We study it on the benchmark of Johnston's recent embedding of $3+1$ causal sets into Minkowski spacetime [Phys. Rev. D **111**, 106020 (2025)]. With oracle substitutions we show that almost all of the residual error of that method comes from spacelike pairs whose shared causal regions contain fewer than ten elements; these pairs concentrate at the waist of the causal diamond, and their errors decrease only slowly with density. We therefore fit the spatial coordinates to the interval counts of related pairs with Casasola's Poisson likelihood, starting from a map built from causal-overlap counts alone. On 44 sprinklings with 1000–15000 elements the fit improves on Johnston's embedding in every sprinkling and removes 72–83% of its residual error; it lowers the median error of individual spatial distances by a third and more than halves the upper tail of their distribution. The remaining error lies along a nearly flat valley of the likelihood: maps of very different accuracy differ in log-likelihood by only $O(1)$, so the likelihood carries almost no information that would let an optimizer prefer the more accurate one. In a region without a waist the gain is smaller, because intervals are truncated at its timelike boundary.

I. INTRODUCTION

In causal set theory spacetime is modeled as a locally finite partially ordered set, whose elements are spacetime events and whose order is the causal order [1, 2]. For a causal set obtained by a Poisson sprinkling into a Lorentzian manifold, the number of elements in a region estimates its spacetime volume, and the order encodes the light cones; the kinematic program is to recover the remaining geometry from these two ingredients. Timelike quantities are comparatively accessible: the longest chain between two related elements estimates their proper time [3], and interval counts give dimension and curvature [4–6]. Spatial distances are harder, because two spacelike elements bound no interval.

Several constructions address this. Brightwell and Gregory and, in more detail, Rideout and Wallden defined spatial distance through intervals between common past and future elements [3, 7]. Major, Rideout and Surya represented spatial hypersurfaces by thickened antichains and recovered their topology from the order [10, 11]; Eichhorn, Surya and Versteegen defined distances on such hypersurfaces from the volume of the smallest common future of two elements, minimized over paths with a mesoscale cutoff, and showed that, because no element is exactly null related to another, short distances are overestimated ("discrete asymptotic silence") [12]. Boguñá and Krioukov measured spatial distance through the overlap of the causal pasts of two elements as seen from a common past element [8], and Hu, Meyer and Meyer gave an exact $1+1$ reconstruction [9]. Embeddings of whole causal sets into Minkowski spacetime were constructed by Henson [13], by Clough and Evans through Lorentzian multidimensional scaling [14], and by Johnston, first through factorization of matrices of inner products [15] and recently through a simpler method [16] that is the current benchmark for $3+1$ sprinklings into a causal diamond. In that method the spatial separation of a spacelike pair is the shortest detour through an element related to both.

Because the benchmark is itself flat, it is worth stating why the accuracy of a flat embedding matters. First, coordinates are the input of most further geometric constructions on a causal set: spatial distances on hypersurfaces [12], local charts from which curvature is reconstructed [31], and the comparison of discrete fields with their continuum counterparts at the positions of elements. Second, embedding tests of manifoldlikeness [14, 18] can only reject a causal set when the embedding error on a genuine sprinkling is small compared with the misfit of a non-manifoldlike one, so a less noisy embedding is a sharper test. Third, independently of any particular method, locating the residual error tells us which information the order carries poorly: we find that it is the spacelike pairs with few shared elements, the same pairs whose distances are overestimated in the construction of Ref. [12]. We do not demonstrate a downstream application in this paper; the claims concern accuracy on the benchmark and the source of its error.

In this paper we ask where the remaining error of Johnston's embedding comes from, and how to remove it. Our results are as follows.

1. We reproduce Johnston's benchmark and show by oracle substitution that the residual error is entirely due to spacelike pairs, and almost entirely due to those whose shared causal regions contain fewer than ten elements (Secs. III and IV).

2. With a causal-overlap distance estimator built on

* roberto@sapereia.com

Boguñá and Krioukov's construction, extended to reference elements at unequal depth, we show that these low-count pairs also limit the classical-scaling embedding of inverted distances, and give a scaling argument for why their errors decrease only slowly with density (Sec. IV).

3. We fit the spatial coordinates directly to the interval counts of related pairs, with their Poisson noise. The likelihood is Casasola's [17]. Our contributions are an intrinsic starting map; a controlled comparison with Ref. [16] up to $n = 15000$, in which the fit removes 72–83% of the residual error; and ablations that show which ingredients matter and that the remaining error lies along a nearly flat valley of the likelihood, which gives an optimizer almost no signal toward more accurate maps (Sec. V).

4. We repeat the comparison in a region without a waist at equal density, and identify the timelike boundary as the source of the reduced gain (Sec. VI).

The cost of the methods is compared in Sec. VII, and Sec. VIII discusses novelty, limitations and extensions. Other approaches that we tested are summarized in Appendix A. A companion paper develops exact counting tools in $1+1$ dimensions [18].

II. SETUP AND BENCHMARK

Following Ref. [16], a Poisson number of points with mean n is sprinkled with density ρ into a causal diamond of height 1 in $3+1$ Minkowski spacetime (waist radius $\frac{1}{2}$, volume $\pi/24$), and its bottom and top tips, v_1 and v_n, are added as two further elements. We call each sprinkling a universe. The order is induced by the light cones; the coordinates are then discarded and used only for scoring. We write $\ell = \rho^{-1/4}$ for the discreteness length, so that a region of volume ℓ^4 contains one element on average; at $n = 3000$ the diamond height is $\approx 12\,\ell$. For a related pair $a \prec b$, N_{ab} denotes the number of elements strictly between them; in a flat interval of proper time τ its mean is $\rho\,\pi\tau^4/24$.

A method returns coordinates from the order alone. As in Table V of Ref. [16], the score is the Pearson correlation $\mathcal{C}$ between embedded and true spatial distances over all pairs. We report the residual $1-\mathcal{C}$ and, for a new method, the fraction of Johnston's residual that it removes, $(\mathcal{C}-\mathcal{C}_\mathrm{J})/(1-\mathcal{C}_\mathrm{J})$, always on the same universes. Because a correlation close to one is dominated by distant pairs and insensitive to an overall rescaling, we also report per-pair distance errors and position errors in units of ℓ (Sec. V).

Table I lists the sets of sprinklings, and each result below names the set it comes from. Quoted uncertainties are standard errors over universes.

TABLE I. Sets of sprinklings. Time coordinate: Johnston's (J) or the true one (true). All sets use independent seeds.

set	n (universes)	time	used in
development	500–1500 (10 per n), 400–2000	J/true	Sec. III, check (i)
diagnostic	1500 (3), 3000 (2)	J	Sec. IV
check	1500 (3), 3000 (2)	J	checks (ii), (iv)
large-n	1000, 3000, 6000 (10 each), 10000 (8), 15000 (6)	J	Sec. V, Fig. 1
shape	diamond 1500, 3000; slab 1500, 6000, 12000 (3–4 each)	true	Sec. VI

III. JOHNSTON'S EMBEDDING

Johnston assigns each element a time coordinate from volumes and the proper time of related pairs from interval counts, so that the spatial separation of a related pair is $R = \sqrt{\Delta t^2 - \tau^2}$. For a spacelike pair he takes the shortest detour through an element z related to both,

$$r(x,y) = \min_z \big[R(x,z) + R(z,y)\big], \tag{1}$$

which is an upper bound on the spatial distance by the triangle inequality and is computed as a min-plus matrix product. Classical multidimensional scaling of r then gives three spatial coordinates.

Our implementation reproduces his Tables III and V, provided that the proper time is computed from the elements strictly between the two endpoints: for Table V we obtain $\mathcal{C} = 0.786$, 0.887, 0.923 at $n = 500$, 1000, 1500 (ten sprinklings each), against his 0.789, 0.887, 0.928. Including the endpoints gives $\mathcal{C} = 0.72$ at $n = 1000$, as also noted by Pellegrin [19]. For the correlation of squared Minkowski intervals (his Table IV) we obtain 0.82, 0.90, 0.93, against his 0.958, 0.977, 0.985, also when the true time coordinate is used. None of the candidate statistics we tried reproduces his values: at $n = 1000$ we obtain 0.969 for the squared intervals of related pairs only, 0.932 for the signed square roots of the intervals, 0.916 for the rank correlation, 0.880 for four-dimensional Euclidean distances and 0.856 for spacelike pairs only, against his 0.977. We therefore use the spatial correlation throughout; no conclusion of this paper depends on Table IV.

On the development set, replacing the spacelike values r by the true spatial distances, and keeping every other step, raises $\mathcal{C}$ from 0.903 to 0.9999 at $n = 1000$ (from 0.915 to 0.9999 at $n = 1500$). Replacing instead the related-pair separations by their true values leaves $\mathcal{C}$ essentially unchanged ($0.903 \to 0.906$). The residual error of the method therefore resides in the spacelike pairs. Pellegrin, studying a link-restricted version of the detour, attributes much of its error to noisy related-pair distances of near-collinear pairs [19]; for Eq. (1) itself, correcting the related pairs does not help.

IV. OVERLAP DISTANCES AND LOW-COUNT PAIRS

Estimator. Let a, b be spacelike and c an element in their common past. In flat spacetime the volumes of $I(c,a)$, $I(c,b)$ and of their overlap C depend only on the proper times $\tau_{ca} \geq \tau_{cb}$ and on the spatial distance d. The overlap in units of the smaller interval,

$$w = \frac{V_C}{V_b} = w\left(\frac{\tau_{cb}}{\tau_{ca}}, \frac{d}{\tau_{ca}}\right) \in [0,1], \tag{2}$$

depends on two ratios only. We tabulate it once by Monte Carlo integration ($\tau_{cb}/\tau_{ca} \in [0.05, 1]$, $d/\tau_{ca} \leq 8$, common random numbers) and invert it in d, reading both proper times from interval counts through $V = \pi\tau^4/24$. On the causal set we use $w = (N_C + \frac{1}{2})/(N_b + 1)$, which keeps the inversion finite at $N_C = 0$, and cap the result at the detour of Eq. (1), an upper bound for exact separations. For the benchmark we take $c = v_1$, so that $d(a,b)$ requires only the sizes of the two pasts and of their intersection, and we use the future side ($c = v_n$) when it contains more elements. For all pairs, these counts follow from two matrix products. The same three counts define Müller's past-cone metric [20], without an inversion to a distance.

This is the construction of Boguñá and Krioukov [8], who introduced the overlap of two causal intervals with a common past element as a measure of spatial distance and proposed embedding the resulting distance matrix. They invert the overlap exactly, numerically and in any dimension, for c at equal proper time below a and b [their Eq. (22)], and on a causal set keep only reference elements at equal depth [their Eqs. (32)–(34)]. Our extension is the table for unequal depths, $\tau_{cb}/\tau_{ca} < 1$, which allows a single reference element for all pairs. The Euclidean analogue, the distance between two sensors from the numbers of shared and unshared neighbors, is known in sensor-network localization, both as a deterministic inversion [21] and as a maximum-likelihood estimate with a Cramér–Rao bound [22]. For late elements of large sprinklings ($n = 2\times10^4$–8×10^4), whose shared pasts are large, the estimator has a median relative error of 1–2% and a bias below 1% for $d \geq 4\ell$, and 6–8% below 2ℓ.

Per pair and as an embedding. On the benchmark most pairs are not late, and the counts are much smaller. Table II compares the two estimates over all spacelike pairs, by true separation. The detour is poor for close pairs, where no element lies between them (its +40% bias at separations below 0.1 is the overestimate of short distances described in Ref. [12]), and accurate for distant pairs, where many elements lie near the straight line; the overlap estimate behaves in the opposite way and is biased low by 5–17%. Restricted to pairs with at least 20 elements in the shared region, its median per-pair error is lower than that of the detour (0.112 against 0.132 at $n = 1000$; 0.062 against 0.083 at $n = 3000$). As a diagnostic, we embed the overlap estimates directly by classical scaling: the result is much worse than Johnston's ($1-\mathcal{C} = 0.18$ against 0.049 at $n = 3000$; Fig. 1). Per-pair accuracy does not translate into an accurate embedding.

TABLE II. Spacelike spatial distances by true separation, $n = 1500$ (all spacelike pairs of one universe): median signed bias and median absolute relative error of the overlap estimate and of Johnston's detour, Eq. (1).

true $\lvert\Delta\mathbf{x}\rvert$	pairs	overlap		detour	
		bias	error	bias	error
0–0.1	3 872	−0.05	0.16	+0.40	0.48
0.1–0.2	43 341	−0.16	0.21	−0.08	0.34
0.2–0.35	217 706	−0.17	0.18	−0.12	0.22
0.35–0.5	295 991	−0.14	0.15	−0.07	0.13
0.5–1	230 652	−0.11	0.12	+0.01	0.06

Low-count pairs. Let k be the number of elements in the shared region on the side used by the overlap estimator. Pairs with $k < 10$ are 50% of the spacelike pairs at $n = 1500$ and 31% at $n = 3000$; most of them straddle the waist of the diamond, where both the common past and the common future are small. A second oracle, on the diagnostic set, shows that Johnston's error lies on these pairs: replacing his detour by the true distance only on the $k < 10$ pairs removes 97% ($n = 1500$) and 94% ($n = 3000$) of his residual, whereas the truth on all other spacelike pairs removes 3% and 4% (with the threshold $k < 30$, 97–99% against 2%). The same holds for the overlap estimates on the same universes: with the truth on the $k < 10$ pairs, their direct embedding removes 96% ($n = 1500$) and 88% ($n = 3000$) of Johnston's residual, while the truth on all other pairs leaves it far worse than his. Classical scaling gives the noisy low-count distances the same weight as all others, whichever estimator supplies them.

Why higher density helps slowly. The information on d is carried by the elements in the past of b that are not in the past of a. For small d this region is a shell of thickness $\sim d$ on a cone of height τ, containing $\kappa\rho\, d\, \tau^3$ elements with $\kappa = O(1)$. Its Poisson fluctuation gives, as an order-of-magnitude estimate,

$$\frac{\delta d}{d} \sim \frac{1}{\sqrt{\kappa\rho\, d\, \tau^3}}. \tag{3}$$

At fixed shape all lengths grow as $n^{1/4}$ in units of ℓ, so $d\tau^3 \propto n$ and $\delta d/d \propto n^{-1/2}$ for a typical pair; the measured median errors fall as $n^{-0.54}$ (overlap) and $n^{-0.42}$ (detour) between $n = 1000$ and 3000. The tail of the error distribution, however, is set by geometry: a pair on opposite sides of the waist has a small common past *and* a small common future, so its τ in Eq. (3) is set by its position rather than by the density. Between $n = 6000$ and 15000, the range least affected by small-n effects, the residual of the direct overlap embedding falls approximately as $n^{-0.4}$, against $n^{-0.85}$ for Johnston's (large-n set, Fig. 1); between $n = 3000$ and 6000 it still halves.

This motivates using the counts themselves, with their noise, instead of distances inverted from them.

V. FITTING COORDINATES TO INTERVAL COUNTS

Likelihood. We keep Johnston's time coordinates t_i and treat the $3n$ spatial coordinates $\mathbf{x}_i$ as unknowns. For a related pair $a \prec b$, N_{ab} is Poisson-distributed with mean $\rho\,\pi\tau_{ab}^4/24$, where $\tau_{ab}^2 = (t_b - t_a)^2 - |\mathbf{x}_b - \mathbf{x}_a|^2$. We use it in a Gaussian form for τ^2. With

$$\widehat{\tau^2}_{ab} = \left(\frac{N_{ab} + \frac{1}{2}}{\rho\pi/24}\right)^{1/2}, \qquad s_{ab} = \frac{\widehat{\tau^2}_{ab}}{2\left(N_{ab}+1\right)^{1/2}}, \quad (4)$$

which correspond to a relative error $\delta N/2N$ of τ^2 (the shifts $\frac{1}{2}$ and 1 keep pairs with $N_{ab} = 0$ finite), the log-likelihood is

$$\ln L(\mathbf{x}) = -\frac{1}{2}\sum_{a\prec b}\left(\frac{\tau_{ab}^2(\mathbf{x}) - \widehat{\tau^2}_{ab}}{s_{ab}}\right)^2, \quad (5)$$

summed over all related pairs (whether all are needed is tested below). Spacelike pairs do not enter. Each element is constrained by its $O(n)$ related partners: a pair with many elements between its members constrains it tightly, and a pair with few constrains it loosely. This is the weighting that classical scaling lacks. The Gaussian form is a convenience: the exact Poisson log-likelihood $\sum[N\ln\mu - \mu]$ gives the same maps (check (iv) below), provided it also encodes that the pair is related; otherwise a pair with $N = 0$ is best fitted as spacelike at no cost, and the fit drifts. The Gaussian form, whose target $\widehat{\tau^2}$ is always positive, needs no such term. We maximize Eq. (5) with the Adam optimizer and the analytic gradient (checked against finite differences), for 400 iterations with a step of 1% of the spatial spread, reduced by a factor 0.3 halfway; the tips are fixed at $\mathbf{x} = 0$. Each iteration costs $O(\#\text{related pairs}) = O(n^2)$.

Starting map. The fit requires a start that is roughly correct and is itself derived from the order. We embed the overlap estimates of Sec. IV by weighted stress majorization (SMACOF [23]; accuracy-weighted scaling is standard in sensor localization [24]), with weight $k/(k+100)$ for each spacelike pair and 1 for each related pair, so that low-count distances have little influence. This map alone removes 23–59% of Johnston's residual, less at large n (Table III).

Relation to previous work. Fitting causal-set coordinates to interval counts with a Poisson likelihood was proposed by Casasola [17], who used a pseudo-likelihood $\sum[N\ln\mu - \mu]$ over related pairs near the light cone ($\tau^2 < 6\ell^2$) in a $3+1$ diamond, with Langevin sampling started from the *true* positions displaced by about ℓ; an intrinsic start was left as an open problem, and no comparison with Ref. [16] was made. Maximum-likelihood coordinates are standard in other contexts: multidimensional scaling [25], latent-space network models [26], hyperbolic maps of complex networks, including a geometric first guess refined by likelihood as here [27, 28], and directed graphs in Minkowski space [29]. What we add is the intrinsic start, the large-n test on Johnston's benchmark, and the ablations below. Casasola's preprint has not been peer reviewed; we take from it the form of the likelihood and the near-cone set of pairs, and reproduce both here.

Results. Table III and Fig. 1 show the fit on the large-n set, paired with Johnston's embedding on the same universes. The fit improves on his embedding in each of the 44 universes. It removes 72% of his residual at $n = 1000$, 74% at $n = 3000$, 80% at $n = 6000$, and 79–83% at $n = 10000$ and 15000; across the ten universes of each size up to 6000 the removed fraction ranges from 61 to 85% (standard deviation 7, 6 and 3 percentage points at $n = 1000$, 3000, 6000). The removed fraction rises at most slowly with n, so the gap does not close. At $n = 15000$ the spatial correlation is 0.9980 against 0.9886. The oracle value with exact spacelike distances, 0.9999, remains higher.

Per-pair errors and causal fidelity. In units of the discreteness length, the median absolute error of the embedded spatial distances is 0.33, 0.23, 0.18 for Johnston's embedding and 0.20, 0.14, 0.12 for the fit at $n = 1000$, 3000, 6000 (ten universes each); the 90th percentiles are 1.14, 0.76, 0.59 against 0.46, 0.33, 0.27, and the 99th percentiles 3.1, 2.9, 2.4 against 1.4, 0.7, 0.5. Because spacelike pairs do not enter Eq. (5), we also checked how well the fitted coordinates reproduce the causal order: the fit misclassifies 1.24%, 0.68% and 0.48% of all pairs (related as spacelike or vice versa), against 3.3%, 1.7% and 1.1% for Johnston's embedding.

Where the error remains. Figure 2 shows the rms position error per element, after removing the best orthogonal map, against the number of related partners of the element and against its time. In both embeddings the error is concentrated on the tenth of the elements with the fewest related partners and, in time, at the waist ($t \approx \frac{1}{2}$); the fit lowers it at every decile and every time, but these elements remain the largest contribution. The rms position error therefore improves less than the median position error. On the ten universes of Fig. 2 the rms falls from 0.66 to $0.57\,\ell$ at $n = 3000$ and from 0.55 to $0.42\,\ell$ at $n = 6000$, while the median falls from 0.27 to $0.18\,\ell$ and from 0.21 to $0.15\,\ell$. The tenth of the elements with the fewest related partners carries an rms of $1.69\,\ell$ (1.43–2.14 over the universes) at $n = 3000$ and $1.24\,\ell$ (0.96–1.51) at $n = 6000$ in the fit, against 1.83 and $1.57\,\ell$ in Johnston's embedding; the other nine tenths carry 0.20 and $0.17\,\ell$, against 0.32 and $0.24\,\ell$. Neither a global rescaling nor an affine map removes this part.

Is the residual set by the optimizer or by the likelihood? Adam is not fully converged after 400 iterations: continuing with L-BFGS raises the log-likelihood by 45 on average at $n = 3000$ (up to 98) and by 176 at $n = 6000$ (up

to 410), but changes $\mathcal{C}$ by only $+1\times10^{-4}$ and $+2\times10^{-4}$. Started from the intrinsic map, from Johnston's map, or from the intrinsic map with every coordinate displaced by $0.5\,\ell$, and run to convergence, the fit reaches the same log-likelihood within a few units out of 4.7×10^5 terms at $n=3000$ (within 10 out of 1.9×10^6 at $n=6000$), and the same $\mathcal{C}$ within the spread between universes (Johnston's start: $+0.0007\pm0.0005$ at $n=3000$, -0.0003 ± 0.0005 at $n=6000$; choosing the start with the highest likelihood: 75.4% of Johnston's residual removed instead of 74.5% at $n=3000$, 81.0% instead of 80.6% at $n=6000$).

Started from the true positions (with Johnston's time coordinate), the fit stays much closer to the truth: it removes 92% of Johnston's residual at $n=3000$ and 93% at $n=6000$. Its log-likelihood differs from that of the converged intrinsic fit by only $O(1)$: it is higher in 4 of 10 universes at $n=1000$ (all differences below 0.8 in magnitude), in 9 of 10 at $n=3000$ (by 0.9 on average, at most 5.4) and in 10 of 10 at $n=6000$ (2.3, at most 6.5). These differences are of the same order as what further iterations add to the intrinsic fit (below), and the two fits were not continued equally, so they do not tell which of the two maps the likelihood prefers. To see how the two maps are connected, we aligned them by the best orthogonal map about the time axis, under which Eq. (5) is invariant; they then differ by $0.63\,\ell$ rms at $n=3000$ and $0.48\,\ell$ at $n=6000$. On the straight line between them the log-likelihood drops by 2.7×10^3–1.3×10^4 at the midpoint. This dip is expected even in the absence of a barrier: for each pair, the norm of the average of two separation vectors is at most the average of their norms (triangle inequality), so at the straight midpoint every related pair is more timelike than the average of the two maps unless its two separation vectors are parallel. Started from the points a quarter, half and three quarters of the way along the line and run to convergence, L-BFGS reaches, in every universe, a map whose accuracy lies between those of the two ends and increases along the path: it removes 78%, 86% and 92% of the residual at $n=3000$ (83%, 89% and 93% at $n=6000$), against 75% and 92% (81% and 93%) at the ends. Each of these maps has a log-likelihood within $O(1)$ of both ends (-3.8 to $+0.6$ from the truth-started fit and -0.9 to $+5.3$ from the intrinsic fit at $n=3000$; -1.1 to $+0.7$ and $+0.25$ to $+6.7$ at $n=6000$). The relaxed midpoint lies only $0.15\,\ell$ rms from the straight midpoint at $n=3000$ ($0.11\,\ell$ at $n=6000$), yet recovers 5.0×10^3 (7.3×10^3) in log-likelihood, as expected from a second-order shortening effect and not from a barrier. The two maps are thus joined by a curved valley along which the log-likelihood changes by $O(1)$ while the removed fraction goes from 75% to 92% (81% to 93% at $n=6000$); no barrier was found along it, although the segments between the relaxed points were not scanned. Continuing L-BFGS from the converged intrinsic fit for 2000 iterations without a stopping tolerance raises the log-likelihood by 0.5 at $n=3000$ and by 2.4 at $n=6000$, of the same order as the difference between the two maps, and leaves $\mathcal{C}$ unchanged (0.98855 → 0.98853 and 0.99544 → 0.99543). Maps of very different accuracy thus differ in log-likelihood by only $O(1)$, so the likelihood carries almost no information that would let an optimizer prefer the more accurate one.

TABLE III. Likelihood fit, Eq. (5), against Johnston's embedding on the large-n set, paired on the same universes. Gain: $\mathcal{C}-\mathcal{C}_\mathrm{J}$ with the standard error of the paired differences in parentheses. Last three columns: fraction of Johnston's residual removed by the fit (mean, and range over universes) and by its starting map. Every row uses the fixed schedule of 400 Adam iterations, which is not fully converged (Sec. V: continuing to convergence changes $\mathcal{C}$ by at most 2×10^{-4}).

n	univ.	$\mathcal{C}_\mathrm{J}$	$\mathcal{C}$ (fit)	gain	fit	range	start
1000	10	0.8855	0.9678	0.082(14)	72%	61–81%	59%
3000	10	0.9551	0.9884	0.0334(19)	74%	64–81%	43%
6000	10	0.9764	0.9953	0.0189(7)	80%	76–85%	33%
10000	8	0.9846	0.9968	0.0122(3)	79%		23%
15000	6	0.9886	0.9980	0.0094(2)	83%		26%

Which pairs are needed? Casasola restricted the likelihood to pairs near the light cone [17]. We ran his exact Poisson form, with the soft relation term of check (iv), once on all related pairs and once on the pairs with $\widehat{\tau^2}<6\,\ell^2$ only (64%, 47% and 37% of related pairs at $n=1000$, 3000, 6000), from the same intrinsic start. The near-cone version is slightly better: $\Delta\mathcal{C}=+0.0004$ at $n=3000$, in ten of ten universes, and $+0.0003$ at $n=6000$, again in ten of ten; the Gaussian form restricted to the same pairs gives $+0.0002$. Distant related pairs therefore add no information that the fit uses, and using all related pairs is not a contribution of this paper. The near-cone set also lowers the cost per iteration by a factor of two to three. For applications we therefore recommend the exact Poisson form on the near-cone pairs. With the same 400-iteration Adam schedule as Table III, on the large-n set it removes 72%, 76% and 82% of Johnston's residual at $n=1000$, 3000, 6000; paired with the all-pairs Gaussian fit of Table III it gains $+0.0008$ and $+0.0006$ in $\mathcal{C}$ at $n=3000$ and 6000, in ten of ten universes at both sizes ($+0.0003$, seven of ten, at $n=1000$). Its cost is estimated in Sec. VII. Table III keeps the all-pairs Gaussian form, whose settings were fixed before these comparisons.

Time coordinate. The fit keeps Johnston's time coordinate. Fitting the time coordinates jointly with the spatial ones (the tips fixed, same likelihood, Adam and L-BFGS) changes $\mathcal{C}$ by $+0.0004\pm0.0001$ at $n=3000$ and $+0.0002\pm0.0001$ at $n=6000$ and does not reduce the time error (0.074 against $0.076\,\ell$ rms at $n=3000$, 0.062 against $0.066\,\ell$ at $n=6000$): Johnston's time coordinate is already accurate to a tenth of a discreteness length, and the related-pair counts do not improve it.

Checks. (i) *Consistency.* Started from the true positions (with true times), the fit departs from the truth only by finite-n noise and approaches it as n grows:




FIG. 1. Residual $1-\mathcal{C}$ of the spatial-distance correlation on Johnston's benchmark (large-n set; $n = 10^3$–1.5×10^4 with 10, 10, 10, 8 and 6 universes; the overlap-distance curve uses a subset of 1, 2, 4, 8, 6 of these universes): his embedding [16], the direct embedding of overlap distances (Sec. IV), the weighted starting map, and the likelihood fit, Eq. (5). Error bars: standard error over universes.

$\mathcal{C} = 0.966$, 0.991, 0.9965 at $n = 400$, 1000, 2000. (ii) *Start independence*, shown above on the large-n set. (iii) *No tuning on the test universes.* The large-n runs used settings fixed beforehand on other universes, including the start's weight scale 100. (iv) *Form of the likelihood and convergence.* On the check sets we compared the Gaussian form with the exact Poisson log-likelihood plus a soft relation term $\sum \ln \mathrm{sig}(\tau^2/s_0)$, with sig the logistic function and $s_0 = 0.1\,\ell^2$ (cf. the soft walls of Ref. [17]), each at 400 and 1000 iterations. All four variants remove 76–77% ($n = 1500$) and 74–75% ($n = 3000$) of Johnston's residual, with paired differences of at most 3×10^{-4} in $\mathcal{C}$. (These fractions differ from those of Table III because the universes differ.) Started from the truth, the exact form remains slightly closer ($\mathcal{C} = 0.973$, 0.993, 0.997 at $n = 400$, 1000, 2000). The log-likelihood still increases slowly after 400 iterations, but the map no longer changes.

VI. A REGION WITHOUT A WAIST

Johnston's error concentrates at the waist of the diamond. We therefore repeat the comparison in a cylinder ("slab") $t \in [0,1]$, $|\mathbf{x}| \le \frac{1}{2}$, without tips. There his embedding is already accurate ($\mathcal{C} = 0.985$ at $n = 1500$, against 0.926 in the diamond). The likelihood needs no reference element and applies unchanged, with no element fixed (we remove the mean of the gradient). To isolate the spatial step, all methods in this section use the true time coordinate. Without tips, 0.4% of spacelike pairs at $n = 1500$ and 0.1% at $n = 6000$ have no element related to both, so Johnston's method is undefined for them; for these pairs his detours fall back to four legs. Every other pair keeps its two-leg value, so the fallback only completes his method. The starting map uses, for each pair, one of 64 reference elements near the lower and upper boundaries.

We compare the shapes at equal density: slab $n = 6000$ and 12000 against diamond $n = 1500$ and 3000, four universes each. The fit removes $58\pm9\%$ and $65\pm6\%$ of Johnston's residual in the slab, against $75\pm5\%$ and $80\pm6\%$ in the diamond through the same pipeline. The difference arises at the timelike boundary. A diamond contains every interval between its elements (it is causally convex); a slab does not, and an interval that extends beyond the boundary contains too few elements, so that its pair appears closer to null than it is, as Pellegrin observed for slab-shaped subsets of a diamond [19]. In the slab, 45% of related pairs lose part of their interval and 14% lose more than a fifth, at every n. Dividing each count by the true fraction of its interval inside the slab (an oracle used for diagnosis only) raises the slab result to 80% and 86% (means over the same four universes), level with the diamond within the uncertainties quoted above: the boundary is sufficient to account for the difference. Its cost, about 20 percentage points, does not decrease with n, because the fraction of truncated intervals is fixed by the shape. On interior pairs (both elements in the top quartile of $\min(|\mathrm{past}|, |\mathrm{future}|)$) the fit does at least as well in the slab as in the diamond ($80\pm12\%$ and $86\pm2\%$, against $69\pm10\%$ and $71\pm14\%$).

Two caveats apply. First, these are fractions of a small residual: in the slab Johnston's $1-\mathcal{C}$ is 0.0069 and 0.0047, about ten times smaller than in the diamond. Second, the gain requires a minimum density: in the slab it drops to $43\pm10\%$ at $n = 1500$ (three universes), and in a single universe at $n = 600$ the fit is worse than his embedding.

VII. COMPUTATIONAL COST

Johnston's min-plus step costs $O(n^3)$ operations; the overlap counts require two dense matrix products and a table look-up; the likelihood costs $O(n^2)$ per iteration at a fixed 400 iterations. We timed both pipelines on one universe per size, on an exclusive node with four BLAS threads (the min-plus step is single-threaded). Johnston's pipeline (matrix product, min-plus step, eigendecomposition) took 44, 525, 2002 and 7691 s at $n = 3000$, 6000, 10000, 15000; the likelihood pipeline (counting step, starting map, set-up, 400 iterations) took 307, 1328, 3647 and 8239 s. Over $n = 6000$–15000 the wall times scale as $n^{2.92}$ and $n^{1.99}$. The likelihood pipeline is slower at every size we ran, by a factor that falls from 7.0 to 1.07; extrapolating the two power laws, the times would cross near $n \approx 16000$.

These timings are for the all-pairs Gaussian form of Table III. The variant we recommend (Sec. V), the exact Poisson form on near-cone pairs, is cheaper per iteration,




FIG. 2. Where the error remains: rms position error per element (after the best orthogonal map), in units of ℓ, for Johnston's embedding and the likelihood fit (converged), large-n set at $n = 3000$ and 6000 (ten universes each). Left: by decile of the element's number of related partners. Right: by true time.

because the share of near-cone pairs falls with n, roughly as $n^{-0.3}$ (0.64, 0.47, 0.37, 0.30, 0.27 at $n = 1000$, 3000, 6000, 10000, 15000), while the exact form costs about 1.7 times the Gaussian form per pair. We did not rerun it on the timing node; scaling the measured time of the 400 iterations of the timing runs by the ratio of the per-iteration costs measured on a laptop, and keeping the other steps, gives an estimated 250, 860, 1900 and 3900 s at the four sizes, against 44, 525, 2002 and 7691 s for Johnston's pipeline: still slower up to $n = 6000$, comparable at $n = 10000$, and about twice as fast at $n = 15000$. The claim of this paper concerns accuracy, not speed.

VIII. DISCUSSION

What is new. The overlap construction is that of Boguñá and Krioukov [8], the time coordinate and the benchmark are Johnston's [16], and the likelihood of related-pair counts is Casasola's [17]; maximum-likelihood coordinates are standard in other fields [25–29]. To our knowledge the following are new: the tabulated overlap inversion for unequal depths; the reproduction and oracle analysis of Ref. [16], with the localization of its error on low-count spacelike pairs; the scaling argument for why inverted distances improve slowly; the intrinsic starting map; the large-n comparison, with the analysis of a second region; and the ablations, which show that pairs near the light cone carry the information and that the remaining error lies along a nearly flat valley of the likelihood, in which maps of very different accuracy differ in log-likelihood by only $O(1)$. Our literature search (October 2026) covered the citing papers of Refs. [8, 14, 16, 17] and sources beyond arXiv and INSPIRE; at the time of writing Ref. [16] has one citing paper on INSPIRE, which reconstructs curvature from counts with local charts rather than distances [31].

Limitations. The likelihood assumes a flat interval volume; in curved spacetime volumes acquire corrections of order $R\tau^2$, which would bias the fit increasingly with interval size (the $1 + 1$ analogue is the depth drift of the companion paper [18]). The time coordinate is Johnston's; refitting it changes little. The benchmark is a flat diamond, with ten universes per size up to $n = 6000$ and six to eight above; in the slab the boundary costs about 20 percentage points at every size, and the gain requires a minimum density.

Outlook. The oracle value ($\mathcal{C} = 0.9999$ against 0.998 at $n = 15000$) leaves room for improvement. The remaining error lies along a nearly flat valley of the likelihood, in which maps of very different accuracy differ in log-likelihood by only $O(1)$, and longer optimization did not move the fit along it. In practice the error is therefore more likely to be reduced by additional information than by further optimization of Eq. (5). The spacelike overlap counts are the obvious candidate, but adding them naively made the fit worse (Appendix A). Other next steps are a model of the boundary of regions that are not causally convex (in $1+1$ dimensions timelike boundaries can be identified from the order alone [30], which might supply the fractions that our boundary correction took from the truth) and a curved interval volume, perhaps via the normal-coordinate construction of Ref. [31], which would turn the fit into a curvature estimator.

DATA AVAILABILITY

The code that produces every number in this paper, and the research notebook in which the

work was carried out, are openly available at https://github.com/presindev/causet-counting, and the version used for this paper is archived at doi:10.5281/zenodo.23856330. The overlap table is in `sims/r11_4d_distance.py`; the reproduction of Ref. [16] and the first oracle in `sims/r12_johnston_compare.py`; the low-count oracles in `sims/r18_waist_oracle.py` and `sims/r30_johnston_lowk_oracle.py`; the starting map in `sims/r19_cluster_wfit.py`; the likelihood in `sims/r23_likelihood.py` and its large-n runs in `sims/r24_cluster_lik.py`; the per-pair errors in `sims/r29_perpair_error.py`, the per-element analysis in `sims/r33_time_vs_position.py` and the likelihood comparison and causal fidelity in `sims/r34_lnl_causal_fidelity.py`; the slab in `sims/r22_slab.py` and `sims/r25_slab_lik.py`; the large-n runs (ten universes per size up to 6000, ablations, optimizer and joint-fit checks) in `sims/r35_referee1_3p1.py`, analyzed by `sims/r35_analyse.py`, which also draws Fig. 2; the likelihood path and valley tests, the per-element error numbers and the cost of the near-cone variant in `sims/r39_3p1_round2.py`, summarized by `sims/r39_summary.py`; and Fig. 1 by `sims/fig_3p1.py --prd`.

ACKNOWLEDGMENTS

This work was funded by SapereIA. Simulations were run on the Nyx cluster of the Institute of Cosmos Sciences of the University of Barcelona (ICCUB). An AI system (Claude Opus 5.5, Anthropic) was used for literature search, derivations, code and drafting, under the author's direction and checks; the procedure is described in Ref. [32]. The author takes full responsibility for the content.

Appendix A: Other approaches tested

Before the likelihood fit we tried to make inverted distances competitive, alone or combined with Johnston's detours. Each item was tested on the same universes as the method it is compared with; fractions refer to Johnston's residual.

- *Hard switch*: the overlap estimate when it places the pair closer than $L = 0.25$, Johnston's detour otherwise. With L chosen on other universes, it improved on his embedding in every universe from $n = 500$ to 15000, removing a stable 17–20% of his residual. In the slab it gains nothing, because its gain is located at the waist.
- *Weighted map as a final result*: its share fades with n (Table III) and depends on the weight scale K_0 in $k/(k + K_0)$ (23–36% at $n = 10^4$–1.5×10^4 for $K_0 = 100$–300). We retain it only as the starting map.
- *Binomial term for spacelike overlap counts*, $N_C \,|\, N_b \sim \mathrm{Bin}(N_b, w)$ with w from the table of Sec. IV, added to Eq. (5). Alone it removes 14–25%; added to the fit it lowers the gain to 58–64% at every n of the large-n set. We have not isolated why (the bias of Table II, ratios of noisy counts, and pairs treated as independent are candidates).
- *Map from counts only* (SMACOF with distant pairs at weight 0.1, no detours): matched Johnston's embedding up to $n = 3000$, worse at larger n (-14% to -31%). Dropping the distant pairs instead of down-weighting them: $\mathcal{C}$ lower by 0.012 at $n = 3000$.
- *Laplacian eigenmaps*, suggested in Ref. [8]: limited to $\mathcal{C} \approx 0.93$ even with true distances, because the eigenfunctions of a ball are not linear coordinates.
- *Local trilateration charts* following Ref. [31]: exact on true intervals, but 30–42% per-pair error from counts at $n \leq 3000$ (ill-conditioned Gram matrices).
- *Detour on the low-count pairs only*, overlap estimates elsewhere: no better than Johnston's embedding.
- *Averaging* the overlap estimate with the detour: worse than the detour alone ($\Delta\mathcal{C} = -0.07$ to -0.006).
- *Several reference elements* instead of v_1, averaged: no gain, since deep references see the same region. *Both sides* (past and future references) combined with Poisson weights: worse than taking the side with more elements.
- *Chained detours* through related edges and short overlap estimates: much worse ($\Delta\mathcal{C} = -0.06$ to -0.29); the short legs are biased low and the minimum over many routes selects the lowest.
- *Cheaper detours* through random intermediate elements: at or below Johnston's embedding (-0.008 to -0.033 at $n = 3000$). Through intermediate elements chosen near the midpoint of each distant pair on a rough first map: about two thirds of the switch's gain at $n = 3000$ at a quarter of the cost, but still cubic.

[1] L. Bombelli, J. Lee, D. Meyer, and R. D. Sorkin, Spacetime as a causal set, Phys. Rev. Lett. **59**, 521 (1987).

[2] S. Surya, The causal set approach to quantum gravity,

Living Rev. Relativ. **22**, 5 (2019), arXiv:1903.11544.
[3] G. Brightwell and R. Gregory, Structure of random discrete spacetime, Phys. Rev. Lett. **66**, 260 (1991).
[4] L. Glaser and S. Surya, Towards a definition of locality in a manifoldlike causal set, Phys. Rev. D **88**, 124026 (2013), arXiv:1309.3403.
[5] M. Roy, D. Sinha, and S. Surya, Discrete geometry of a small causal diamond, Phys. Rev. D **87**, 044046 (2013), arXiv:1212.0631.
[6] D. M. T. Benincasa and F. Dowker, The scalar curvature of a causal set, Phys. Rev. Lett. **104**, 181301 (2010), arXiv:1001.2725.
[7] D. Rideout and P. Wallden, Spacelike distance from discrete causal order, Classical Quantum Gravity **26**, 155013 (2009), arXiv:0810.1768.
[8] M. Boguñá and D. Krioukov, Measuring spatial distances in causal sets via causal overlaps, Phys. Rev. D **110**, 024008 (2024), arXiv:2401.17376.
[9] C. A. Hu, D. A. Meyer, and E. J. Q. Meyer, Reconstructing Minkowski geometry from causal separations, J. Math. Phys. **66**, 122502 (2025), arXiv:2601.03280.
[10] S. Major, D. Rideout, and S. Surya, Spatial hypersurfaces in causal set cosmology, Classical Quantum Gravity **23**, 4743 (2006), arXiv:gr-qc/0506133.
[11] S. Major, D. Rideout, and S. Surya, On recovering continuum topology from a causal set, J. Math. Phys. **48**, 032501 (2007), arXiv:gr-qc/0604124.
[12] A. Eichhorn, S. Surya, and F. Versteegen, Induced spatial geometry from causal structure, Classical Quantum Gravity **36**, 105005 (2019), arXiv:1809.06192.
[13] J. Henson, Constructing an interval of Minkowski space from a causal set, Classical Quantum Gravity **23**, L29 (2006), arXiv:gr-qc/0601069.
[14] J. R. Clough and T. S. Evans, Embedding graphs in Lorentzian spacetime, PLoS One **12**, e0187301 (2017), arXiv:1602.03103.
[15] S. Johnston, Embedding causal sets into Minkowski spacetime, Classical Quantum Gravity **39**, 095006 (2022), arXiv:2111.09331.
[16] S. Johnston, Simpler embeddings of causal sets into Minkowski spacetime, Phys. Rev. D **111**, 106020 (2025), arXiv:2502.09701.
[17] G. Casasola, Causal sets that locate and grow themselves: sub-discreteness self-localization and order-intrinsic sequential growth of 3+1 Minkowski orders, Research Square preprint (2026, not peer reviewed), doi:10.21203/rs.3.rs-11223018/v1.
[18] R. Bada, Spatial distance and curvature from an exact counting identity in two-dimensional causal sets (companion paper, 2026), to be submitted.
[19] G. Pellegrin, Link-restricted path filling for causal set embeddings, Zenodo preprint (2026, not peer reviewed), doi:10.5281/zenodo.21813653.
[20] O. Müller, On the Hauptvermutung of causal set theory, arXiv:2503.01719 (2025).
[21] C. Buschmann, D. Pfisterer, and S. Fischer, Estimating distances using neighborhood intersection, in *Proceedings of the IEEE Conference on Emerging Technologies and Factory Automation (ETFA)* (IEEE, 2006).
[22] B. Huang, C. Yu, B. D. O. Anderson, and G. Mao, Estimating distances via connectivity in wireless sensor networks, Wirel. Commun. Mob. Comput. **14**, 541 (2014).
[23] J. de Leeuw and P. Mair, Multidimensional scaling using majorization: SMACOF in R, J. Stat. Softw. **31**(3), 1 (2009).
[24] J. A. Costa, N. Patwari, and A. O. Hero, Distributed weighted-multidimensional scaling for node localization in sensor networks, ACM Trans. Sens. Netw. **2**, 39 (2006).
[25] J. O. Ramsay, Maximum likelihood estimation in multidimensional scaling, Psychometrika **42**, 241 (1977).
[26] P. D. Hoff, A. E. Raftery, and M. S. Handcock, Latent space approaches to social network analysis, J. Am. Stat. Assoc. **97**, 1090 (2002).
[27] M. Boguñá, F. Papadopoulos, and D. Krioukov, Sustaining the Internet with hyperbolic mapping, Nat. Commun. **1**, 62 (2010), arXiv:1009.0267.
[28] G. García-Pérez, A. Allard, M. Á. Serrano, and M. Boguñá, Mercator: uncovering faithful hyperbolic embeddings of complex networks, New J. Phys. **21**, 123033 (2019), arXiv:1904.10814.
[29] A. Sim, M. Wiatrak, A. Brayne, P. Creed, and S. Paliwal, Directed graph embeddings in pseudo-Riemannian manifolds, in *Proceedings of ICML 2021*, arXiv:2106.08678.
[30] W. J. Cunningham, Inference of boundaries in causal sets, Classical Quantum Gravity **35**, 094002 (2018), arXiv:1710.09705.
[31] N. Madsen, Reconstruction of the Riemann tensor from causal order and element counts, arXiv:2610.36695 (2026).
[32] R. Bada, Two causal-set papers written with an AI research partner: methodology note (2026), released with the code at `https://github.com/presindev/causet-counting`.