

Abscense of diffusion in certain random lattices

testing out titles and stuff

P.W. Anderson

1957-10-10

Abstract

This paper presents a simple model for such processes as spin diffusion or conduction in the “impurity band.” These processes involve transport in a lattice which is in some sense random, and in them diffusion is expected to take place via quantum jumps between localized sites. In this simple model the essential randomness is introduced by requiring the energy to vary randomly from site to site. It is shown that at low enough densities no diffusion at all can take place, and the criteria for transport to occur are given.

Bell Telephone Laboratories

Contents

1	Introduction	1
2	Summary of the reasoning	1
	References	2

1. Introduction

A number of physical phenomena seem to involve quantum-mechanical motion, without any particular thermal activation, among sites at which the mobile entities (spins or electrons, for example) may be localized. The clearest case is that of spin diffusion” (Bloembergen 1949; Portis 1956); another might be the so-called impurity band conduction at low concentrations of impurities. In such situations we suspect that transport occurs not by motion of free carriers (or spin waves), scattered as they move through a medium, but in some sense by quantum-mechanical jumps of the mobile entities from site to site. A second common feature of these phenomena is randomness: random spacings of impurities, random interactions with the “atmosphere” of other impurities, random arrangements of electronic or nuclear spins, etc.

Our eventual purpose in this work will be to lay the foundation for a quantum-mechanical theory of transport problems of this type. Therefore, we must start with simple theoretical models rather than with the complicated experimental situations on spin diffusion or impurity conduction. In this paper, in fact, we attempt only to construct, for such a system, the simplest model we can think of which still has some expectation of representing a real physical situation reasonably well, and to prove a theorem about the model. The theorem is that at sufficiently low densities, transport does not take place; the exact wave functions are localized in a small region of space. We also obtain a fairly good estimate of the critical density at which the theorem fails. An additional criterion is that the forces be of sufficiently short range—actually, falling off as $r \rightarrow \infty$ faster than $1/r^3$ —and we derive a rough estimate of the rate of transport in the $V \propto 1/r^3$ case.

...

2. Summary of the reasoning

Since the mathematical development is fairly complicated and involves lengthy consideration of each of a number of points, we should like to summarize the reasoning rather fully in this section, leaving the proofs and details to later sections. First, then, let us set up the simple model which we study. The equation for the time-dependence of the probability amplitude a_j , that a particle is on the site j is:

$$i \dot{a}_j = E_j a_j + \sum_{k \neq j} V_{jk} a_k. \quad (1)$$

References

- Bloembergen, N. 1949. "On the Interaction of Nuclear Spins in a Crystalline Lattice." *Physica* 15 (3): 386–426. [https://doi.org/10.1016/0031-8914\(49\)90114-7](https://doi.org/10.1016/0031-8914(49)90114-7).
- Portis, A. M. 1956. "Spectral Diffusion in Magnetic Resonance." *Phys. Rev.* 104 (November):584–88. <https://doi.org/10.1103/PhysRev.104.584>.