

Dr. Anna R. Giuliano

DISCLOSURE OF CONFLICTS OF INTEREST

GLOBAL EDUCATION GROUP (GLOBAL) REQUIRES INSTRUCTORS, PLANNERS, MANAGERS AND OTHER INDIVIDUALS AND THEIR SPOUSE/LIFE PARTNER WHO ARE IN A POSITION TO CONTROL THE CONTENT OF THIS ACTIVITY TO DISCLOSE ANY REAL OR APPARENT CONFLICT OF INTEREST THEY MAY HAVE AS RELATED TO THE CONTENT OF THIS ACTIVITY. ALL IDENTIFIED CONFLICTS OF INTEREST ARE THOROUGHLY VETTED BY GLOBAL FOR FAIR BALANCE, SCIENTIFIC OBJECTIVITY OF STUDIES MENTIONED IN THE MATERIALS OR USED AS THE BASIS FOR CONTENT, AND APPROPRIATENESS OF PATIENT CARE RECOMMENDATIONS.

THE FACULTY REPORTED THE FOLLOWING FINANCIAL RELATIONSHIPS OR RELATIONSHIPS TO PRODUCTS OR DEVICES THEY OR THEIR SPOUSE/LIFE PARTNER HAVE WITH COMMERCIAL INTERESTS RELATED TO THE CONTENT OF THIS CME ACTIVITY:

DR. ANNA GIULIANO HAS DISCLOSED THE FOLLOWING RELEVANT FINANCIAL RELATIONSHIPS WITHIN THE PAST 12 MONTHS:

Consultant/Independent Contractor: Merck
Grant/Research Support: Merck
Honoraria: Merck
Speaker's Bureau: Merck



Communicating an Integrated HPV Message

Anna R. Giuliano, PhD

Director, Center for Infection Research in Cancer Moffitt Cancer Center



Disclosure

• Dr. Giuliano: Grants/Research Support: Merck & Co., Inc.; Consultant: Merck & Co., Inc.; Speakers Bureau: Merck & Co., Inc.

"Knowledge is not enough; we must apply. Willing is not enough; we must do."


Johann Wolfgang von Goethe

HPV is a Necessary Cause of Cervical Cancer

- Association of cervical neoplasia with sexual activity
- In the mid-1970's Professor zur Hausen identified HPV as a likely viral candidate in cervical neoplasia.
- ~100% of cervical cancer and majority of HgSIL contain HPV DNA
- Strong and consistent associations between HPV and cervical cancer (OR=9 to >100)
- HPV infection precedes onset of disease
- ~50% of cervical tumors contain HPV 16 DNA, an additional ~20% of tumors contain HPV 18, and an additional 20% HPV 31, 45, 33, 52, 58 DNA

There Is a High HPV Disease Burden Among Males and Females Globally

Estimated annual new HPV-related disease cases in males and females globally



Published HPV prevalence rates were applied as follows: Parkin D et al. *Vaccine*. 2006 (penile, vulvar, anal, cervical cancers); WHO/ICO 2010 (head and neck cancer); De Vuyst H et al. *Int J Cancer*. 2009 (vaginal cancer); Greer CE et al. *J Clin Microbiol*. 1995 (genital warts).

^{1.} Parkin DM et al. *Vaccine*. 2006;24(Suppl 3):S3/11–S3/25. 2. WHO/ICO Information Centre on HPV and Cervical Cancer (HPV Information Centre). Human Papillomavirus and Related Cancers in World. Summary Report 2010. http://www.who.int/hpvcentre/en/. Accessed June 21, 2012. 3. World Health Organization (WHO). Executive summary: the state of world health. 1995. http://www.who.int/whr/1995/media_centre/executive_summary1/en/index3.html#. Accessed June 7, 2012.

Disease Endpoints Evaluated in Women

Disease Endpoint	GARDASIL™	Cervarix [™]	
Immediate Cervical Cancer Precursors ¹			
HPV 16/18-related CIN 3	✓		
HPV 16/18-related AIS	✓		
Immediate Vulvar Cancer Precursor ²			
HPV 16/18-related VIN 2/3	✓		
Immediate Vaginal Cancer Precursor ²			
HPV 16/18-related ValN 2/3	✓		
High-grade Cervical Dysplasia ^{1,3}			
HPV 16/18-related CIN 2/3	✓	✓	
HPV 16-related CIN 2/3	✓	✓	
HPV 18-related CIN 2/3	✓	✓	
Low-grade Cervical Dysplasia ³⁻⁵			
HPV 16/18-related CIN 1	✓	✓	
HPV 6/11-related CIN 1	√		
External Genital Lesions ⁴			
Genital warts	✓		

Gardasil Vaccine Efficacy

Per-Protocol Populations

Population	HPV 6/11/16/18-Related Endpoints	Per Protocol Efficacy (%)	95% CI
16-26 year-old women	CIN 2/3	98	93, 100
	VIN 2/3	100	67, 100
	VaIN 2/3	100	55, 100
	Genital warts	99	96, 100


FDA Approval and CDC routine vaccination recommendation for females in 2006

9-Valent HPV Vaccine in Final Year of Phase III Testing in Females

Burden of HPV-Related Disease in Males

- Genital warts
- Recurrent respiratory papillomatosis
- >90% caused by HPV 6 & 11^{1,2}


- Anal cancer
- Penile cancer
- Oropharyngeal and oral cavity cancers



There are no routinely available reliable screening methods for cancers caused by HPV in men


^{1.} Greer CE et al. *J Clin Microbiol*. 1995;33:2058–2063. **2.** Freed GL et al. *Int J Pediatr Otorhinolaryngol*. 2006;70:1799-1803. **3.** De Vuyst H et al. *Int J Cancer*. 2009;124:1626-1636. **4.** Miralles-Guri C et al. *J Clin Pathol*. 2009;62:870-878. **5.** Kreimer AR et al. *Cancer Epidemiol Biomarkers Prev*. 2005;14:467-475.

Age-Adjusted Incidence of Invasive Anal Cancer by Gender and Year of Diagnosis: United States




Howlader N et al. (eds). SEER Cancer Statistics Review, 1975-2009 (Vintage 2009 Populations). http://seer.cancer.gov/csr/1975_2009_pops09/. Accessed June 21, 2012.

US Annual Incidence of Invasive Anal Cancer




Observed and Projected Incidence Rates for Oropharyngeal Cancers and Cervical Cancer




Risk of Acquiring an HPV Infection in Males Does Not Differ by Age


Total 12-month probability=0.39 (95% CI: 0.35, 0.44)



Gender Gaps in HPV 16 Seroprevalence



Low Seroconversion Following HPV HPV 16 and 18 DNA Detection



HPV VACCINATION OF MALES REDUCES DISEASE BURDEN

Efficacy Against HPV 6/11/16/18 Related External Genital Lesions (EGL)

Per-protocol population

	•	ccine 1,397)	Placebo (n = 1,408)				
Endpoint	Cases	Inc. per 100 PY	Cases	Inc. per 100 PY	% Efficacy	95% CI	p-value
All subjects	3	0.1	31	1.1	90.4	69.2, 98.1	<0.001

EGLs include external genital warts, penile/perianal/perineal intraepithelial neoplasia (PIN), penile, perianal, or perineal cancer; case counting began after month 7.

Efficacy Against HPV 6/11/16/18—Related Anal Disease Among MSM

Per-protocol population

	qHPV vaccine (n=299)		ne Placebo (n=299)				
Endpoint	Cases	Inc. per 100 PY	Cases	Inc. per 100 PY	% Efficacy	95% CI	<i>P</i> value
All subjects	5	1.3	24	5.8	77.5	39.6–93.3	<0.001

- Anal intraepithelial neoplasia
- Case counting began at month 7

Vaccine Indications in Men in Several Countries


- Prevention of genital warts 2009
- Anal cancer prevention (males and females) 2010

US ACIP approved routine vaccination of males - 2011


▼ HMO coverage

MDs advised to actively recommend vaccine to boys as well as girls

Proportion of Australian born women under 21 years with genital warts, 2004-2011



US HPV Vaccination[†] Coverage 2011



† Quadrivalent or bivalent vaccine Source: National Immunization Survey (NIS) Teen, US, 2011

Summary

- HPV causes multiple diseases in females and males
- Currently licensed and recommended 4-valent HPV vaccine will prevent in <u>women</u>:
 - Genital warts, cervical, vaginal, and vulvar cancers
- Currently licensed and recommended 4-valent HPV vaccine will prevent in men:
 - genital warts and anal cancer
- HPV vaccine dissemination in the US is <50% in females and <10% in males

Eliminating HPV related disease depends on dissemination of proven technologies

Acknowledgements

HIM Study Team

USA

Anna R Giuliano, Ph.D., Principal Investigator, Moffitt Cancer Center

Drs. Lin, Messina – Co-Investigators

Martha Abrahamsen, Mary Papenfuss, Christine Gage, Kathy Eyring, Nadia Lambermont, Kayoko Kennedy, Kim Isaacs, Andrea Leto, Amanda Sivia, Danelle Smith, Abidemi Ajidahun, Mike Blackner, Bradley Sirak, Alan Nyitray, Beibei Lu, Christine Pierce Campbell, Donna Ingles, Suzette Torres

Ray Viscidi, PhD., Co-Investigator, Hopkins

Cuernavaca, MEXICO

Eduardo Lazcano, Ph.D., Co-Investigator, Instituto Nacional de Salud Pública

Dr. Jorge Salmerón Castro, MC Manuel Quiterio Trenado, Dr. Aurelio Cruz Valdez, Dr. René de Jesús Alvear Vásquez, Dr. Oscar Rojas Juárez, Dra. Rossana del Carmen González Sosa, Enf. Rosangel Ríos Vences, Enf. Martha Huerta Segura, LE Alicia Rodríguez Galván, LE Paula Román Rodríguez, Biol. Ana Laura Landa Vélez, Enf. Griselda Díaz García, LI Verónica Chávez Abarca, Enf. Gisela Flores Quevedo, MSP. María del Pilar Hernández Nevárez, M.C. Guillermina Sánchez Martínez, Biol. Adriana Ortiz Rojas, T.I. Carlos Omar Barrera Flores

São Paulo, BRAZIL

Luisa L. Villa, Ph.D., Co-Investigator, Ludwig Institute for Cancer Research

Maria Luiza Baggio, Roberto C Silva, Lenice Galan, Rubens Matsuo, Elimar Gomes, Filomena Cernicchiaro, Vera Souza, Ricardo Cintra, Viviani Relvas, Birgit Fietzek, Raquel Hessel, Cecília Costa, Fernanda Silva, Juliana Antunes, Graças Ribeiro, Roberta Bocalon, Rosária Otero, Rossana Terreri, Sandra Araujo

Protocol 020 Male HPV Vaccine Trial Investigators, Merck

National Cancer Institute

Funding

Aimee R. Kreimer, Ph.D. Allan Hildesheim, PhD

NCI, NIH CA R01CA098803 (PI: AR Giuliano) Merck & CO, Inc

Questions and Comments?