

CONSTRUCTING 3-LIE ALGEBRAS

RUIPU BAI AND YONG WU

ABSTRACT. 3-Lie algebras are constructed by Lie algebras, derivations and linear functions, associative commutative algebras, whose involutions and derivations. Then the 3-Lie algebras are obtained from group algebras $F[G]$. An infinite dimensional simple 3-Lie algebra $(A, [, ,]_{\omega, \delta_0})$ and a non-simple 3-Lie algebra $(A, [, ,]_{\omega_1, \delta})$ are constructed by Laurent polynomials $A = F[t, t^{-1}]$ and its involutions ω and ω_1 and derivations δ and δ_0 . At last of the paper, we summarize the methods of constructing n -Lie algebras for $n \geq 3$ and provide a problem.

1. INTRODUCTION

To construct the generalized Hamiltonian dynamics, Nambu [1, 2] first proposed the notion of 3-bracket. Thus Nambu dynamics is described by the phase flow given by Nambu-Hamilton equations of motion which involves two Hamiltonians. The notion of n -Lie algebra was introduced by Filippov in 1985 ([3]). It is a natural generalization of the concept of a Lie algebra to the case where the fundamental multiplication is n -ary, $n \geq 2$ (when $n = 2$ the definition agrees with the usual definition of a Lie algebra). 3-Lie algebras have close relationships with many important fields in mathematics and mathematical physics (cf. [1, 2, 4, 5, 6, 7, 8]). For example, the metric 3-Lie algebras are used to describe a world volume of multiple $M2$ -branes [4, 7] (BLG).

Since the multiple multiplication, the structure of n -Lie algebras is more complicated than that of Lie algebras. Especially, the realizations of n -Lie algebras is hardness.

Filippov in [9] constructed n -Lie algebra structure on commutative associative algebras by commuting derivations $\{D_1, \dots, D_n\}$. Let A be a commutative associative algebra, then A is an n -Lie algebra with the multiplication:

$$(1.1) \quad [x_1, \dots, x_n] = \det \begin{pmatrix} D_1(x_1) & \cdots & D_1(x_n) \\ \cdots & \cdots & \cdots \\ D_n(x_1) & \cdots & D_n(x_n) \end{pmatrix},$$

which is called a *Jacobian algebra* defined by $\{D_1, \dots, D_n\}$.

2010 *Mathematics Subject Classification.* 17A30, 17B60.

Key words and phrases. 3-Lie algebra, Lie algebra, associative commutative algebra, involution, derivation.

Filippov and Pozhidaev in papers [9, 11, 12] defined *monomial n -Lie algebra* $A_G(f, t)$ with the multiplication

$$(1.2) \quad [e_{a_1}, \dots, e_{a_n}] = f(a_1, \dots, a_n)e_{a_1+\dots+a_n+t},$$

where G is an Abelian group, A_G is a vector space with a basis $\{e_g | g \in G\}$, $t \in G$, $f : G^n \rightarrow F$.

Dzhumadil'daev in [10] constructed an $(n+1)$ -Lie algebra from a strong n -Lie-Poisson algebra (L, \cdot, ω) by endowing a skew-symmetric $(n+1)$ -multiplication

$$(1.3) \quad \bar{\omega} = Id_L \wedge \omega,$$

where Id_L is the identity map of L .

Bai and collaborators in [13, 14] constructed n -Lie algebras by two-dimensional extensions of metric n -Lie algebras. And obtain $(n+1)$ -Lie algebras by n -Lie algebras and linear functions $f \in L^*$, which satisfies $f(L^1) = 0$.

In this paper, we pay our main attention to construct 3-Lie algebras. The paper is organized as follows. Section 2 introduces some basic notions of n -Lie algebras. Section 3 provides a construction of 3-Lie algebras from a commutative associative algebras, involutions and derivations. Section 4 constructs 3-Lie algebras from group algebras. Section 5 studies 3-Lie algebras constructed by Laurent polynomials.

2. Fundamental notions of n -Lie algebra

An n -Lie algebra L (cf.[3]) is a vector space endowed with an n -ary multilinear skew-symmetric multiplication satisfying the n -Jacobi identity: $\forall x_1, \dots, x_n, y_2, \dots, y_n \in L$

$$(2.1) \quad [[x_1, \dots, x_n], y_2, \dots, y_n] = \sum_{i=1}^n [x_1, \dots, [x_i, y_2, \dots, y_n], \dots, x_n].$$

The n -ary skew-symmetry of the operation $[x_1, \dots, x_n]$ means that

$$(2.2) \quad [x_1, \dots, x_n] = \text{sgn}(\sigma)[x_{\sigma(1)}, \dots, x_{\sigma(n)}], \quad \forall x_1, \dots, x_n \in L,$$

for any permutation $\sigma \in S_n$.

A subspace B of L is called a *subalgebra* if $[B, \dots, B] \subseteq B$. In particular, the subalgebra generated by the vectors $[x_1, \dots, x_n]$ for any $x_1, \dots, x_n \in L$ is called the *derived algebra* of L , which is denoted by L^1 . If $L^1 = 0$, then L is called an *abelian n -Lie algebra*.

A *derivation* of an n -Lie algebra is a linear transformation D of L into itself satisfying

$$(2.3) \quad D([x_1, \dots, x_n]) = \sum_{i=1}^n [x_1, \dots, D(x_i), \dots, x_n],$$

for $x_1, \dots, x_n \in L$. Let $Der L$ be the set of all derivations of L . Then $Der L$ is a subalgebra of the general Lie algebra $gl(L)$ and is called *the derivation algebra* of L .

The map $\text{ad}(x_1, \dots, x_{n-1}) : L \rightarrow L$ given by

$$\text{ad}(x_1, \dots, x_{n-1})(x_n) = [x_1, \dots, x_{n-1}, x_n], \forall x_n \in L,$$

is referred to as a *left multiplication* defined by elements $x_1, \dots, x_{n-1} \in L$. It follows from identity (2.1) that $\text{ad}(x_1, \dots, x_{n-1})$ is a derivation. The set of all finite linear combinations of left multiplications is an ideal of $\text{Der}L$, which is denoted by $\text{ad}(L)$. Every derivation in $\text{ad}(L)$ is by definition an inner derivation.

An *ideal* of an n -Lie algebra L is a subspace I such that $[I, L, \dots, L] \subseteq I$. If $L^1 \neq 0$ and L has no ideals except for 0 and itself, then L is called a *simple n -Lie algebra*.

3. 3-LIE ALGEBRAS CONSTRUCTED BY COMMUTATIVE ASSOCIATIVE ALGEBRAS

R.Bai and collaborators (cf [14]) constructed 3-Lie algebra $(L, [,], \alpha)$ by a Lie algebra $(L, [,])_{\alpha}$ and linear function $\alpha \in L^*$, where $\alpha([L, L]) = 0$, and $[,], \alpha$ defined as follows

$$[a, b, c]_{\alpha} = \alpha(c)[a, b] + \alpha(a)[b, c] + \alpha(b)[c, a], \text{ for } a, b, c \in L.$$

And it is proved that the 3-Lie algebra $(L, [,], \alpha)$ is a two step 3-solvable 3-Lie algebra, that is, $L^{(2)} = [L^1, L^1, L^1]_{\alpha} = 0$, where $L^1 = [L, L, L]_{\alpha}$.

In this section, we try to construct 3-Lie algebras by commutative associative algebras and their derivations, involutions and linear functions.

Let A be a commutative associative algebra over a field F . A derivation Δ of A is a linear mapping of A satisfying $\Delta(xy) = \Delta(x)y + x\Delta(y)$ for every $x, y \in A$. If a linear mapping $\omega : A \rightarrow A$ satisfying for every $a, b \in A$, $\omega(ab) = \omega(a)\omega(b)$ and $\omega^2(a) = a$, then ω is called an *involution* of A . If $\text{ch}F \neq 2$, then for every $c \in A$,

$$\omega(c + \omega(c)) = c + \omega(c), \omega(c - \omega(c)) = -(c - \omega(c)), \text{ and } c = \frac{1}{2}(c + \omega(c)) + \frac{1}{2}(c - \omega(c)).$$

Therefore,

$$A = A_1 \dot{+} A_{-1}, \text{ where } A_1 = \{a \mid a \in A, \omega(a) = a\}, A_{-1} = \{b \mid b \in A, \omega(b) = -b\}.$$

First we give the following result.

Lemma 3.1. *Let A be a commutative associative algebra, ω be an involution of A and $\Delta \in \text{Der}A$. Then $(A, [,]_{\Delta})$ and $(A, [,]_{\omega})$ are Lie algebras, where*

$$(3.1) \quad [a, b]_{\Delta} = (Id_A \wedge \Delta)(a, b) = a\Delta(b) - b\Delta(a), \quad a, b \in A;$$

$$(3.2) \quad [a, b]_{\omega} = (\omega \wedge Id_A)(a, b) = \omega(a)b - \omega(b)a, \quad a, b \in A.$$

If Δ and ω satisfy $\Delta\omega + \omega\Delta = 0$, then $(A, [,]_{\omega, \Delta})$ is a Lie algebra, where

$$(3.3) \quad [a, b]_{\omega, \Delta} = ((Id_A - \omega) \wedge \Delta)(a, b) = (a - \omega(a))\Delta(b) - (b - \omega(b))\Delta(a), \quad a, b \in A.$$

Proof. Eq. (3.1) is well know result. By the direct computation, Eqs. (3.2) and (3.3) satisfy Eq (2.1), respectively. \square

Theorem 3.2. *Let A be a commutative associative algebra, $\Delta \in \text{Der}(A)$ and ω be an involution of A . If $\alpha, \beta, \gamma \in A^*$ satisfy*

$$(3.4) \quad \alpha(ab) = 0, \quad \beta(a\Delta(b) - b\Delta(a)) = 0, \quad \gamma(a\Delta(b) - b\Delta(a)) = \gamma(\omega(a)\Delta(b) - \omega(b)\Delta(a)),$$

then $(A, [, ,]_{\alpha, f})$ and $(A, [, ,]_{\beta, \Delta})$ are 3-Lie algebras, where for arbitrary $a, b, c \in A$,

$$(3.5) \quad [a, b, c]_{\alpha, \omega} = (\alpha \wedge \text{Id}_A \wedge \omega)(a, b, c) = \alpha(a)[b, c]_{\omega} + \alpha(b)[c, a]_{\omega} + \alpha(c)[a, b]_{\omega},$$

$$(3.6) \quad [a, b, c]_{\beta, \Delta} = (\beta \wedge \text{Id}_A \wedge \Delta)(a, b, c) = \beta(a)[b, c]_{\Delta} + \beta(b)[c, a]_{\Delta} + \beta(c)[a, b]_{\Delta}.$$

If Δ and ω satisfy $\Delta\omega + \omega\Delta = 0$, then $(A, [, ,]_{\gamma, \omega, \Delta})$ is a 3-Lie algebra, where

$$(3.7) \quad [a, b, c]_{\gamma, \omega, \Delta} = (\gamma \wedge (\text{Id}_A - \omega) \wedge \Delta)(a, b, c) = \gamma(a)((b - \omega(b))\Delta(c) - (c - \omega(c))\Delta(b)) \\ + \gamma(b)((c - \omega(c))\Delta(a) - (a - \omega(a))\Delta(c)) + \gamma(c)((a - \omega(a))\Delta(b) - (b - \omega(b))\Delta(a)).$$

Proof. Let $\alpha \in A^*$. Then for every $a, b \in A$,

$$\alpha(\omega(a)b - \omega(b)a) = \begin{cases} 0, & \text{if } a, b \in A_1; \text{ or } a, b \in A_{-1}, \\ 2\alpha(ab), & \text{if } a \in A_1, b \in A_{-1}, \\ -2\alpha(ab), & \text{if } b \in A_1, a \in A_{-1}. \end{cases}$$

Therefore, if $\alpha(ab) = 0$, then $\alpha(\omega(a)b - \omega(b)a) = 0$. From Lemma 3.1, Eq. (3.1), and Theorem 3.1 in [14], the result holds. \square

Theorem 3.3. *Let A be a commutative associative algebra over a field F , Δ be a derivation of A and $\omega : A \rightarrow A$ be an involution of A satisfying*

$$f\Delta + \Delta\omega = 0.$$

Then A is a 3-Lie algebra in the multiplication $[, ,]_{\omega, \Delta} : A \otimes A \otimes A \rightarrow A$, $\forall a, b, c \in A$,

$$(3.8) \quad [a, b, c]_{\omega, \Delta} = \omega \wedge \text{Id}_A \wedge \Delta(a, b, c) = \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta(a) & \Delta(b) & \Delta(c) \end{vmatrix} \\ = \omega(a)(b\Delta(c) - c\Delta(b)) + \omega(b)(c\Delta(a) - a\Delta(c)) + \omega(c)(a\Delta(b) - b\Delta(a)).$$

Proof. It is clear that $[, ,]_{\omega, \Delta}$ is a 3-ary linear skew-symmetric multiplication on A . Now we prove that $[, ,]_{\omega, \Delta}$ satisfies Eq.(2.1). Since A is commutative and ω is an involution of A which satisfies $\omega\Delta + \Delta\omega = 0$, by Eq.(3.8), $\forall a, b, c, d, e \in A$

$$\begin{aligned}
\omega([a, b, c]_{\omega, \Delta}) &= \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta\omega(a) & \Delta\omega(b) & \Delta\omega(c) \end{vmatrix}, \\
\Delta([a, b, c]_{\omega, \Delta}) &= \begin{vmatrix} \Delta\omega(a) & \Delta\omega(b) & \Delta\omega(c) \\ a & b & c \\ \Delta(a) & \Delta(b) & \Delta(c) \end{vmatrix} + \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta^2(a) & \Delta^2(b) & \Delta^2(c) \end{vmatrix}, \\
[[a, b, c]_{\omega, \Delta}, d, e]_{\omega, \Delta} &= \begin{vmatrix} \omega([a, b, c]_{\omega, \Delta}) & \omega(d) & \omega(e) \\ [a, b, c]_{\omega, \Delta} & d & e \\ \Delta([a, b, c]_{\omega, \Delta}) & \Delta(d) & \Delta(e) \end{vmatrix} \\
&= \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta\omega(a) & \Delta\omega(b) & \Delta\omega(c) \end{vmatrix} \begin{vmatrix} d & e \\ \Delta(b) & \Delta(c) \end{vmatrix} - \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta(a) & \Delta(b) & \Delta(c) \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ \Delta(d) & \Delta(e) \end{vmatrix} \\
&+ \begin{vmatrix} \Delta\omega(a) & \Delta\omega(b) & \Delta\omega(c) \\ a & b & c \\ \Delta(a) & \Delta(b) & \Delta(c) \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ d & e \end{vmatrix} + \begin{vmatrix} \omega(a) & \omega(b) & \omega(c) \\ a & b & c \\ \Delta^2(a) & \Delta^2(b) & \Delta^2(c) \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ d & e \end{vmatrix}.
\end{aligned}$$

Then we have

$$\begin{aligned}
&[[a, d, e]_{\omega, \Delta}, b, c]_{\omega, \Delta} + [[b, d, e]_{\omega, \Delta}, c, a]_{\omega, \Delta} + [[c, d, e]_{\omega, \Delta}, a, b]_{\omega, \Delta} \\
&= \circlearrowleft_{a, b, c} \Delta\omega(a) \begin{vmatrix} b & c \\ \Delta(b) & \Delta(c) \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ d & e \end{vmatrix} + \circlearrowleft_{a, b, c} a \begin{vmatrix} \omega(b) & \omega(c) \\ \Delta(b) & \Delta(c) \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ \Delta(d) & \Delta(e) \end{vmatrix} \\
&+ \circlearrowleft_{a, b, c} \Delta\omega(a) \begin{vmatrix} \omega(b) & \omega(c) \\ b & c \end{vmatrix} \begin{vmatrix} d & e \\ \Delta(d) & \Delta(e) \end{vmatrix} + \circlearrowleft_{a, b, c} \Delta^2(a) \begin{vmatrix} \omega(b) & \omega(c) \\ b & c \end{vmatrix} \begin{vmatrix} \omega(d) & \omega(e) \\ d & e \end{vmatrix} \\
&= [[a, b, c]_{\omega, \Delta}, d, e]_{\omega, \Delta},
\end{aligned}$$

where $\circlearrowleft_{a, b, c}$ is the circulation of a, b, c , for example

$$\circlearrowleft_{a, b, c} \Delta\omega(a) \begin{vmatrix} b & c \\ \Delta(b) & \Delta(c) \end{vmatrix} = \Delta\omega(a) \begin{vmatrix} b & c \\ \Delta(b) & \Delta(c) \end{vmatrix} + \Delta\omega(b) \begin{vmatrix} c & a \\ \Delta(c) & \Delta(a) \end{vmatrix} + \Delta\omega(c) \begin{vmatrix} a & b \\ \Delta(a) & \Delta(b) \end{vmatrix}.$$

Therefore, $(A, [,],_{\omega, \Delta})$ is a 3-Lie algebra in the multiplication (3.8). \square

Corollary 3.4. *Let A be a commutative associative algebra, $\Delta \in \text{Der}(A)$ and ω be an involution of A satisfying $\Delta\omega + \omega\Delta = 0$. Then 3-Lie algebra $(A, [,],_{\Delta, \omega})$ with the multiplication (3.8) can be decomposed into the direct sum of abelian subalgebras A_1 and A_{-1} , that is, $A = A_1 \dot{+} A_{-1}$, and*

$$[A_1, A_1, A_1]_{\omega, \Delta} = [A_{-1}, A_{-1}, A_{-1}]_{\omega, \Delta} = 0.$$

And $\Delta(A_1) \subseteq A_{-1}$, $\Delta(A_{-1}) \subseteq A_1$.

Proof. For every $a_i \in A_1$, $\omega(a_i) = a_i$ for $i = 1, 2, 3$, $\omega\Delta(a_1) = -\Delta\omega(a_1) = -\Delta(a_1)$, then

$$[a_1, a_2, a_3]_{\omega, \Delta} = \begin{vmatrix} \omega(a_1) & \omega(a_2) & \omega(a_3) \\ a_1 & a_2 & a_3 \\ \Delta(a_1) & \Delta(a_2) & \Delta(a_3) \end{vmatrix} = \begin{vmatrix} a_1 & a_2 & a_3 \\ a_1 & a_2 & a_3 \\ \Delta(a_1) & \Delta(a_2) & \Delta(a_3) \end{vmatrix} = 0.$$

Thus A_1 is an abelian subalgebra, and $\Delta(A_1) \subseteq A_{-1}$. \square

Theorem 3.5. *Let $(A, [,],_{\omega, \Delta})$ be the 3-Lie algebra with the multiplication (3.8). If I is an ideal of the associative algebra A which satisfies $\omega(I) \subseteq I$ and $\Delta(I) \subseteq I$. Then I is an ideal of the 3-Lie algebra $(A, [,],_{\omega, \Delta})$.*

Proof. Let I be an ideal of the associative algebra A satisfying $\omega(I) \subseteq I$ and $\Delta(I) \subseteq I$. Then for every $a \in I$, and $\forall b, c \in A$, by Eq. (3.8)

$$[a, b, c]_{\omega, \Delta} = \omega(a)(b\Delta(c) - c\Delta(b)) + \omega(b)(a\Delta(c) - c\Delta(a)) + \omega(c)(a\Delta(b) - b\Delta(a)) \in I.$$

Therefore, I is an ideal of the 3-Lie algebra $(A, [,],_{\omega, \Delta})$. \square

Theorem 3.6. *Let A_1 and A_2 be commutative associative algebras, ω_i be an involution of A_i and Δ_i be a derivation of A_i which satisfies $\omega_i\Delta_i + \Delta_i\omega_i = 0$ for $i = 1, 2$. If $\sigma : A_1 \rightarrow A_2$ is an associative algebra isomorphism satisfying $\sigma\omega_1 = \omega_2\sigma$, and $\sigma\Delta_1 = \Delta_2\sigma$. Then σ is a 3-Lie algebra isomorphism from $(A_1, [,],_{\omega_1, \Delta_1})$ onto $(A_2, [,],_{\omega_2, \Delta_2})$.*

Proof. For every $a, b, c \in A_1$, by Eq.(3.8),

$$\begin{aligned} \sigma([a, b, c]_{\omega_1, \Delta_1}) &= \sigma(\omega_1(a)(b\Delta_1(c) - c\Delta_1(b)) + \omega_1(b)(c\Delta_1(a) - a\Delta_1(c)) + \omega_1(c)(a\Delta_1(b) - b\Delta_1(a))) \\ &= \omega_2(\sigma(a))(\sigma(b)\Delta_2\sigma(c) - \sigma(c)\Delta_2(\sigma(b))) + \omega_2(\sigma(b))(\sigma(c)\Delta_2(\sigma(a)) - \sigma(a)\Delta_2(\sigma(c))) \\ &\quad + \omega_2(\sigma(c))(\sigma(a)\Delta_2(\sigma(b)) - \sigma(b)\Delta_2(\sigma(a))) \\ &= [\sigma(a), \sigma(b), \sigma(c)]_{\omega_2, \Delta_2}. \end{aligned}$$

It follows the result. \square

Corollary 3.7. *If A is a nilpotent commutative associative algebra, then $(A, [,],_{\omega, \Delta})$ is a nilpotent 3-Lie algebra.*

Proof. Suppose $A^m = 0$ for some positive integer m . Then for every $a_1, \dots, a_m \in A$, $a_1 \cdots a_m = 0$. Thanks to Eq. (3.8), for every $a, b \in A$, $ad^m(a, b)(A) \subseteq A^m = 0$. Therefore, $(A, [,],_{\omega, \Delta})$ is a nilpotent 3-Lie algebra. \square

4. 3-LIE ALGEBRAS CONSTRUCTED BY GROUP ALGEBRAS

Let $(G, +)$ be an additive Abelian group, and $F[G]$ be the group algebra, that is, $F[G]$ is a commutative associative algebra with a basis $\{e_g \mid g \in G\}$, and for every $x = \sum_{g \in G} \lambda_g e_g$,

$$y = \sum_{h \in G} \mu_h e_h \in F[G],$$

$$x + y = \sum_{g \in G} (\lambda_g + \mu_g) e_g, \quad xy = \left(\sum_{g \in G} \lambda_g e_g \right) \left(\sum_{h \in G} \mu_h e_h \right) = \sum_{g, h \in G} \lambda_g \mu_h e_{g+h}.$$

Define linear mapping $\omega : F[G] \rightarrow F[G]$,

$$(4.1) \quad \omega(x) = \omega\left(\sum_{g \in G} \lambda_g e_g\right) = \sum_{g \in G} \lambda_g e_{-g}, \quad \forall x = \sum_{g \in G} \lambda_g e_g \in F[G].$$

Then ω is a linear isomorphism of $F[G]$ and for $x = \sum_{g \in G} \lambda_g e_g, y = \sum_{h \in G} \mu_h e_h \in F[G]$,

$$\omega^2(x) = \omega\left(\sum_{g \in G} \lambda_g e_{-g}\right) = \sum_{g \in G} \lambda_g e_g = x,$$

$$\begin{aligned} \omega(xy) &= \omega\left(\sum_{g \in G} \lambda_g e_g \sum_{h \in G} \mu_h e_h\right) = \omega\left(\sum_{g, h \in G} \lambda_g \mu_h e_{g+h}\right) = \sum_{g, h \in G} \lambda_g \mu_h e_{-g-h} \\ &= \sum_{g \in G} \lambda_g e_{-g} \sum_{h \in G} \mu_h e_{-h} = \omega\left(\sum_{g \in G} \lambda_g e_g\right) \omega\left(\sum_{h \in G} \mu_h e_h\right) = \omega(x) \omega(y). \end{aligned}$$

Therefore, ω is an involution of the commutative associative algebra $F[G]$.

Denote F^+ the addition group of F . For every $\alpha \in \text{Hom}(G, F^+)$, then α satisfies $\alpha(g + h) = \alpha(g) + \alpha(h), \forall g, h \in G$. Define linear mapping $\alpha^* : F[G] \rightarrow F[G]$ as follows:

$$(4.2) \quad \alpha^*\left(\sum_{g \in G} \lambda_g e_g\right) = \sum_{g \in G} \lambda_g \alpha(g) e_g, \quad \forall \sum_{g \in G} \lambda_g e_g \in F[G].$$

Lemma 4.1. *Let $(G, +)$ be an abelian group, $\omega : F[G] \rightarrow F[G]$ be defined as Eq.(4.1). Then for every $\alpha \in \text{Hom}(G, F^+)$, α^* defined as Eq. (4.2) is a derivation of the algebra $F[G]$, and satisfies $\omega\alpha^* + \alpha^*\omega = 0$.*

Proof. By Eqs. (4.1) and (4.2), for arbitrary $x = \sum_{g \in G} \lambda_g e_g, y = \sum_{h \in G} \mu_h e_h \in F[G]$,

$$\begin{aligned} \alpha^*\left(\left(\sum_{g \in G} \lambda_g e_g\right)\left(\sum_{h \in G} \mu_h e_h\right)\right) &= \alpha^*\left(\sum_{g, h \in G} \lambda_g \mu_h e_{g+h}\right) = \sum_{g, h \in G} \lambda_g \mu_h (\alpha(g) + \alpha(h)) e_{g+h} \\ &= \sum_{g, h \in G} \lambda_g \mu_h \alpha(g) e_{g+h} + \sum_{g, h \in G} \lambda_g \mu_h \alpha(h) e_{g+h} = \alpha^*\left(\sum_{g \in G} \lambda_g e_g\right) \left(\sum_{h \in G} \mu_h e_h\right) + \left(\sum_{g \in G} \lambda_g e_g\right) \alpha^*\left(\sum_{h \in G} \mu_h e_h\right), \\ (\omega\alpha^* + \alpha^*\omega)\left(\sum_{g \in G} \lambda_g e_g\right) &= \omega\left(\sum_{g \in G} \lambda_g \alpha(g) e_g\right) + \alpha^*\left(\sum_{g \in G} \lambda_g e_{-g}\right) \\ &= \sum_{g \in G} \lambda_g \alpha(g) e_{-g} + \sum_{g \in G} \lambda_g \alpha(g^{-1}) e_{-g} = \sum_{g \in G} \lambda_g \alpha(g) e_{-g} - \sum_{g \in G} \lambda_g \alpha(g) e_{-g} = 0. \end{aligned}$$

It follows the result. \square

Theorem 4.2. *Let G be an abelian group, $\omega : F[G] \rightarrow F[G]$ be defined as Eq.(4.1), $\alpha \in \text{Hom}(G, F^+)$, α^* be defined as Eq. (4.2). Then $(F[G], [, ,]_{\omega, \alpha^*})$ is a 3-Lie algebra, where for arbitrary $\sum_{g \in G} \lambda_g e_g, \sum_{h \in G} \mu_h e_h, \sum_{q \in G} \nu_q e_q \in F[G]$,*

$$(4.3) \quad \left[\sum_{g \in G} \lambda_g e_g, \sum_{h \in G} \mu_h e_h, \sum_{q \in G} \nu_q e_q \right]_{\omega, \alpha^*} = \sum_{g, h, q \in G} \lambda_g \mu_h \nu_q (\alpha(q - h) e_{h+q-g} + \alpha(g - q) e_{g+q-h} + \alpha(h - g) e_{g+h-q}).$$

Proof. By Lemma 4.1 and Theorem 3.3, for arbitrary $\sum_{g \in G} \lambda_g e_g, \sum_{h \in G} \mu_h e_h, \sum_{q \in G} \nu_q e_q \in F[G]$,

$$\begin{aligned} \left[\sum_{g \in G} \lambda_g e_g, \sum_{h \in G} \mu_h e_h, \sum_{q \in G} \nu_q e_q \right]_{\omega, \alpha^*} &= \begin{vmatrix} \sum_{g \in G} \lambda_g e_{-g} & \sum_{h \in G} \mu_h e_{-h} & \sum_{q \in G} \nu_q e_{-q} \\ \sum_{g \in G} \lambda_g e_g & \sum_{h \in G} \mu_h e_h & \sum_{q \in G} \nu_q e_q \\ \sum_{g \in G} \lambda_g \alpha(g) e_g & \sum_{h \in G} \mu_h \alpha(h) e_h & \sum_{q \in G} \nu_q \alpha(q) e_q \end{vmatrix} \\ &= \sum_{g, h, q \in G} \lambda_g \mu_h \nu_q \{ (\alpha(q) - \alpha(h)) e_{h+q-g} + (\alpha(g) - \alpha(q)) e_{g+q-h} + (\alpha(h) - \alpha(g)) e_{g+h-q} \} \\ &= \sum_{g, h, q \in G} \lambda_g \mu_h \nu_q (\alpha(q - h) e_{h+q-g} + \alpha(g - q) e_{g+q-h} + \alpha(h - g) e_{g+h-q}). \end{aligned}$$

It follows the result. \square

By the above discussions, the products of basis vectors $\{e_g \mid g \in G\}$ of 3-Lie algebra $(F[G], [, ,]_{\omega, \alpha^*})$ are as follows: for arbitrary $g, h, w \in G$,

$$(4.4) \quad [e_g, e_h, e_w]_{\omega, \alpha^*} = \begin{vmatrix} e_{-g} & e_{-h} & e_{-w} \\ e_g & e_h & e_w \\ \alpha(g) e_g & \alpha(h) e_h & \alpha(w) e_w \end{vmatrix} = \alpha(w - h) e_{h+w-g} + \alpha(g - w) e_{g+w-h} + \alpha(h - g) e_{g+h-w}.$$

For $\alpha \in \text{Hom}(G, F^+)$, define mapping

$$(4.5) \quad \phi_\alpha : F[G] \rightarrow F, \quad \phi_\alpha(x) = \sum_{g \in G} \lambda_g \alpha(g), \quad \forall x = \sum_{g \in G} \lambda_g e_g \in F[G].$$

Denote $I_0 = \{x \mid x = \sum_{g \in G} \lambda_g e_g \in F[G], \phi_\alpha(x) = \sum_{g \in G} \lambda_g \alpha(g) = 0\}$. Then I_0 is a subspace of $F[G]$. And we have the following result.

Theorem 4.3. *For $\alpha \in \text{Hom}(G, F^+)$, if $\alpha \neq 0$, then I_0 is a maximal ideal of 3-Lie algebra $(F[G], [, ,]_{\omega, \alpha^*})$. Therefore, $(F[G], [, ,]_{\omega, \alpha^*})$ is a non-simple 3-Lie algebra.*

Proof. For arbitrary $g, h, w \in G$, by Eq. (4.4),

$$\begin{aligned}\phi_\alpha([e_g, e_h, e_w]_{\omega, \alpha^*}) &= \phi_\alpha(\alpha(w-h)e_{h+w-g} + \alpha(g-w)e_{g+w-h} + \alpha(h-g)e_{g+h-w}) \\ &= (\alpha(w) - \alpha(h))(\alpha(h) + \alpha(w) - \alpha(g)) + (\alpha(g) - \alpha(w))(\alpha(g) + \alpha(w) - \alpha(h)) \\ &\quad + (\alpha(h) - \alpha(g))(\alpha(g) + \alpha(h) - \alpha(w)) = 0.\end{aligned}$$

It follows that the derived algebra of $(F[G], [,], \omega, \alpha^*)$ is contained in I_0 . Therefore, I_0 is an ideal of the 3-Lie algebra $(F[G], [,], \omega, \alpha^*)$.

Since $\alpha \neq 0$, without loss of generality suppose $\alpha(d) = 1$ for some non-zero element d of G . Then for every $x \in F[G]$, $x = \phi_\alpha(x)e_d + (x - \phi_\alpha(x)e_d)$. Since

$$\phi_\alpha(x - \phi_\alpha(x)e_d) = \phi_\alpha(x) - \phi_\alpha(x) = 0,$$

we have $F[G] = Fe_d + I_0$ as the direct sum of subspaces. Therefore, I_0 is a maximal ideal of 3-Lie algebra $(F[G], [,], \omega, \alpha^*)$. \square

Example 4.1 Let $G = \{A = (a_{ij}) | a_{ij} \in F, 1 \leq i \leq m, 1 \leq j \leq n\}$ be the set of all $(m \times n)$ -matrices over a field F . Then G is an abelian group in the addition: $\forall A = (a_{ij}), B = (b_{ij}) \in G, A + B = (a_{ij} + b_{ij})$. Define $\alpha : G \rightarrow F^+$ and $\omega : F[G] \rightarrow F[G]$ as follows

$$\alpha(A) = \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^n a_{ij}, \quad \omega(e_A) = e_{-A}.$$

Then $\alpha \in \text{Hom}(G, F^+)$ and ω is an involution of $F[G]$. By Theorem 4.2, $F[G]$ is an mn -dimensional 3-Lie algebra with the multiplication: $\forall A = (a_{ij}), B = (b_{ij}), C = (c_{ij}) \in G$,

$$[e_A, e_B, e_C]_{\omega, \alpha^*} = \sum_{i=1}^m \sum_{j=1}^n (c_{ij} - b_{ij})e_{B+C-A} + \sum_{i=1}^m \sum_{j=1}^n (a_{ij} - c_{ij})e_{C+A-B} + \sum_{i=1}^m \sum_{j=1}^n (b_{ij} - a_{ij})e_{A+B-C}.$$

Example 4.2 Let $G = \{A = (a_{ij}) | a_{ij} \in F, 1 \leq i, j \leq n\}$ be the group of all $(n \times n)$ -matrices over a field F with the addition: $\forall A = (a_{ij}), B = (b_{ij}) \in G, A + B = (a_{ij} + b_{ij})$. Let $\beta \in \text{Hom}(G, F^+)$, $\beta(A) = \text{tr}(A) = \sum_{i=1}^n a_{ii}$, $\omega : F[G] \rightarrow F[G]$, $\omega(e_A) = e_{-A}$. Then by Theorem 4.2, $F[G]$ is an n^2 -dimensional 3-Lie algebra in the multiplication: $\forall A = (a_{ij}), B = (b_{ij}), C = (c_{ij}) \in G$,

$$[e_A, e_B, e_C]_{\omega, \beta^*} = \text{tr}(C - B)e_{B+C-A} + \text{tr}(A - C)e_{A+C-B} + \text{tr}(B - A)e_{A+B-C}.$$

Example 4.3 Let $G = Z_p^+$ be the addition group of the prime field Z_p , $ch F_p = p$. Then the multiplication of the group algebra $Z_p[G]$ is

$$e_{\bar{r}}e_{\bar{s}} = e_{\overline{r+s}}, \quad \forall \bar{s}, \bar{r} \in G.$$

Define

$$\begin{aligned}\alpha : G &\rightarrow Z_p^+, \alpha(\bar{r}) = \bar{r}, \forall \bar{r} \in Z_p; \\ \omega : Z_p[G] &\rightarrow Z_p[G], \omega(e_{\bar{r}}) = e_{-\bar{r}}, \bar{r} \in Z_p.\end{aligned}$$

By Theorem 4.2, $(Z_p[G], [\cdot, \cdot]_{\omega, \alpha^*})$ is a p -dimensional 3-Lie algebra with the multiplication as follows

$$[e_{\bar{r}}, e_{\bar{s}}, e_{\bar{k}}] = \begin{vmatrix} e_{-\bar{r}} & e_{-\bar{s}} & e_{-\bar{k}} \\ e_{\bar{r}} & e_{\bar{s}} & e_{\bar{k}} \\ \bar{r}e_{\bar{r}} & \bar{s}e_{\bar{s}} & \bar{k}e_{\bar{k}} \end{vmatrix} = \overline{k-s}e_{\overline{s+k-r}} + \overline{r-k}e_{\overline{k+r-s}} + \overline{s-r}e_{\overline{r+s-k}}.$$

5. 3-LIE ALGEBRAS CONSTRUCTED FROM LAURENT POLYNOMIALS

In this section we study 3-Lie algebras constructed by Laurent polynomials. In the following, denote $A = F[t^{-1}, t]$, the set of Laurent polynomials over a field F .

We know that the derivation algebra $Der A = \{t^s \frac{d}{dt} \mid s \in Z\}$ with the product

$$(5.1) \quad [t^m \delta, t^n \delta] = (n-m)t^{m+n} \delta, \quad m, n \in Z,$$

where $\delta = t \frac{d}{dt}$, Z is the set of all integer numbers.

Let $\omega : A \rightarrow A$ be an algebra homomorphism and satisfy $\omega^2 = Id_A$. Since $\omega(1) = 1$ and $\omega(t)\omega(t^{-1}) = 1$, we have $\omega(t) = \lambda t^r$, $\lambda \in F$ and $\lambda \neq 0$. Then $\omega(t^m) = \lambda^m t^{rm}$,

$$t^m = \omega^2(t^m) = \omega(\omega(t^m)) = \omega((\omega(t))^m) = \omega(\lambda^m t^{rm}) = \lambda^{m+rm} t^{r^2 m}.$$

We obtain $r = -1$, $\lambda \neq 0$; or $r = 1$, $\lambda = \pm 1$. Therefore, we get the following result.

Lemma 5.1. *Let $\omega : A \rightarrow A$ be a linear map. If $ch F \neq 2$, then ω is an involution of A if and only if ω satisfies*

$$(5.2) \quad \omega(t^m) = \varepsilon^m t^m, \varepsilon = \pm 1, \forall m \in Z; \text{ or } \omega(t^m) = \lambda^m t^{-m}, \lambda \in F, \lambda \neq 0, \forall m \in Z.$$

Proof. The result follows from the above discussions. \square

Lemma 5.2. *Let ω be an involution of A , $\delta = t^l \frac{d}{dt} \in Der F[t^{-1}, t]$ and $ch F \neq 2$. Then $\omega\delta + \delta\omega = 0$ if and only if ω, δ satisfy the following one possibilities*

- (i) $\omega(t^m) = (-1)^m t^m, \forall m \in Z, \delta = t^{2k} \frac{d}{dt}, k \in Z.$
- (ii) $\omega(t^m) = \lambda^m t^{-m}, \lambda \in F, \lambda \neq 0, \forall m \in Z, \delta = t \frac{d}{dt}.$
- (ii) If $ch F = 2$, $\omega(t^m) = t^{-m}, \forall m \in Z, \delta = t \frac{d}{dt}.$

Proof. The result follows from Lemma 5.1 and the direct computation. \square

Theorem 5.3. *Let $\delta = t \frac{d}{dt}$, $ch F \neq 2$, $\omega_\lambda : A \rightarrow A$,*

$$\omega_\lambda(t^m) = \lambda^m t^{-m}, \lambda \in F, \lambda \neq 0, \forall m \in Z.$$

Then $(A, [, ,]_{\omega_\lambda, \delta})$ is a 3-Lie algebra in the multiplication: $\forall t^l, t^m, t^n \in L$,

$$(5.3) \quad [t^l, t^m, t^n]_{\omega_\lambda, \delta} = \begin{vmatrix} \lambda^l t^{-l} & \lambda^m t^{-m} & \lambda^n t^{-n} \\ t^l & t^m & t^n \\ lt^l & mt^m & nt^n \end{vmatrix} \\ = \lambda^l(n-m)t^{m+n-l} + \lambda^m(l-n)t^{n+l-m} + \lambda^n(m-l)t^{l+m-n}.$$

Proof. The result follows from Theorem 3.3 and Lemma 5.2. \square

Corollary 5.4. Let $\lambda = 1$ in Theorem 5.3. Then the multiplication of the 3-Lie algebra $(A, [, ,]_{\omega_1, \delta})$ is as follows: $\forall t^l, t^m, t^n \in A$,

$$(5.4) \quad [t^l, t^m, t^n]_{\omega_1, \delta} = \begin{vmatrix} t^{-l} & t^{-m} & t^{-n} \\ t^l & t^m & t^n \\ lt^l & mt^m & nt^n \end{vmatrix} \\ = (n-m)t^{m+n-l} + (l-n)t^{n+l-m} + (m-l)t^{l+m-n}.$$

$$\text{And} \quad \omega([t^l, t^m, t^n]_{\omega_1, \delta}) = -[\omega(t^l), \omega(t^m), \omega(t^n)]_{\omega_1, \delta}.$$

Theorem 5.5. Let $chF \neq 2$. Then the 3-Lie algebra $(A, [, ,]_{\omega_\lambda, \delta})$ with the multiplication (5.3) for some $\lambda \neq 0$ is isomorphic to the 3-Lie algebra $(A, [, ,]_{\omega_1, \delta})$ with the multiplication (5.4), where $\delta = t \frac{d}{dt}$.

Proof. Denote $\omega_\lambda : A \rightarrow A$, $\omega_\lambda(t^m) = \lambda^m t^{-m}$, $\lambda \in F$, $\lambda \neq 0$. Then we have $\omega_\lambda \delta + \delta \omega_\lambda = 0$.

Define $\sigma : A \rightarrow A$, $\sigma(t^m) = \lambda^{\frac{m}{2}} t^m$, $\forall m \in \mathbb{Z}$. Then

$$\sigma(t^m t^n) = \sigma(t^m) \sigma(t^n), \quad \delta \sigma(t^m) = \delta(\lambda^{\frac{m}{2}} t^m) = m \lambda^{\frac{m}{2}} t^m = \sigma \delta(t^m), \\ \omega_1 \sigma(t^m) = \omega_1(\lambda^{\frac{m}{2}} t^m) = \lambda^{\frac{m}{2}} t^{-m} = \lambda^{-\frac{m}{2}} (\lambda^m t^{-m}) = \sigma \omega_\lambda(t^m).$$

$$\delta \sigma(t^m) = \delta(\lambda^{\frac{m}{2}} t^m) = m \lambda^{\frac{m}{2}} t^m = \sigma \delta(t^m).$$

Follows from Theorem 3.7, the result holds. \square

Theorem 5.6. If $chF = p > 2$, then for every integer $k \in \mathbb{Z}$ and $k \neq 0$,

$$I_k = \{ (t^{kp} + t^{-kp})h(t) \mid \forall h(t) \in A \}, \quad J_k = \{ (t^{kp} - t^{-kp})h(t) \mid \forall h(t) \in A \}$$

are non-zero proper ideals of the 3-Lie algebra $(A, [, ,]_{\omega_1, \delta})$.

Proof. Since for every integer $k \in \mathbb{Z}$ and $k \neq 0$, I_k, J_k are ideals of the associative algebra $A = F[t^{-1}, t]$, and satisfy

$$\omega(I_k) \subseteq I_k, \quad \omega(J_k) \subseteq J_k, \quad \Delta(I_k) \subseteq I_k, \quad \Delta(J_k) \subseteq J_k.$$

By Theorem 3.5, I_k and J_k are proper ideals of the 3-Lie algebra. \square

By the above discussions, A_1 and A_{-1} are two abelian subalgebras of $(A, [,], \omega_1, \delta)$, and $A = A_1 \dot{+} A_{-1}$, where

$$A_1 = \{ p(t) \mid p(t) \in A, \omega(p(t)) = p(t) \} = \left\{ \sum_{i=r}^s a_i(t^i + t^{-i}), a_i \in F, r, s \in Z \right\},$$

$$A_{-1} = \{ p(t) \mid p(t) \in A, \omega(p(t)) = -p(t) \} = \left\{ \sum_{i=r}^s a_i(t^i - t^{-i}), a_i \in F, r, s \in Z \right\}.$$

If $B = F[t_1^{-1}, \dots, t_k^{-1}, t_1, \dots, t_k]$ is the commutative associative algebra of k variable Laurent polynomials over a field F of characteristic zero. Then for every $1 \leq j \leq k$, $\delta_j = t_j \frac{\partial}{\partial t_j}$ are derivations of B , where for every $p(t_1, \dots, t_k) = \sum_{i_1 \dots i_k} a_{i_1 \dots i_k} t_1^{i_1} \dots t_k^{i_k} \in B$, $\delta_j(p(t_1, \dots, t_k)) = \sum_{i_1 \dots i_k} i_j a_{i_1 \dots i_k} t_1^{i_1} \dots t_k^{i_k}$.

Theorem 5.7. *Let $B = F[t_1^{-1}, \dots, t_k^{-1}, t_1, \dots, t_k]$ with $chF = 0$, $\Delta_j = t_j \frac{\partial}{\partial t_j}$ be a derivation of B . For an algebra homomorphism $\omega : B \rightarrow B$, ω satisfies $\Delta_j \omega + \omega \Delta_j = 0$ and $\omega^2 = id_B$ if and only if*

$$(5.5) \quad \omega(t_1^{r_1} \dots t_j^{r_j} \dots t_k^{r_k}) = (\lambda_1^{r_1} \dots \lambda_j^{r_j} \dots \lambda_k^{r_k}) t_1^{-r_1} \dots t_j^{-r_j} \dots t_k^{-r_k},$$

where $\lambda_s \in F$, $\lambda_s \neq 0$, $r_s \in Z$, $1 \leq s \leq k$. Therefore, $(B, [,], \omega, \delta_j)$ is a 3-Lie algebra in the multiplication: $\forall t_1^{r_1} \dots t_j^{r_j} \dots t_k^{r_k}, t_1^{i_1} \dots t_j^{i_j} \dots t_k^{i_k}, t_1^{n_1} \dots t_j^{n_j} \dots t_k^{n_k} \in B$,

$$(5.6) \quad [t_1^{r_1} \dots t_j^{r_j} \dots t_k^{r_k}, t_1^{i_1} \dots t_j^{i_j} \dots t_k^{i_k}, t_1^{n_1} \dots t_j^{n_j} \dots t_k^{n_k}]_{\omega, \Delta_j} \\ = (\lambda_1^{r_1} \dots \lambda_j^{r_j} \dots \lambda_k^{r_k})(n_j - i_j) t_1^{i_1 + n_1 - r_1} \dots t_j^{i_j + n_j - r_j} \dots t_k^{i_k + n_k - r_k} \\ + (\lambda_1^{i_1} \dots \lambda_j^{i_j} \dots \lambda_k^{i_k})(r_j - n_j) t_1^{r_1 + n_1 - i_1} \dots t_j^{r_j + n_j - i_j} \dots t_k^{r_k + n_k - i_k} \\ + (\lambda_1^{n_1} \dots \lambda_j^{n_j} \dots \lambda_k^{n_k})(i_j - r_j) t_1^{i_1 - n_1 + r_1} \dots t_j^{i_j - n_j + r_j} \dots t_k^{i_k - n_k + r_k}.$$

Proof. The proof is completely similar to Theorem 5.3. □

In the following we study the 3-Lie algebra (A, ω, δ_{2k}) , where the derivation

$$\delta_{2k} = t^{2k} \frac{d}{dt} \in Der A, \quad k \in Z.$$

From Lemma 5.2, if $chF \neq 2$, for $\delta_{2k} = t^{2k} \frac{d}{dt} \in Der(F[t^{-1}, t])$, $k \in Z$, then the involution $\omega : A \rightarrow A$ satisfies

$$\delta_{2k} \omega + \omega \delta_{2k} = 0$$

if and only if ω is defined as $\omega(t^m) = (-1)^m t^m$, $\forall m \in Z$

Therefore, we have the following result.

Theorem 5.8. *If $chF \neq 2$, then A is a 3-Lie algebra in the multiplication $[\cdot, \cdot]_{\omega, \delta_{2k}} : \text{for arbitrary } t^l, t^m, t^n \in A$,*

$$(5.7) \quad [t^l, t^m, t^n]_{\omega, \delta_{2k}} = \begin{vmatrix} (-1)^l t^l & (-1)^m t^m & (-1)^n t^n \\ t^l & t^m & t^n \\ lt^{2k+l-1} & mt^{m+2k-1} & nt^{2k+n-1} \end{vmatrix} \\ = \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\} t^{2k+l+m+n-1}.$$

Proof. The result follows from Lemma 5.1. Lemma 5.2 and Theorem 3.3. \square

Define linear functions $\alpha, \beta, \gamma : A \rightarrow F$:

$$\alpha(t^m) = (-1)^m, \quad \beta(t^m) = 1, \quad \gamma(t^m) = m, \quad \forall t^m \in A,$$

that is, for every $p(t) = \sum_{i=m}^n a_i t^i \in A$,

$$\alpha(p(t)) = \sum_{i=m}^n (-1)^i a_i, \quad \beta(p(t)) = \sum_{i=m}^n a_i, \quad \gamma(p(t)) = \sum_{i=m}^n i a_i.$$

Then Eq.(5.7) can be written as: $\forall t^l, t^m, t^n \in A$,

$$[t^l, t^m, t^n]_{\omega, \delta_{2k}} = (\alpha \wedge \beta \wedge \gamma)(t^l, t^m, t^n) t^{l+m+n+2k-1},$$

$$\text{where } (\alpha \wedge \beta \wedge \gamma)(t^l, t^m, t^n) = \begin{vmatrix} \alpha(t^l) & \alpha(t^m) & \alpha(t^n) \\ \beta(t^l) & \beta(t^m) & \beta(t^n) \\ \gamma(t^l) & \gamma(t^m) & \gamma(t^n) \end{vmatrix} = \begin{vmatrix} (-1)^l & (-1)^m & (-1)^n \\ 1 & 1 & 1 \\ l & m & n \end{vmatrix}.$$

Remark References [11, 12, 9] studied n -Lie algebras $A(G, f, t)$, where G is an additive Abelian group, $f : G^n \rightarrow F$. By the above discussions, the 3-Lie algebra $(A, [\cdot, \cdot]_{\omega, \delta_{2k}})$ is isomorphic to the 3-Lie algebra $A(Z, f, 2k+1)$ in [12], where Z is the set of all integers,

$$f(l, m, n) = \begin{vmatrix} (-1)^l & (-1)^m & (-1)^n \\ 1 & 1 & 1 \\ l & m & n \end{vmatrix}, \quad \forall l, m, n \in Z.$$

Theorem 5.9. *Let $A = F[t^{-1}, t]$ be the Laurent polynomials over the field of complex numbers. Then for any integer k , $k \neq 0$, 3-Lie algebra $(A, [\cdot, \cdot]_{\omega, \delta_{2k}})$ with the multiplication (5.7) is isomorphic to the 3-Lie algebra $(A, [\cdot, \cdot]_{\omega, \delta_0})$, where $\delta_0 = \frac{d}{dt}$, and for every*

$t^l, t^m, t^n \in A$,

$$(5.8) \quad [t^l, t^m, t^n]_{\omega, \delta_0} = \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\}t^{l+m+n-1}.$$

Proof. If $k = 2s$, define linear mapping $\sigma : (A, [,],_{\omega, \delta_0}) \rightarrow (A, [,],_{\omega, \delta_k})$, $\sigma(t^m) = t^{m-k}$, $\sigma(1) = 1$, for every $t^m \in A$. Then for every $t^l, t^m, t^n \in A$,

$$\sigma([t^l, t^m, t^n]_{\omega, \delta_0}) = \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\}t^{l+m+n-k-1},$$

$$\begin{aligned} [\sigma(t^l), \sigma(t^m), \sigma(t^n)]_{\omega, \delta_{2k}} &= [t^{l-k}, t^{m-k}, t^{n-k}]_{\omega, \delta_{2k}} \\ &= \{(-1)^{l-k}(n-m) + (-1)^{m-k}(l-n) + (-1)^{n-k}(m-l)\}t^{l+m+n-k-1} \\ &= \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\}t^{l+m+n-k-1}. \end{aligned}$$

If $k = 2s + 1$, define linear mapping $\sigma : A \rightarrow A$, $\sigma(t^m) = it^{m-k}$, $\sigma(1) = 1$, where $i^2 = -1$. Then for every $t^l, t^m, t^n \in A$,

$$\begin{aligned} \sigma([t^l, t^m, t^n]_{\omega, \delta_0}) &= i \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\}t^{l+m+n-k-1}, \\ [\sigma(t^l), \sigma(t^m), \sigma(t^n)]_{\omega, \delta_{2k}} &= [it^{l-k}, it^{m-k}, it^{n-k}]_{\omega, \delta_{2k}} \\ &= -i \{(-1)^{l-k}(n-m) + (-1)^{m-k}(l-n) + (-1)^{n-k}(m-l)\}t^{l+m+n-k-1} \\ &= i \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\}t^{l+m+n-k-1} \\ &= \sigma([t^l, t^m, t^n]_{\omega, \delta_0}). \end{aligned}$$

The result holds. \square

Theorem 5.10. *Let $A = F[t^{-1}, t]$ over the field F of complex numbers. Then 3-Lie algebra $(A, [,],_{\omega, \delta_0})$ is a simple 3-Lie algebra.*

Proof. By Eq.(5.8), for every $t^l, t^m, t^n \in A$,

$$(5.9) \quad [t^l, t^m, t^{-m+1}]_{\omega, \delta_0} = \{(-1)^l(-m+1-m) + (-1)^m(l+m-1) + (-1)^{-m+1}(m-l)\}t^l \\ = \begin{cases} 0, & \text{if } l = m \text{ or } l = -m+1, \\ \{(-1)^{l+1}2m + (-1)^m2l + (-1)^l + (-1)^{m+1}\}t^l \neq 0, & \text{others.} \end{cases}$$

If $m = 0, n = 1$, then we have

$$[t^l, 1, t^1]_{\omega, \delta_0} = \{(-1)^l + (l-1) + 1\}t^l = \begin{cases} 0, & \text{if } l = 1, \\ \{2l + (-1)^l - 1\}t^l, & \text{others.} \end{cases}$$

Let I be a non-zero ideal of the 3-Lie algebra $(A, [,],_{\omega, \delta_0})$. For every non-zero vector $p(t) = \sum_{i=r}^s a_i t^i \in I$, where $a_s \neq 0, a_r \neq 0$, and m is a positive integer such that $m > s$ and $-m+1 < r$. Thanks to Eq.(5.8) and the Vandermonde determinant, we have $t^l \in I$ if $a_l \neq 0$ for $r \leq l \leq s$. We conclude that there is an integer l such that $t^l \in I$.

Now we prove $I = A$. If $t^m \in I$, then by Eq.(5.9), for every $l \neq m$ and $l \neq -m + 1$, we have $t^l \in I$. Therefore, we can choose j satisfying $j \neq \pm m$, $j \neq -m + 1$ (and then $-m + 1 \neq -j + 1$) such that $t^j \in I$. Again by Eq.(5.8), we have $t^m \in I$ and $t^{-m+1} \in I$.

Summarizing above discussions, we get $I = A$. Therefore, $(A, [,], \omega, \delta_0)$ is a simple 3-Lie algebra. \square

Theorem 5.11. *If $chF = p > 2$, then the 3-Lie algebra $(A, [,], \omega, \delta_0)$ in Theorem 5.8 is a non-simple 3-Lie algebra.*

Proof. Suppose $I_k = \{(t^{kp} + t^{-kp})h(t) \mid \forall h(t) \in A\}$, $J_k = \{(t^{kp} - t^{-kp})h(t) \mid \forall h(t) \in A\}$, $k \neq 0$. Then $\omega(I_k) \subseteq I_k$, $\delta_0(I_k) \subseteq I_k$, $\omega(J_k) \subseteq J_k$ and $\delta_0(J_k) \subseteq J_k$. Thanks to Theorem 3.4, I_k and J_k are non-zero proper ideals of the 3-Lie algebra $(A, [,], \omega, \delta_0)$. Therefore, the result holds. \square

By the above discussions, if $chF = p > 2$, then $J_1 = \{(t^p - t^{-p})h(t) \mid \forall h(t) \in A\}$ is an ideal of the 3-Lie algebra $(A, [,], \omega, \delta_0)$, and satisfies $\omega(J_1) \subseteq J_1$ and $\delta_0(J_1) \subseteq J_1$. Then we get the quotient 3-Lie algebra of $(A, [,], \omega, \delta_0)$ relating to the ideal J_1 , which is denoted by $(\bar{A}, [,], \omega, \delta_0)$. The multiplication of $\bar{A} = A/J_1$ in the basis $\bar{t}^{-p+1}, \dots, \bar{t}^{-1}, \bar{1}, \bar{t}, \dots, \bar{t}^p$ as follows

$$(5.10) \quad [\bar{t}^l, \bar{t}^m, \bar{t}^n]_{\omega, \delta_0} = \{(-1)^l(n-m) + (-1)^m(l-n) + (-1)^n(m-l)\} \bar{t}^{l+m+n-1},$$

where $\bar{t}^p = \bar{t}^{-p}$.

Theorem 5.12. *The 3-Lie algebra $(\bar{A}, [,], \omega, \delta_0)$ is a simple 3-Lie algebra, where $[,], \omega, \delta_0$ is defined as Eq. (5.10) and $\dim \bar{A} = 2p$.*

Proof. Let \bar{I} be a nonzero ideal of the 3-Lie algebra $(\bar{A}, [,], \omega, \delta_0)$. Suppose $h(\bar{t}) = \sum_{i=1-p}^p a_i \bar{t}^i \in \bar{I}$ and $h(\bar{t}) \neq 0$.

Case I. If $h(\bar{t}) = \sum_{i=1-p}^p a_i \bar{t}^i = \bar{t}^p$. For every l satisfying $1-p < l < p$, since

$$[\bar{t}^l, \bar{t}^p, \bar{t}^{1-p}] = \{(-1)^l - 2l + 1\} \bar{t}^l \in \bar{I}, \quad (-1)^l - 2l + 1 \neq 0, \quad l \neq 1-p,$$

we get $\bar{t}^l \in \bar{I}$ for $1-p < l \leq p$.

Thanks to $p \geq 3$, $\bar{t}^2 \in \bar{I}$, then $-2\bar{t}^{p-1} = [\bar{t}^2, \bar{t}^{-1}, \bar{t}^{1-p}] \in \bar{I}$. It follows $\bar{I} = \bar{A}$.

Case II. If $h(\bar{t}) = \sum_{i=1-p}^p a_i \bar{t}^i \in \bar{I}$ satisfies $a_p \neq 0$, and there is an integer k satisfying $1-p \leq k < p$ and $a_k \neq 0$. Without loss of generality, we suppose $a_p = 1$. By Eq.(5.10)

$$[h(\bar{t}), \bar{t}^{p-1}, \bar{t}^{2-p}] = \sum_{i=1-p}^p a_i (3(-1)^i + 2i - 1) \bar{t}^i \in \bar{I}.$$

Since $3(-1)^i + 2i - 1 = 0$ if and only if $i = p - 1$ or $i = 2 - p$, and $3(-1)^i + 2i - 1 = 3(-1)^j + 2j - 1$ if and only if $i = j$ for $i \neq p - 1, i \neq 2 - p, j \neq p - 1, j \neq 2 - p$, we obtain $\bar{t}^p \in I$ (using Vandermonde determinant). Follows the discussions of the Case I, $\bar{I} = \bar{A}$.

Case III. If $h(\bar{t}) = \sum_{i=1-p}^p a_i \bar{t}^i = \sum_{i=1-p}^s a_i \bar{t}^i$, where $s < p$, $a_s = 1$ (that is, $a_p = a_{p-1} = \dots = a_{s+1} = 0$) and there is an integer k satisfying $1 - p \leq k < s$ such that $a_k \neq 0$. Then $s > 1 - p$ and

$$[h(\bar{t}), \bar{t}^p, \bar{t}^{1-s}] = \sum_{i=1-p}^s a_i \{(-1)^i(1-s) - i - s + 1 + i(-1)^s\} \bar{t}^{i+p-s} = \sum_{j=1-s}^p b_j \bar{t}^j \in \bar{I},$$

where $b_i = a_i((-1)^i(1-s) - i - s + 1 + i(-1)^s)$. We obtain

$$b_p = (-1)^s(1-s) - s - s + 1 + s(-1)^s = -2s + (-1)^s + 1 \neq 0 \text{ since } 1 - p < s < p.$$

Follows from Case II, $\bar{I} = \bar{A}$.

Case IV. If $h(\bar{t}) = \sum_{i=1-p}^p a_i \bar{t}^i = \bar{t}^l$, where $l < p$. If $l < p - 1$, then

$$[\bar{t}^l, \bar{t}^p, \bar{t}^{l+1}] = \{(-1)^l(2l+1) + 1\} \bar{t}^p \in \bar{I}, \text{ and } (-1)^l(2l+1) + 1 \neq 0, \text{ we obtain } \bar{t}^p \in \bar{I}.$$

If $l = p - 1$, then $[\bar{t}^{p-1}, \bar{t}^p, \bar{t}^{-p+2}] = 4\bar{t}^p \in \bar{I}$. Therefore, $\bar{t}^p \in \bar{I}$.

Summarizing above discussions, $\bar{I} = \bar{A}$. It follows the result. \square

6. CONCLUSIONS AND DISCUSSIONS

Since the multiple multiplication, constructions of n -Lie algebras is a continuously difficult problem in the structure theory of n -Lie algebras, for $n \geq 3$.

In [3, 10, 11, 12], n -Lie algebras are realized by associative commutative algebras and its arbitrary n pairwise commuting derivations, and linear functions.

Papadopoulos in ([7]) constructed 3-Lie algebras by Dirac γ -matrices. Let A be spanned by the four-dimensional γ -matrices (γ^μ) and let $\gamma^5 = \gamma^1 \dots \gamma^4$. Then the product

$$(6.1) \quad [x, y, z] = [[x, y]\gamma^5, z], \quad \forall a, b, c \in A.$$

defines a 3-Lie algebra which is isomorphic to the unique simple 3-Lie algebra ([3]).

In [17] 3-Lie algebras are constructed from metric Lie algebras. Let (\mathfrak{g}, B) be a metric Lie algebra over a field \mathbb{F} , that is, B is a nondegenerate symmetric bilinear form on \mathfrak{g} satisfying $B([x, y], z) = -B(y, [x, z])$ for every $x, y, z \in \mathfrak{g}$. Suppose $\{x_1, \dots, x_m\}$ is a basis of \mathfrak{g} and $[x_i, x_j] = \sum_{k=1}^m a_{ij}^k x_k$, $1 \leq i, j \leq m$. Set

$$(6.2) \quad \mathfrak{g}_0 = \mathfrak{g} \oplus \mathbb{F}x^0 \oplus \mathbb{F}x^{-1} \quad (\text{the direct sum of vector space}).$$

Then there is a 3-Lie algebra structure on \mathfrak{g}_0 given by

$$(6.3) \quad [x_0, x_i, x_j] = [x_i, x_j], \quad 1 \leq i, j \leq m; \quad [x^{-1}, x_i, x_j] = 0, \quad 0 \leq i, j \leq m;$$

$$(6.4) \quad [x_i, x_j, x_k] = \sum_{s=1}^m a_{ij}^s B(x_s, x_k) x^{-1}, \quad 1 \leq i, j, k \leq m.$$

And A is a metric 3-Lie algebra in the multiplication (6.4) and (6.5).

In [14], 3-Lie algebras are realized by Lie algebras and linear functions. Let $(L, [,])$ be a Lie algebra, $f \in L^*$ satisfying $f([x, y]) = 0$ for every $x, y \in L$. Then L is a 3-Lie algebra in the multiplication

$$(6.5) \quad [x, y, z]_f = f(x)[y, z] + f(y)[z, x] + f(z)[x, y], \quad \forall x, y, z \in L.$$

And it is proved in [14] that every m -dimensional 3-Lie algebras can be obtained by the multiplication (6.2) and (6.6) for $m \leq m$.

Awata, Li and et al in [18] constructed a 2-step solvable 3-Lie algebra from $(n \times n)$ -matrices. Let $\mathfrak{g} = gl(m, \mathbb{F})$ be the general linear Lie algebra. Then there is a 3-Lie algebra structure on \mathfrak{g} defined by

$$(6.6) \quad [A, B, C] = (tr A)[B, C] + (tr B)[C, A] + (tr C)[A, B], \quad \forall A, B, C \in \mathfrak{g}.$$

In this paper we construct 3-Lie algebras by associative commutative algebras and their derivations and involutions. From Example 4.1 and 4.2 we can obtain 3-Lie algebras from any $\alpha \in Hom(G^+, F^+)$ which is not isomorphic to the 3-Lie algebra obtain by Eq.(6.7), where G is the set of all $(n \times n)$ -matrices over a field F .

So we may provide a problem that is how can we realize 3-Lie algebras by Lie algebras, and associative commutative algebras with multilinear functions and general linear mappings.

ACKNOWLEDGEMENTS

The first author (R. Bai) was supported in part by the Natural Science Foundation of Hebei Province (A2010000194).

REFERENCES

- [1] Y. Nambu, Generalized Hamiltonian dynamics, *Phys. Rev. D* 7 (1973) 2405-2412.
- [2] L. Takhtajan, On foundation of the generalized Nambu mechanics, *Comm. Math. Phys.* 160 (1994) 295-315.
- [3] V.T. Filippov, n -Lie algebras, *Sib. Mat. Zh.*, 26 (1985) 126-140.
- [4] J. Bagger and N. Lambert, Modeling multiple $M2$ s, *Phys. Rev. D* 75 (2007) 045020 [hep-th/0611108]; Gauge symmetry and supersymmetry of multiple $M2$ -branes, *Phys. Rev. D* 77 (2008) 065008 [arXiv:0711.0955]; Comments on multiple $M2$ -branes, *JHEP* 02 (2008) 105 [arXiv:0712.3738].
- [5] P. Ho, R. Hou and Y. Matsuo, Lie 3-algebra and multiple M_2 -branes, arXiv: 0804.2110.

- [6] P. Ho, M. Chebotar and W. Ke, On skew-symmetric maps on Lie algebras, *Proc. Royal Soc. Edinburgh A* 113 (2003) 1273-1281.
- [7] A. Gustavsson, Algebraic structures on parallel M2-branes, arXiv: 0709.1260.
- [8] G. Papadopoulos, M2-branes, 3-Lie algebras and Plucker relations, arXiv: 0804.2662.
- [9] V.T. Filippov, On n -Lie algebra of Jzcpbians, *J. Siberian Mathe.*, 39(3) (1998): 576-581.
- [10] A.S. Dzhumadil'daev, Identities and derivations for Jacobian algebras, arXiv: 0202040v1[math.RA]
- [11] A. P. Pozhidaev, Monomial n -Lie algebras, *Algebra and Logic*, 37(5) (1998): 307-322.
- [12] A. P. Pozhidaev, Simple n -Lie algebras, *Algebra and Logic*, 38(3) (1999): 181-192.
- [13] R. Bai, Y. Wu, J. Li and H. Zhou, Constructing $(n + 1)$ -Lie algebras from n -Lie algebras, *J. Phys. A: Math. Theor.* 45 (2012)
- [14] R. Bai, C. Bai and J. Wang, Realizations of 3-Lie algebras, *J. Math. Phys.* 51, 063505 (2010). 475206.
- [15] S. Kasymov, On a theory of n -Lie algebras, *Algebra i Logika* 26 (1987) 277-297.
- [16] J.T. Hartwig, D. Larsson and S. Silvestrov, Deformations of Lie algebras using σ -derivations, arXiv: 5036 2003
- [17] P. Ho, Y. Imamura, Y. Matsuo, M2 to D2 revisited, *J. High Energy Phys.* 0807,003 2008; e-print arXiv:0805.1202.
- [18] H. Awata, M. Li, D. Minic and T. Yoneya, On the quantization of Nambu brackets, *JHEP* 2 (2001) 013 (17pp).

COLLEGE OF MATHEMATICS AND COMPUTER SCIENCE, HEBEI UNIVERSITY, BAODING 071002,
P.R. CHINA

E-mail address: bairuipu@hbu.cn

COLLEGE OF MATHEMATICS AND COMPUTER SCIENCE, HEBEI UNIVERSITY, BAODING 071002,
P.R. CHINA

E-mail address: wuyg1022@sina.com