



[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Farmers' Innovation](#) | [Publications](#) | [Contact](#)

Crop protection :: IPM :: Chickpea

Monitoring for Pest and Disease

- Undertake community level monitoring to know change in distribution and abundance of pest.
- Organise regular pest monitoring and assess bio control potential at every 10 days interval.
- Record pest incidence at every 5 to 10 km distance at 12 spots/ha and 5 plants/spot selected randomly.
- Use pheromone traps for monitoring gram pod borer. Install pheromone traps at a distance of 50 m @ 5 traps/ha for each insect pest.
- Use specific lure for each pest species and change the lure after every 15-20 days. Trapped moths should be removed daily.

Pre-sowing stage

- Deep ploughing in summer.
- Apply well decomposed FYM or neem cake
- Synchronised sowing single recommended variety in village/area
- Solar treatment to the soil using transparent polythene mulching for at least for 15 days.
- Removal of crop stubbles
- Inter cropping with coriander/mustard/wheat/sorghum (Rabi)
- Early planting mid October to escape the peak activity of *H. armigera*
- Marigold plantation should be adopted as trap crop/antagonistic crop for nematode.
- Use tolerant/resistant varieties

Sowing date

- Use Rhizobium culture @ 1 pkt +(200g), 190 kg seed for effective modulation
- Thinning should be done in case of dense plant population
- Seed treatment with Trichoderma @ 4 gm/kg of seed.
- Seed treatment with Carbendazim 25% DS@ 0.75 gm./ kg. of seed.
- Seed treatment with Carbosulfan 25 EC @ 3%.
- Follow recommended agronomic practices for land preparation seed ratio proper and balanced fertilizer crop stand.

Vegetative stage

- Inter culture and band weeding for keeping the crop weed free for 6 to 8 weeks.
- Detopping to reduce pest occurrence
- Regular monitoring of gram pod borer in south zone during entire crop season.
- Conservation of predatory wisp, spiders, insect parasitoids etc.

Flowering stage

- NPV 250 LE or B.t. var Kurstaki 1 kg/ha NSKE 5% or neem pesticides.
- Conservation of predators like spiders, chrysopa and other natural enemies by avoiding indiscriminate use of pesticides.
- Installing of bird perches for predatory birds.

Fruiting stage

- NPV 250 LE or B.t. var Kurstaki 1 kg/ha NSKE 5% or neem pesticides.
- Conservation of predators like spiders, chrysopa and other natural enemies by avoiding indiscriminate use of pesticides.
- Installing of bird perches for predatory birds.

[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Publications](#) | [Contact Us](#)

© TNAU 2008-2024 All Rights Reserved.