



[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Farmers' Innovation](#) | [Publications](#) | [Contact](#)

Crop protection :: IPM :: Red gram

Monitoring for Pest & Disease

- Under take survey at every 10 km distance initially at weekly intervals and there after at 10 days intervals.
- Record incidence of pod borer on all host crops of the locality.
- Field scouting for pests and bio control fauna by extension agencies and farmers once in 7 days should be undertaken to workout ETL.
- Two hours light exposure up to 8-8.30 p.m. for pest monitoring through light trap should be undertaken in the field if possible.
- Install pheromone traps at a distance of 50 m @ 5 traps/ha.
- Use helioverpa lure and change it after every 20 days. Trapped should be removed daily.

Pre-sowing stage

- Apply FYM or neem cake.
- Synchronise sowing with a single variety in a village/area
- Inter crop with sorghum for reducing Helicoverpa, wilt and nematode incidence.
- Early planting (mid-June) in NWPZ for avoiding peak infestation period of *H. armigera* on short duration pigeonpea.
- Resistant/tolerant varieties.
- Cultivating early maturing varieties.
- Planting of castor or tall sorghum/maize varieties on borders for conserving natural enemies. These plants also function as live perches for predatory birds.
- Plant short stature crop like cowpea, mungbean, urdbean, fodder soybean etc. in 1 m wide band after 8-10 rows of pigeonpea.

Sowing stage

- During visual observation, if wilt and blight incidence is above 20%, the field may be considered as sick and growing pigeon pea should be discouraged there.
- Growing trap crop like marigold on the borders and in between rows as inter crop Their flowers shall attract oviposition which can then be plucked and disposed.
- Ridge Planting + cover crops like soybean or cowpea.
- Treatment of the seed with *Trichoderma viride* @ 4 grams/kg seed.
- Seed treatment with carbendazim + thiram* (1 gm + 2 gm/kg seed, or carbendazim (2 gm/kg seed) or thiram (3 gm/kg seed).
- Seed treatment with metalaxy1 for phytophthora blight.
- Soil treatment with carbofuran granule @ 8 kg a.i./ha at planting or seed treatment.

Vegetative stage

- Inter-culture and hand weeding for keeping the crop, weed free for 6-8 weeks.
- Hand picking, jarring on cloth or in bags is effective in a limited way in small areas.
- Conservation of predatory spiders and wasps etc.
- Installation of bird perches for the predatory birds.

Flowering stage

- Removal and destruction of sterility mosaic affected plants.
- Hand picking and jarring on cloth or in bags is effective in a limited way in small areas.
- Conservation of predatory spiders and wasps etc.
- Installation of bird perches for the predatory birds.
- Application of NPV @ 250 LE per ha. on noticing eggs and 1st instars larvae (2-3 eggs or 1 larva per twigs).
- B.t Spray@ 0.07-1.0 kg/ha
- Spray of NSKE 5% at pre-flowering stage 3 times at 15 days interval.
- Spraying of 0.07% endsulfan (2ml of 35 EC/lit of water) or chlorpyrifos 20 EC @ 3.5 ml./lit. of water at 600-1000 lit of spray material per ha. with hand sprayer or 200-300 lit/ha with power sprayer.

Fruiting stage

- HANPV @ 250 LE per ha. on noticing eggs and 1st instars larvae (2-3 eggs or 1 larvae/5 twigs)
- Spraying the crop with chlorpyrifos 20 EC 3.5 ml/lit. of water or only one time fenvalerate 75 gm a. i./ha. in 600 to 1000 of spray liquid per ha.

Storage

- Make harvest at the correct maturity time

[Home](#) | [About Us](#) | [Success Stories](#) | [Farmers Association](#) | [Publications](#) | [Contact Us](#)

© TNAU 2008-2024 All Rights Reserved.