भारत सरकार / **Government of India**

कृषि एवं किसान कल्याण मंत्रालय / **Ministry of Agriculture & Farmers Welfare**

कृषि एवं किसान कल्याण विभाग

Department of Agriculture & Farmers Welfare

वनस्पति संरक्षण, संगरोध और संग्रह निदेशालय

Directorate of Plant Protection, Quarantine & Storage

केंद्रीय कीटनाशी बोर्ड एवं पंजीकरण समिति

Central Insecticide Board & Registration Committee

एन. एच. - 4, फरीदाबाद001 121 – (हरियाणा)

N.H.-IV, Faridabad-121 001 (Haryana)

MAJOR USES OF PESTICIDES

(Registered under the Insecticides Act, 1968)

(UPTO – 30.09.2026)

(Based on certificate issued)

***Disclaimer**: The document has been compiled on the basis of available information for guidance and not for legal purposes.*

INSECTICIDES

1. **Insecticides registered for Agriculture use (Page No. – 02)**
2. **Insecticides combination registered for agriculture use (Page No. – 53)**
3. **Insecticides registered for Public Health use (Page No. – 85)**
4. **Insecticides registered for Household use (Page No. – 90)**
5. **Recommended chemicals by FAO for Locust Control (Page No. – 105)**

Approved Uses of Registered Insecticides.

Agricultural Use					
Crop	**Common Name of the pest**	**Dosage/ha**			**Waiting Period (days)**
		a.i (gm)	**Formulation (gm/ml)**	**Dilution in Water (Liter)**	
Abamectin 01.90 % EC					
Rose (Ornamental)	Red spider mites (*Tetranychus urticae*)	0.00048-0.00096%	0.025-0.050%	500	03
Grapes	Mites	0.014/L	0.75 ml/L water	500 – 1000	03
Apple	European Red Mite and Spotted Red Spider Mite	0.00095%	0.05%	6-7 litre water/ tree	7
Tomato	Red spider mites and Leaf Miner	8.6 – 11.4	450 - 600	500	3
Acequinocyl 15% w/v SC					
Rose	Two spotted mites (Tetranychus urticae)	75g	500 ml	500L	10 days
Acephate 75 % SP					
Cotton	Jassids	292	390	500 – 1000	15
	Bollworms	584	780	500 – 1000	15
Safflower	Aphids	584	780	500 – 1000	15
Rice (Paddy)	Yellow stem borer, Leaf folder, Plant Hoppers, Green leafhopper	500 – 750	666 – 1000	300 – 500	15
Acephate 97 % DF					
Cotton	Jassids & Boll wormcomplex	436.50 – 582	450 – 600	500	48
Paddy (Rice)	Yellow stem borer,Leaf folder, Plant hoppers, Green leafhopper	727.50	750	500	21
Acephate 95 % SG					
Rice (Paddy)	Stem borer, Leaf folder, Brown planthopper	562.50	592	500	30
Cotton	Jassids	750	790	500	18

Chilli	Thrips, Fruit borer (*Helicoverpa armigera*), Aphid	750	790	500	07
Cardamom	Thrips, Shoot, Panicle & Capsule borer	950	1000	1000	14
Acetamiprid 20 % SP					
Cotton	Aphids, Jassids	10	50	500 – 600	15
	Whiteflies	20	100	500 – 600	15
Cabbage	Aphids	15	75	500 – 600	07
Okra (Bhindi)	Aphids	15	75	500 – 600	03
Chilli	Thrips	10 – 20	50 – 100	500 – 600	03
Rice (Paddy)	Brown plant hopper	10 – 20	50 – 100	500 – 600	07
Tea	Tea Mosquito bug (*Helopeltis theivora*)	24	120	400	07
Acetamiprid 25% + Bifenthrin 25 % WG					
Cotton	Jassid, aphids, thrips, whiteflies, *Pectinophora gossypiella*, *Helicoverpa armigera*, *Earias vitella*	80	160	500	33
Soybean	Whitefly, girdle beetle, semi looper and tobacco caterpillar	125	250	500	28
Acynonapyr 18.19 % w/w SC					
Tea	Red Spider Mite	100-150	500-750	500	7
Afidopyropen 50 g/L DC					
Brinjal	Whitefly, Jassids	50	1000	500 – 750	01
Cotton	Whitefly, Jassids	50	1000	500 – 750	25
Cucumber	Whitefly	35 – 50	700 – 1000	500	05
Alphacypermethrin 10.00% EC					
Cotton	Boll Worms	15 – 25	165 – 280	600 – 1000	07

Alphacypermethrin 10.00% SC					
Cotton	Boll Worms	25 – 30	250 – 300	500 – 1000	10

Name of Commodity	**Common name of the pest**	**Dose**	**Exposure Period**	**Aeration Waiting period**
Aluminum Phosphide 56 % (3g Tablet, 10g Pouch)				
Stored Whole Cereals and Seed Grains Millet, Pulses Dry Fruits, Nuts Spices & Oil Seeds	Rice Weevil (*Sitophilus oryzae*),Lesser Grain Borer, Khapra Beetle (*Trogoderma granarium*), Rust Red Flour Beetle, Saw Toothed GrainBeetle, Caddle Beetle, Drug Store Beetle, Cigarette Beetle, Pulse Beetle	03 tablets (03 gm) per ton or 150 gm per 100m^3 or 10 gm Pouch Per ton of commodity or 150 gm per 100 m^3.	Minimum 05 Days (*Sitophilus oryzae*) or 07 Days (*Trogoderma granarium*)	One hour of partial aeration in case non-polyethylene packed commodities allowed by 6-8 hrs of full aeration. For polyethylene packed commodities minimum aeration period is 48 hrs. The waiting period for the release of stock is 48hrs in both the cases. Recommendation for bag stock 15 days.
Mild Products: De-oiled Cakes, Rice Bran Flour, Grain Animal & Poultry Food Split Pulses (Dal) & other Processed Food	Long Headed Floor Beetle, Coffee Borer,Dried Fruit Beetle, Flat Grain Beetle, Carpet Beetle	03 tablets/10 gm per ton or225 gm/100 m^3	05 days	Aeration is waiting Period07 days to be checked PH^3 detector strips.
Empty Godowns & Sheds	Rice Moth, Almond Moth, Mites, Fruit Fly, Granary Weevil, Caddle or Flour worm, Red Flour Beetle, Indian Meal Moth, Larger cabinetMoth, Wheat Kernel Damage in the field Cockroach.	14 tablets/1000 m^3 or 150 gm/100 m^3 or 4pouch 10 gms each/1000 CFTor 150 gm/100 $_m3$	72 hrs.	Aeration Period24 hrs detectorstrips or 4hosphine detect tubes should be used in the premises to signal safety of atmosphere.
Rodents Burrows	Rodents	01 Tablet / Burrow	-	-
Aluminum Phosphide 15 % (12g Tablet)				

Stored whole cereals and seed grains.	Rice weevil, Rust redflower beetle	1 tablet (12 g)per ton or 600100 m^3	Non polythene Packed commodities: Partial-1 hour. Full-(6-8) hour. Polythene Packed commodities: Minimum 48 hrs.	07-14
Millets, pulses, dry fruits, nuts, spices & oilseeds (Air tight cover or godowns)	Lesser Grain Borer, Khapra Beetle, Saw Toothed Grain Beetle, Rice Moth, Almond Moth	900 g/100 m^3	-	05
Milled products: De-oiled cakes, Rice bran	Rust red flower beetle	3 tablets/ton	48 hrs.	05
Flour Suji meals and Crushed grain (Animal & poultry feed), Split Pulses Dals)	Saw Toothed Grain, Beetle , Rice Moth, Almond Moth, long headed flour beetle & Mites	900 g/100 m3	48 hrs.	03
Other processed food and Empty Godowns & Sheds (under air tight condition)	All insect pests.	14 tablets/1000 tons or 600 g/1000 m^3	48 hrs. 24 hrs.	03
Aluminium Phosphide 77.50 % GR				
Stored Grain	Red Rust Flour Beetle, Lesser Grain Borer, Rice Weevil, Khapra Beetle	3.35 gm	07 days	24 hours
Aluminum Phosphide 06 % Tablet				
Crop & Non-Crop area	Field rodents	0.72 g a.i./burrow	One tablet of 12gm/burrow	-
Barium Carbonate 1% P				
Godowns, Residential Premises, Public halls	Rats, Mice, Field Rodents	10-20% Technical material to be mixed with bait	-	-	-
Beta-cyfluthrin 02.45 % SC				
Cotton	Bollworm	12.5-18.75	500 – 750	500 – 1000	20
Benfuracarb 03 % GR				

Rice (Paddy)	Stem borer, Leaf folder, Brown plant hopper	1000	33000	-	20
Benfuracarb 40 % EC					
Red gram (Tur or Arhar)	Pod borer	1000	2500	500	20
Benzpyrimoxan 10% SC					
Rice	Brown Plant Hopper, White Backed Plant Hopper	75-100	750-1000	500	31
Bifenazate 50 % WP					
Rose	Two Spotted Mite (*Tetranychus urticae*)	375	750	3000	-
Bifenazate 22.60 % SC					
Rose	Two Spotted Mite (*Tetranychus urticae*)	120	500	2000	-
Grapes	Mites	120	500	1000	15
Bifenthrin 08 % SC					
Tea	Red spider mite, Tea Mosquito bug	40.00	500	400	11
Apple	Mites	60 gm (0.006% Conc.)	7.50 ml/tree	10 lit/tree	21
Bifenthrin 08.80 % CS					
Rice (Paddy)	Stem borer, Leaf folder	44	500	500	21
Bifenthrin 10 % EC					
Cotton	Bollworms, Whitefly	80	800	500	15
Rice (Paddy)	Stem borer, Leaf folder, Green leaf hopper	50	500	500	21
Sugarcane	Termites	100	1000	500	300
Bifenthrin 02.50 % EC					
Pre and post construction: Bifenthrin 2.5% EC shall be applied at 0.05% a.i. conc. i.e. 20.0 ml formulated product diluted in 1 liter of water for the control of termites in building during pre and post construction. Treatment should be as per IS 6313 (Part 2):2001 for pre construction chemical treatment and IS 6313 (Part-3): 2001 for post construction treatment of the existing building. Recommendation for use of control of Wood borer (Powder Post Beetle) in plywood, veneer and wood					

Use	Method of application	Dosage (a.i.)	Dilution
Plywood	Glue Line Poisoning	10 g/ meter3 of wood	400 ml formulation per meter3 of wood
	Dipping	0.025% Solution	Mix 01 lit of formulation in 99 lit of waterto make 0.025% Solution
Veneer	Dipping	0.025% Solution	Mix 01 lit of formulation in 99 lit of waterto make 0.025% Solution
Wood	Dipping /Brushing	0.025% Solution	Mix 01 lit of formulation in 99 lit of waterto make 0.025% Solution

Brodifacoum 0.005 %w/w BB

Pest	Dose rate	Manner of application /use pattern
Field rats/Bandicootrats (*Bandoicota bengalensis; B. indica*) Indian houserat / Black, Indian house rat/black rat/roof rat (*Rattus rattus; R. meltade*),	One bait of 0.005% (a block of 20 gm each) per baiting station as a single feed	In and around premises (Residential, commercial, institutional, industrial public service premises, cold storage , Godowns, ware house, municipal locations, grain mandis, crop store rooms, burrow baiting, livestock rearing facilities, damp premises such as sewer etc.)
House mouse/ Fieldmouse (*Mus musculus*)		

Broflanilide 300 g/l SC					
Chilli	Fruit borer (*Helicoverpa armigera*)	12.6- 18.6	42-62	500	1
	Thrips (*Scirtothrips dorsalis*)	18.6- 25.2	62-84	500	1
Brinjal	Shoot and fruit borer (*Leucinodes orbonalis*)	12.6- 18.6	42-62	500	1
	Thrips (*Thrips tabaci*)	18.5- 25.2	62-84	500	1
Tomato	Fruit borer (*Helicoverpa armigera*) & leaf miner (*Liriomyza trifolii*)	18.6-25.2	62-84	500	1
Soybean	*Helicoverpa armigera*, *Spodoptera litura*, Semilooper (*Chrysodeixis acuta*)	12.6-18.6	42-62	500	37
Red gram	*Helicoverpa armigera* & *Maruca vitrata*	12.6-18.6	42-62	500	25
Chilli	Thrips (*Scirtothrips dorsalis*)	18.6 - 25.2	42-62	500	1
	Fruit and Shoot Borer (*Helicoverpa armigera*)	12.6 – 18.6	62-84		
Broflanilide 20% SC					
Brinjal	Shoot and fruit borer (*Leucinodes orbonalis*), Thrips (*Thrips tabaci*) & Jassids (*Amrasca devastans*)	25	125	500	1
Cabbage	Diamond back moth (*Plutella*	25	125	500	1

	xylostella) & Tobacco caterpillar (*Spodoptera litura*)				
Chilli	Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura*), Thrips (*Thrips tabaci*) and Jassids (*Amrasca devastans*)	25	125	500	1
Okra	Fruit borer (*Helicoverpa armigera*), Thrips (*Thrips tabaci*) and Jassids (*Amrasca devastans*)	25	125	500	1
Maize	Fall Army Worm	25	125	500	29
Cotton	Thrips & Jassids	25	125	500	10
Pomegranate	Thrips & Fruit borer	25	125	500	1
Soybean	Pod borer, Semi-looper & Tobacco Caterpillar	25	125	500	23
Tomato	Thrips, Leaf miner & Fruit borer	25	125	500	1
Bromadiolone 00.25 % CB					
Paddy (Rice)	Field Rat, Large Bandicota Indian house rat, Indian field mouse	0.005	-	-	-
Wheat , Gram	Field Rat, Indian house rat	0.005	-	-	-
Groundnut, Sugarcane	Field Rat, LargeBandicota	0.005	-	-	-
Coconut/ Bamboo	Indian house rat	0.005	-	-	-
Residential premises	Field Rat, LargeBandicota	0.005	-	-	-
Poultry Farm	Indian house rat,House mouse	0.005	-	-	-
Bromadiolone 00.005 % RB					
Paddy (Rice)	Field Rat, Large Bandicota, Indianhouse rat	0.005	-	-	-
Wheat	Indian Field mouse,Field Rat	0.005	-	-	-
Gram	Indian house rat,Field Rat, Indianhouse rat	0.005	-	-	-
Groundnut,Sugarcane	Field Rat, LargeBandicota	0.005	-	-	-
Coconut/ Bamboo	Indian house rat,Field Rat Large Bandicota	0.005	-	-	-
Residentialpremises	Indian House rat,House mouse	0.005	-	-	-

Poultry Farm	Indian house rat, House mouse, LargeBandicota	0.005	-	-	-

Buprofezin 25 % SC					
Cotton	Whitefly Aphids, Jassids, Thrips	250.0	1000	500 – 750	20
Chilies	Yellow Mite	75.0-150.0	300-600	500 – 750	05
Mango	Hoppers	0.025% - 0.05%	1-2 ml/liter of water	5-15 liter per tree	20
Grapes	Mealy bugs	250 – 375	1000 – 1500	500 – 1000	07
Rice	Brown plant hopper, Green leaf hopper, White Back Plant Hopper	200	800	400 – 500	20
Buprofezin 70 % DF					
Okra (Bhindi)	Jassids	200	286	500	05
Cotton	Jassids, Whitefly	250 – 300	357 – 429	500	20
Rice	Brown plant hopper	175	250	500	24
Carbofuran 03 % CG					
Barley	Aphid, Cystnematode	1000	33300	-	-
	Jassids	1250	41600	-	-
Bajra	Shoot fly	1500	50000	-	-
Sorghum	Shoot fly	1000	33300	-	-
	Stem borer	250	8300	-	-
Jute	Nematodes	1000	33300	-	-
Groundnut	Pod borer	1500	50000	-	-
	White grub	1000	33300	-	-
French bean	White grub	700	23300	-	-
Potato	Aphid	500	16600	-	-
	Jassids	1000	33300	-	-
Tomato	Whitefly fly	1200	40000	-	-
Apple	Woolly aphid	05/tree	166/tree	-	-
Citrus	Nematode	360	12000	-	-
	Leaf miner	1500	50000	-	-
Maize	Stem borer, Shootfly, Thrips	1000	33300	-	-
Paddy (Rice)	Brown plant hopperGall midge, Stem borer, Green leaf hopper, Hispa	750	25000	-	-
	Nematodes	1500	50000	-	-

Mustard	Mustard leaf miner	2000	66600	-	-
	Whitefly	1000	33300	-	-
Soybean	Root knot nematode	1500	50000	-	-
Sugarcane	Top borer	2000	66600	-	-
Bhindi (Okra)	Jassids	1000	33300	-	-
Chilli	Aphid , Thrips	1000	33300	-	-
Cabbage	Nematode	1000	50000	-	-
Wheat	Ear cockle nematode	3000	10000	-	-
	Cereal cyst nematode	2000	66600	-	-
Brinjal	Root knot nematode,Reni form nematode	2000	66600	-	-
Banana	Rhizome weevil	01 g/ suckers	33 g/sucker	-	-
	Aphid	50 g/ suckers	166 g/sucker	-	-
	Nematode	1.5 g/suckers	50 g/suckers	-	-
Peach	Leaf curl aphid	1000	33300	-	-
Mandarins	Soft greens scale	0.40 g/plant	13.30 g/plant	-	-
French bean	White grubs	750	23300	-	-
	Grey & Stem weevil	1000	33300	-	-
Pea	Shoot fly & Aphid	1000	-	-	-
Tea	Cock chafer grub	0.30 g/plant	33.10 g/plant	-	-
Carbosulfan 06 % Granules					
Rice (Paddy)	Stem borer, Gall midge, Green leaf hopper, Leaf folder	1000	16700	-	37
Carbosulfan 25% EC					
Rice (Paddy)	Green leaf hopper, White Back Plant Hopper, Brown plant hopper, Gall midge, Stem borer, leaf folder	200 – 250	800 – 1000	500 – 1000	14
Chilli	White aphid	200 – 250	800 – 1000	500 – 1000	08
Cumin	Aphid, Thrips	312.5	1250	500	17
Brinjal	Fruit and Shoot borer	312.5	1250	500	5
Cotton	Aphid, Thrips	312.5	1250	500	70
Carbosulfan 25 % DS					
Cotton	Jassid, Aphids,Thrips	15 gm/kg seed	60 gm/kg seed	Not required	-

Cartap Hydrochloride 04 % Granules					
Rice (Paddy)	Stem borer	750.0	18750	-	-
	Leaf folder, Whorl maggot	750-1000	18750-25000	-	-
Cartap Hydrochloride 50 % SP					
Rice (Paddy)	Stem borer, Leaffolder	500	1000	500 – 1000	21
Cartap Hydrochloride 75 % SG					
Rice	Yellow stem borer,Leaf folder	318.75 – 375	425 – 500	250 – 500	35-89
Chlorantraniliprole 18.50 % SC					
Rice	Stem borer, Leaf folder	30	150	500	47
Cabbage	Diamond back moth	10	50	500	03
Cotton	American bollworm, Spotted bollworm, Tobacco caterpillar	30	150	500	09
Sugarcane	Termite	100 – 125	500 – 625	1000	208
	Early shoot borer, Top borer	75	375	1000	208
Tomato	Fruit borer	30	150	500	03
Chilli	Fruit borer and Tobacco caterpillar	30	150	500	03
Brinjal	Shoot & Fruit borer	40	200	500 –750	22
Pigeon pea	Gram Pod borer and Pod Fly	30	150	500 –750	29
Soybean	Green Semi looper, Stem fly, Girdle beetle and Tobacco caterpillar	30	150	500 –750	22
Bengal gram	Pod borers	25	125	500	11
Black gram	Pod borers	20	100	500	20
Bitter gourd	Fruit borers & Leaf Caterpillars	20 – 25	100 – 125	500	07
Okra (Bhindi)	Fruit Borer	25	125	500	05
Maize	Spotted stem borer (*Chilo partellus*), Pink stem borer (*Sesamia inferens*), Fall army worm (*Spodoptera frugiperda*)	40	200	500	10
Groundnut	Tobacco caterpillar (*Spodoptera litura*)	30	150	500	28
Green gram	Pod borers (*Helicoverpa armigera, Spodoptera litura and Maruca testulatis*)	25	125	400-600	14

Red Gram	Gram Pod borer (Helicoverpa armigera), Pod fly (Melanogromyza obtusa	30	150	500	22
Chlorantraniliprole 00.40 % GR					
Rice (Paddy)	Yellow stem borer (*Scirpophaga incertulas*), Leaf folder (*Cnaphalocrosis medinalis*)	40	10 kg	Broadcast	53
Sugarcane	Early shoot borer,Top borer	75	18.75 kg	Broadcast	147
Maize	Fall armyworm (*Spodoptera frugipeda*), stem borer (*Chilo partellus*) and pink stem borer (*Sesamia inferns*)	50	12.5	Whorl Application	65
CHLORANTRANILIPROLE 0.53% w/w GR					
Rice	Yellow stem borer (*Scirpophaga incertulas*), Leaf folder (*Cnaphalocrosis medinalis*)	39.75	7500 g	Direct Broadcasting	63
Chlorantraniliprole 5% w/w DT					
Rice	Yellow Stem borer (*Scirpophaga incertulas*), rice leaf folder (*Cnaphalocrosis medinalis*)	40-50	1000-1250 (no of tablets)	NA	47
Chlorantraniliprole 35 % WG					
Okra	Fruit borer (*Helicoverpa armigera & Earias vittella*)	25	71	500	05
Tomato	Fruit borer (*Helicoverpa armigera*)	30	86	500	03
CHLORANTRANILIPROLE 47.85 % w/w SC					
Red gram	Pod borer (*Helicoverpa armigera*) Spotted pod borer (*Maruca spp*.) Pod fly (*Melanagromyza obtusa*)	30	50	500-750	31
Chickpea	Pod borer (*Helicoverpa armigera*) Tobacco caterpillar (*Spodoptera spp*.)	25	41.66	500	26
Sugarcane	Early shoot borer (*Chilo infuscatellus*), Top Borer (*Scirpophaga excerptalis*), Stalk borer	75	125	1000	214

	(*Chilo auricilius*)				
Soybean	Tobacco caterpillar (*Spodoptera litura*) Girdle beetle (*Obereopsis brevis*) Stem fly (*Melanagromyza spp*.) Green semilooper (*Chrysodeixis acuta*)	30	50	500	44
Maize	Fall armyworm (*Spodoptera frugiperda*) Stem borer (*Chilo partellus*) Pink stem borer (*Sesamia inference*)	40-50	67-83	500	87
Rice	Yellow stem borer (*Scirpophaga incertulas*), Rice leaf folder (*Cnaphalocrocis medinalis*)	30	50	500	47
Cotton	American Boll worm (*Helicoverpa armigera*) Tobacco caterpillar (*Spodoptera litura*)	30	50	500	67
Chili	Fruit borer (*Helicoverpa armigera*) Tobacco caterpillar (*Spodoptera litura*) Beet armyworm (*Spodoptera Exigua*)	30	50	500	3
Groundnut	Tobacco caterpillar (*Spodoptera litura*) Groundnut leaf miner (*Aproaerema modicella*)	30	50	400-500	34
Tomato	Tobacco caterpillar (*Spodoptera litura*) Fruit borer (*Helicoverpa armigera*)	30	50	400-500	3
Green gram	Pod borer (*Helicoverpa* armigera) Tobacco caterpillar (*Spodoptera litura*) Green semilooper (*Crysodeixis acuta*) Spotted pod borer (*Maruca testulalis*)	25	41.67	400-600	26
Chlorantraniliprole 50% W/w FS					
Rice	Yellow Stemborer (*Scirpophaga incertulas*), Leaf folder (*Cnaphalocrocis medinalis*)	75	120	Seed Dresser	-
Maize	Spotted Stemborer *(Chilo*	40	64	Seed	-

	partellus)			Dresser	
	Fall army worm (*Spodoptera frugiperda*)	70	112		
Chlorfenapyr 10 % SC					
Cabbage	Diamond back moth (*Plutella xylostella*)	75 – 100	750 –1000	500	07
Chilli	Mites (*Polyphagotarsonemus latus*)	75 – 100	750 –1000	500	05
Chlorfenapyr 240 g/L SC					
Red gram	*Helicoverpa armigera, Maruca vitrata*	192-240	800-1000	500	41
Cotton	*Spodoptera litura* and Thrips	75-102	312.5-425	500	48
Chickpea	Pod borer (*Helicoverpa armigera*)	192-240	800-1000	500	49
Brinjal	Fruit and Shoot Borer and Mites	192-240	800-1000	500	10
Chilli	Mites, Thrips and *Spodoptera litura*	75-102	312.5-425	500	7
	Helicoverpa armigera	192-240	800-1000	500	7
Soybean	*Spodoptera litura* and Green Semilooper	78-102	312.5-425	500	21
Chlorfluazuron 05.40 % EC					
Cabbage	Diamond back moth, Tobacco leaf eating caterpillar	75	1500	500	07
Cotton	American bollworm, Tobacco leaf eating caterpillar	75 – 100	1500 - 2000	500	10
Soybean	Fruit Borer (*Helicoverpa amigera*), Tobacco Caterpillar (*Spodoptera litura*)	75	1500	500	34
Chilli	Fruit Borer (*Helicoverpa amigera*), Tobacco Caterpillar (*Spodoptera litura*)	100	2000	500	3
Tomato	Fruit Borer (*Helicoverpa amigera*), Tobacco Caterpillar (*Spodoptera litura*)	100	2000	500	3
Chlorpyrifos 10 % Granules					

Rice (Paddy)	Stem borer, Leaf folder, Gall midge	1000	10000	-	30
Chlorpyriphos 75 % w/w WG					
Rice	Yellow stem borer *(Scirpophaga 16incertulas)*	375 – 400	500 – 533	500 –1000	15
Chlorpyrifos 19% ME					
Paddy	Stem Borer	270.75	1425	500	30
Chlorpyrifos 20 % EC					
Paddy (Rice)	Hispa	250	1250	500 –1000	-
	Leaf folder	375	1875	500 –1000	-
	Gall midge, Stem borer, Whorl maggot	250	1250	500 –1000	-
Beans	Pod borer, Black bug	600	3000	500 –1000	-
Gram	Cut worm	500	2500	500 –1000	-
Sugarcane	Black bug	150	750	500 –1000	-
	Early shoot & stalkborer	250 – 300	1250 –1500	500 –1000	-
	Pyrilla	300	1500	500 –1000	-
Cotton	Aphid, Bollworm,Whitefly	250	1250	500 –1000	-
	Cut worm	750	3750	500 –1000	-
Groundnut	Aphid	200	1000	500 –1000	-
	Root grub	225	1125	500 –1000	-
Mustard	Aphid	100	500	500 –1000	-
Brinjal	Shoot & fruit borer	200	1000	500 –1000	-
Cabbage	Diamond back moth	400	2000	500 –1000	-
Onion	Root grub	1000	5000	500 –1000	-
Apple	Aphid	0.05%	3750-5000	1500 –2000	-
Ber	Leaf hopper	0.03%	2250-3000	1500 –2000	-
Citrus	Black citrus, Aphid	0.02%	1500-2000	1500 –2000	-
Tobacco	Ground beetle	350	1750	500 – 1000	-

Termite control

Non cropped area: Building (Pre & Post construction treatment @1.0% a.i.)

Forestry @1.0% a.i.

Cropped area: Wheat: 3-4 ml/kg seed

Barley: 4-6 ml/kg seed

Gram: 15-30 ml/kg seed

Soil treatment: Wheat: 2-3 lit/ha.

Sugarcane: 6.25 lit/ha.					
Chlorpyrifos 50 % EC					
Rice (Paddy)	Stem borer, Leaffolder	375-400	750-800	500-1000	15
Cotton	Bollworms	500-600	1000-1200	500-1000	30
For non- agricultural use: - For protecting building from termite attack at pre and postsconstruction stages, apply Chlorpyriphos 50% EC @ 0.5% and 1.0% concentration.					
Chlorpyrifos 01.50 % DP					
Paddy (Rice)	Stem borer, Green leaf hopper, Brownplant hopper, Leaf folder, Gall midge, Grass hopper	375	25000	-	07
Bengal gram	Pod borer (*Helicoverpa armigera*)	375	25000	-	07
Chromafenozide 80 % WP					
Paddy (Rice)	Leaf folder, Stemborer	75-100	94-125	500	32
Clothianidin 0.5 % GR					
Okra	Jassids & White fly	40 – 60	8 - 12	--	01
Clothianidin 50 % WDG					
Rice (Paddy)	Brown plant hopper	10 – 12	20 – 24	500	12
Cotton	Jassids	15 – 20	30 – 40	500	20
	Whitefly	20 – 25	40 – 50	500	20
Cotton (Soil drench)	Jassids, Aphids,Thrips, Whitefly	100 – 125	200 – 250	1000	76
Sugarcane (Soil drench)	Termite	125	250	1000	310
Tea	Mosquito Bug (*Helopeltis theiovora*)	60	120	500	05
Grapes (Soil drench)	Thrips, Jassids and Mealy Bugs	250	500	1000	15
Groundnut (Soil Drench)	White Grubs	125	250	1000	62
Coumatetralyl 0.75 % w/w Gel					
Indoor or outdoor	Rats (*Rattus rattus,Rattus norvegicus, Bandicota bengalensis, Bandicota indica,*	01 mg per spot	2.50 per spot	-	-
	Tetra indica, Meriones hurrianae)				

Indoor	Mice	01	2.50	-	-
Coumatetralyl 0.0375 % Bait					
Indoor or outdoor	Rats (*Rattus rattus, Rattus norvegicus, Bandicota bengalensis, Bandicota indica, Tetra indica, Meriones hurrianae*)	01 mg per spot	02.50 per spot	-	-
Indoor	Mice	01	02.50	-	-
Cyantraniliprole 10.26 % OD					
Grapes	Thrips (*Scirtothrips dorsalis),* Flea beetle(*Scelodonta strigicollis)*	70	700	1000	05
Pomegranate	Thrips (*Scirtothripsdorsalis),* Pomegranate butterfly (*Deudorixossypi18s)*	75 (0.0075%)	750 (0.075%)	1000	05
	Whitefly (*Siphoninus phillyreae),* Aphids (*Aphis punicae*)	90 (0.009%)	900 (0.09%)	1000	05
Cabbage	Cabbage Aphid (*Brevicoryne brassicae*), MustardAphid (*Lipaphis erysimi*), Diamond back moth (*Plutellaxylostella*), Tobaccocaterpillar (*Spodoptera litura*)	60	600	500	05
Chilli	Thrips (*Scirtothrips dorsalis*), Fruit borer(*Helicoverpa armigera*), Tobaccocaterpillar (*Spodoptera litura*)	60	600	500	03
Tomato	Leaf miner (*Liriomyza trifolii*), Aphids (*Aphis ossypi*), Thrips (*Thrips tabaci*), Whitefly (*Bemesia tabaci*), Fruit borer (*Helicoverpa armigera*)	90	900	500	03
Gherkins	Leaf miner (*Liriomyzatrifolii*), Red pumpkinbeetle (*Aulacophora foveicolli*s), Aphids (*Aphis gossypi*), Thrips (*Thrips palmi*), Whitefly (*Bemesia tabaci*), Pumpkin caterpillar (*Diaphaniaindica*), Fruit fly (*Bactrocera cucurbitae*)	90	900	500	05

Okra	Whitefly- *Bemisia tabaci* Aphid- *Aphis gossypii* Shoot & fruit borer- *Earias vitella* Tobacco caterpillar- *Spodoptera litura* Fruit borer- *Helicoverpa armigera*	90	900	500	3
Brinjal	Whitefly- *Bemisia tabaci* Shoot and fruit borer – *Leucinodes orbonalis* Aphids- *Aphis gossypii* Thrips- *Thrips tabaci*	90	900	500	3
Cotton	Whitefly- *Bemisia tabaci* Aphids- *Aphis gossypii* Thrips- *Thrips tabaci* Tobacco caterpillar- *Spodoptera litura* Bollworms- *Helicoverpa armigera* & *Earias vitella*	90	900	500	7
Bittergourd	Thrips- *Thrips palmi* White fly- *Bemisia tabaci* Aphids- *Aphis gossypii* Pumpkin Caterpillar- *Diaphania indica* Leaf miner- *Liriomyza trifolii*	90	900	500	5
Ridgegourd	Thrips- *Thrips palmi* White fly- *Bemisia tabaci* Aphids- *Aphis gossypii* Pumpkin Caterpillar- *Diaphania indica* Leaf miner- *Liriomyza trifolii*	90	900	500	5
Watermelon	Thrips- *Thrips palmi* White fly- *Bemisia tabaci* Aphids- *Aphis gossypii* Leaf miner- *Liriomyza trifolii*	90	900	500	5
Potato	Thrips *(Thrips tabaci*) & Aphids (*Aphis gossypii)*	60	600	500	12
Capsicum	Thrips *(Scirtothrips dorsalis*), Tobacco caterpillar (*Spodoptera litura)* & Fruit borer (*Helicoverpa armigera)*	60	600	500	3
Citrus	Citrus Psylla (*Diaphorina citri)* & Thrips (*Scirtothrips dorsalis*)	60	600	500	5
CYANTRANILIPROLE 18.41% w/w SC					
Chilli	Thrips, , White fly, Tobacco caterpillar	**Dosage/1000 seedlings**		10 L	80
		8.75	43.75		
Capsicum	Thrips, Aphid, White fly, Leaf miner, Tobacco caterpillar	8.75	43.75	10	61

Tomato	Thrips, Aphid, White fly, Leaf miner	8.75	43.75	10	75
Cucumber	Thrips, Aphid, White fly, Leaf miner, Cucumber Moth	12.5	62.5	50	42
Watermelon	Thrips, Aphid, White fly, Leaf miner	15	75	50	47
Cyantraniliprole 48% w/w FS					
Maize	Fall Army Worm	1.8	3	-	NA
	Stem Borer	1.2	2		
	Cut Worm	1.2	2		
Cyclaniliprole 9.3% W/W DC (Cyclaniliprole 10.0% W/V DC)					
Chick pea	Pod borer (Helicoverpa armigera)	30	300	500	17
Soybean	Tobacco caterpillar (*Spodoptera litura*) and green semi looper (*Chrysodeixis acuta*)	40	400	500	41
Maize	Fall armyworm (*Spodoptera frugiperda*)	40	400	500	39
Sugarcane	Early shoot borer (*Chilo infuscatellus*) and top borer (*Scirpophaga excerptalis*)	100	1000	500	107
Cyclaniliprole 8.0 % SL					
Rice	Stem borer and Leaf Folder	35- 40	437.5 - 500	500	30
Cyclobutrifluram 38.4% w/w SC (Import only)					
Banana	Burrowing Nematode: *Radopholus similis*	225	500	1.2 L/ Plant	-
Okra	Root Knot Nematode (*Meloidogyne incognita*)	168.75	375	1.2 L/ Plant	-
Cucumber	Root Knot Nematode (*Meloidogyne incognita*)	168.75	375	50-100 ml/Plant	-
Tomato	Root Knot Nematode (*Meloidogyne incognita*)	168.75	375	50-100 ml/Plant	-
Cyenopyrafen 30 % SC					
Apple	Mite	60 – 90	200 – 300	1000	15
Chilli	Mite	60 – 90	200 – 300	400 – 600	07
Rose	Mite (*Tetranychus urticae*)	90	300	500	10
Brinjal	Mite (*Tetranychus urticae*)	90	300	500	10

Cyflumetofen 20 % SC					
Tea	Red spider mite	125 – 150	625 – 750	400 – 500	05
Cypermethrin 00.25% DP					
Brinjal	Fruit & shoot borer	50 – 60	20000 – 24000	-	03
Cypermethrin 10 % EC					
Cotton	Spotted bollworm, American bollworm, Pink bollworm	50 – 70	550 – 760	150 – 1000	07
Cabbage	Diamond back moth	60 – 70	650 – 760	100 – 400	07
Okra (Bhindi)	Fruit borer	50 – 70	550 – 760	150 – 400	03
Brinjal	Fruit & shoot borer	50 – 70	550 – 760	150 – 400	03
Wheat	Shoot fly	50	550.0	500 – 800	14
Sunflower	Bihar hairy caterpillar	60 – 70	650 – 760	500 – 700	14
Cypermethrin 25 % EC					
Cotton	Bollworms	40 – 70	160 – 280	400 – 800	-
	Jassids, Thrips	20 – 30	80 – 120	200 – 300	-
Bhindi (Okra)	Shoot & fruit borer,Jassids	37 – 50	150 – 200	500	03
Brinjal	Shoot & fruit borer,Jassids, *Epilachna* grub (Hadda beetle)	37 – 50	150 – 200	500	01
Dazomet					
Tobacco (Nursery)	Root knot nematode,Stunt nematode, Reni-form nematode	30 – 40	30 – 40	-	-
Tomato nursery	Root knot nematode	30 – 40	30 – 40	-	
Floriculture (Carnation &Gerbera)	Root-knot nematode	30 – 40	30 – 40		-
Deltamethrin 11% w/w EC					
Cotton	Bollworms	12.50	125	400 – 600	30

Rice (Paddy)	Stem borer, Leaf folder, Green leafhopper, Whorl maggot	15	150	500	13
Tea	Tea Thrips	10	100	400	15
Rice	Leaf folder	15 – 18.75	150 – 187.5	500	13
Tomato	Fruit borers	10 – 12.5	100 – 125	375 – 500	3
Okra	Fruit Borers	10-12.5	100-125	375 – 500	3
Chilli	Fruit borers	17.5	175	500	5
Onion	Thrips	15	150	500	5
Deltamethrin 25 % Tablet					
Cotton	Bollworms	12.50	50	400 – 600	30
Deltamethrin 01.80 % EC					
Cotton	Bollworms	12.50	781	400 – 600	30
	Sucking insects	10	625	400 – 600	30
Rice (Paddy)	Stem borer, Leaffolder	10 – 12.50	625 – 780	500	07
Deltamethrin 02.50 % WP					
Wheat & Rice (Grain & seed in stacks)	Rice weevil, Leasergrain borer, Khaprabeetle, Red flour beetle, Saw toothedgrain beetle, Rice moth, Almond moth	30	1200	1 litre/30 m^2	-
Walls, ceilings floors of Godowns	Rice weevil, Lesser grain borer, Khaprabeetle, Red flour beetle, Saw toothed grain beetle, Rice moth, Almond moth	30	1200	1.5-2.5 litre/50 m^2	-
Public health	Mosquito	625 – 1250	25000 – 50000	-	-
Deltamethrin 02.80 % EC					
Cotton	Bollworm	12.50	500	400 – 600	-
	Sucking Insects	10	400	400 – 600	-
Tea	Thrips, Caterpillar	3-4	120 – 150	400 – 600	03
	Leaf folder	10	400	400 – 600	03
	Lopper	2.50 – 3.75	100 – 150	400 – 600	03
Bhindi (Okra)	Shoot & fruit borer	10 – 15	400 – 600	400 – 600	01

	Jassid	10	400.0	400 – 600	01
Groundnut	Leaf miner	12.50	500.0	400 – 600	03
Mango	Hoppers	0.03 – 0.05 %	0.33-0.5 ml/lit	As per spray field requirement	01
Chilli	Fruit borer	10 – 12.5	400 – 500	400 – 600	05
Brinjal	Shoot & Fruit Borer	10 – 12.5	400 – 500	500	03
Red Gram (Arhar/Tur)	Pod Borer & Pod Fly	12.50	500.0	500	10

Deltamethrin 25% WDG

Name Of Insect Pest	**Dosage**			
	Infestation	**Requirement mg a.i./m²**	**Formulation (ml)/ litre water**	**Spray solution (ml)/ m²**
Mosquitoes (*Aedes aegypti, Anopheles stephensi, Culex quinquefasciatus*) **Housefly (**Musca domestica) **Cockroaches** (*Periplanata americana, blatella germinica*), **Bed Bugs** (*Cimex hemipterus*)	Low	10	0.8 g/ litre water	50
	Moderate	20	1.6/ litre water	50
	High	25	2.0/ litre water	50

Diafenthiuron 47.80 % SC

Cotton	Whiteflies, Aphids,Thrips, Jassids	250	500	500	41
Potato	Whiteflies, Leaf hoppers, Mites	250	500	500	44
Soyabean	Whiteflies, mites	250	500	500	44

Diafenthiuron 50 % WP

Cotton	Whiteflies, Aphids,Thrips, Jassids	300	600	500 – 1000	21
Cabbage	Diamond back moth	300	600	500 – 750	07
Chilli	Mites	300	600	500 – 750	03
Brinjal	Whitefly	300	600	500 – 750	03

Cardamom	Thrips, Capsule borer	400	800	1000	07
Citrus	Mites	1.0 g/l	2.0 g/l	2-3 Liter/ ha.	30
Cotton	Whiteflies, Aphids,Thrips, Jassids	239	500	500	30
watermelon	Whiteflies and Red spider mites	300	600	500	05
Okra	Whiteflies, Red Spider mites and Jassids	300	600	500	05
Tomato	Whiteflies and Redspider mites	300	600	500	05
Diflubenzuron 25 % WP					
Cotton	Tobacco Caterpillar	75 – 87.50	300 – 350	500 – 1000	-
	Bollworms	75	300	500 – 1000	-
Dimethoate 30 % EC					
Bajra	Milky weed bug	180-200	594-660	500 -1 000	-
Cotton	Aphis, Jassids, Thrips	200	660	500 - 1000	24
	Grey weevil	300	1000		
Cauliflower	Painted bug, MustardAphid	200	660	500 – 1000	-
Maize	Stem borer	200	660	500 – 1000	-
	Shoot fly	350	1155	500 – 1000	-
Sorghum	Midge	500	1650	500 – 1000	-
Castor	Jassids, Mites	250	825	500 – 1000	-
	Semi looper	350	1155	500 – 1000	-
Mustard	Leaf minor, Aphid,Sawfly	200	660	500 – 1000	-
Safflower	Aphid	200	660	500 – 1000	-
Potato	Thrips	200	660	500 – 1000	-
	Aphid	200	660	500 – 1000	-
Bhindi (okra)	Aphid	700	2310	500 – 1000	-
	Leaf hopper, Jassids	600	1980	500 – 1000	-
Brinjal	Jassids	600	1980	500 – 1000	-
	Shoot borer	200	660	500 – 1000	-
Rose	Scale	750	2475	500 – 1000	-
	Thrips	400	1320	500 – 1000	-
DIMPROPYRIDAZ 120 g/l SL					
Brinjal	White fly	108 -120	900 – 1000	500	3
	Jassids, Aphids	84	700	500	

Cotton	White fly	108 – 120	900 – 1000	500	31
	Jassids, Aphids	84	700	500	
Cucumber	White fly	108 – 120	900 – 1000	500	3
	Jassids, Aphids	84	700	500	
Tomato	White fly, Jassids, Aphids	108 - 120	900 – 1000	500	3
Chilli	White fly, Aphids	108 - 120	900 -1000	500	3
Dinotefuran 20 % SG					
Rice (Paddy)	Brown plant hopper	30 – 40	150 – 200	500	21
Cotton	Whitefly, Jassids,Aphids & Thrips	25 – 30	125 – 150	500	15
Dinotefuran 70 % WG					
Cotton	Aphids, Whitefly, Leaf Hopper	61.2	87	500	17
Brinjal	Jassids, Aphids and Whitefly	61.2	87	500	7
Rice	Brown Plant Hopper, White Backed Plant Hopper and Green leaf Hopper	61.2	88	500	32
Emamectin benzoate 05 % SG					
Cotton	Boll worms	9.5 –11.0	190 – 220	500	10
Okra (Bhindi)	Fruit & Shoot Borer	6.75 –8.50	135 – 170	500	05
Cabbage	Diamond back moth	7.5 – 10	150 – 200	500	03
Chilli	Fruit borer, Thrips,Mites	10	200	500	03
Brinjal	Fruit and Shoot borer	10	200	500	03
Red gram (Arhar/Tur)	Pod borer	11	220	500 – 750	14
Chickpea	Pod borer	11	220	500	14
Grapes	Thrips	11	220	500 – 1000	05
Tea	Tea looper	10	200	500	01
Emamectin benzoate 01.90 % EC					
Cotton	Boll worms	11	580	500	15
Chilli	Fruit borer , Thrips	07.13	375	500	03
Chick pea	Pod borer	07.13	375	500	14

Paddy	Leaf folder & Hispa	8.08	425	500	48
Soybean	Green semi looper,pod borer, Girdle beetle & Tobacco caterpillar	8.08	425	500	20
Ethion 50 % EC					
Tea	Red spider mites, Purple mites, Yellowmite, Thrips, Scale	250	500	500 – 1000	03
Cotton	Whitefly	750–1000	1500 –2000	500 – 1000	-
	Bollworms	1000	2000	500 – 1000	25
Chilli	Mites & thrips	75 – 1000	1500 –2000	500 – 1000	05
Gram	Pod borer	500 –750	1000 –1500	500 – 1000	21
Pigeon pea or Redgram (Arhar/Tur)	Pod borer	500 – 750	1000 –1500	500 – 1000	21
Soybean	Girdle beetle & stemfly	750	1500	500 – 1000	30
Ethofenoprox 10 % EC					
Rice	Brown plant hopper,Green leaf hopper, Stem borer, Leaf folder, Gall midge, Whorl maggot, White backed plant hopper	50 – 75	500 – 750	500	15

Ethylene dichloride + Carbon tetrachloride (3:1)

Crop	Common name of the pest	Cond.	Weight of volume	Exposure period	Conc. In air (ppm)	Aeratio n / Waitin g
Stored whole cereals Millets Pulses	Rice weevil, Lesser grain Borer, Khapra Beetle, Rust red flour beetle, Pulse beetle, Dried fruit Beetle	Air tight cover	300 –400 gm/m^3 (230 – 307 ml)	48 – 72 Hr. forcover fumigation	10 ppm	Partial aeration Forat least 1 hr. followed by 24 hr. complete Aeration waiting period of 24 hr.
Godown fumigation	Rice weevil, Lesser grain	Air tight cover	150 gm/m3	07 days	10 ppm	Partial aeration For at least 1

	Borer, Khapra Beetle, Rust red flour beetle, Pulse beetle, Dried fruitBeetle					hr. followed by 24 hr. complete Aeration waiting period of 24 hr.
Etoxazole 10 % SC						
Brinjal	Red spider mite		40	400	400 – 500	05
Tea	Red spider mite		40	400	400	05
Apple	European Red Spider Mite		40	400	1000	43
Fenazaquin 10 % EC						
Tea	Red spider mite, PinkMite, Purple mite		100	1000	400 – 600	07
	Scarlet mite		125	1250	400 – 600	07
Chilli	Yellow mite		125	1250	400 – 600	10
Apple	Red spider mite, Twospotted mite		40	400	1000	30
Okra (Bhindi)	Red spider mite		125	1250	500	07
Brinjal	Red spider mite		125	1250	500	07
Tomato	Two spotted spidermite		125.0	1250	500	07
Fenazaquin 18.3 % SC						
Brinjal	Red spider mite		114.375	625	400 – 500	10
Tea	Red spider Mite (*Oligonychus coffeae*)		80 – 100	400-500	400	07
Apple	European Red mite		40	200	1000	30
Chilli	Yellow Mite (*Polyphagotarsonemus latus*)		114.375	625	500	10
Okra	Red Spider Mite (Tetranychus urticae)		114.375	625	500	7
Tomato	Two Spotted Spider Mite (*Tetranychus spp*)		114.375	625	500	7
Fenmezoditiaz 200g/L SC						
Rice	Brown Plant Hopper (Nilaparvata lugens), White Backed Plant Hopper (Sogatella		25-30	125-150	500	37

	furcifera)				
Fenobucarb (BPMC) 50 % EC					
Rice	Brown plant hopper,Green leaf hopper	250 – 750	500 – 1500	500	30
Fenpropathrin 10 % EC					
Cotton	Pink boll worm, Spotted boll worm, American boll worm	75 – 100	750 – 1000	750 –1000	14
Fenpropathrin 30 % EC					
Cotton	Pink boll worm, Spotted boll worm, American boll worm, White fly	75 - 100	250 – 340	750 – 1000	14
Chilli	Thrips, Whitefly,Mites	75 – 100	250 – 340	750 – 1000	07
Brinjal	Whitefly, Shoot andFruit borer, Mites	75 – 100	250 – 340	750 – 1000	10
Okra (Bhindi)	Whitefly, Shoot andFruit borer, Mites	75 – 100	250 – 340	750 – 1000	07
Tea	Mites	50 – 60	165 – 200	400 – 500	07
Paddy (Rice)	Yellow stem borer,Leaf folder	100	333	500	30
Fenpyroximate 05 % EC					
Tea	Red spider mite, PinkMite, Purple mite	15 – 30	300 – 600	400 – 500	07
Chilli	Yellow mite	15-30	300 – 600	300 – 500	07
Cotton	Jassids, Mites	37.50	750	500	15
Coconut	Eriophyid mites	0.50 gm/tree (Root feeding)	10ml/lit.	As required	-
	Eriophyid mites	0.056-0.075 gm/tree	0.75-01.0 ml/lit.	As required	-
Fenpyroximate 05 % SC					
Chilli	Yellow mite	15 – 30	300 – 600	500 – 750	03
Tea	Red spider mite, Pinkmite, Purple mite	30 – 60	600 – 1200	400	07

Fenpropathrin 10% EW					
Rice	Stem borer (*Scirpophaga incertulas*) and leaf folder (*Cnaphalocrocis medinalis*)	100	1000	5000	58
Fenvalerate 20 % EC					
Cauliflower	Diamond back moth,American boll worm,Aphids, Jassids	60 -75	300 – 375	600 – 750	07
Cotton	Boll worm	75 -100	375 – 500	700 – 900	07
	Aphids, Jassids,Thrips	25 -40	125 – 200	250 – 400	07
Brinjal	Shoot & fruit borer,Aphids	75 – 100	375 – 500	600 – 800	05
Okra (Bhindi)	Shoot & fruit borer,Jassids	60 – 75	300 – 375	600 – 750	07
Fenvalerate 02 % Conc.					
Cotton	Spotted & Spiny, Pink American, Egyptian boll worm	80 – 100	4000 – 5000	-	-
Fenvalerate 00.40 % DP					
Cotton	Spotted Bollworm,Pink Bollworm	80 – 100	20000-25000	-	07
Fipronil 0.6% w/w WG					
Rice	Yellow stem borer (*Scirpophaga incertulas*)	50-75	8333-12500	Direct Broadcasting	49
Sugarcane	Early shoot borer (*Chilo infuscatellus*) and Termites (*Odontotermes obesus*)	80	13333	Direct Broadcasting	238
Fipronil 05 % SC					
Rice	Stem borer, Brown plant hopper, Green leaf hopper, Rice leaf hopper, Rice Gall midge, Whorl maggot, White backed plant hopper	50 – 75	1000 – 1500	500	32
Cabbage	Diamond back moth	40 – 50	800 – 1000	500	07
Chilli	Thrips, Aphids, Fruitborers	40 – 50	800 – 1000	500	07

Sugarcane	Early shoot borer &Root borer	75 – 100	1500 –2000	500	270
Cotton	Aphid, Jassid, Thrips, White fly	75 – 100	1500 –2000	500	06
	Boll worms	100	2000	500	07
Fipronil 18.87 % w/w SC					
Cotton	Thrips	75	375	375 – 500	21
Chilli	Thrips, Aphids, *Helicoverpa armigera*	50	250	500	5
Rice	Stem Borer, Leaf Folder , Brown Plant Hopper	50	250	500	46
Fipronil 02.92 % EC					
Pre-construction (Building)	Termite	0.25%	100	01	IS:6313 - 2001 (Part-2)
Post-construction (Building)	Termite	0.25%	100	01	IS:6313 -2001 (Part-3)
Fipronil 00.30 % GR					
Rice	Stem borer, Brown plant hopper, Green leaf hopper Rice leaf hopper, Rice gall midge, Whorl maggot, White backed plant hopper	50 – 75	16670 – 25000	-	32
Sugarcane	Early shoot borer, Root borer	75.0 – 100	25000 – 33300	-	09
Wheat	Termites	0.06	20 kg	-	91
Fipronil 00.60 % w/w GR					
Rice	Stem borer & Leaffolder	60	10	-	65
Sugarcane	Early shoot borer, Termite	75	12.5	-	229
Wheat	Pink Stem borer, Armyworm	60	10	-	106
	Termite	60	10	-	117
Fipronil 80 % WG					
Rice	Stem borer, Leaf folder	40 – 50	50 – 62.50	375 – 500	19
Grapes	Thrips	40 – 50	50 – 62.5	750 – 1000	10
Onion	Thrips	60	75	500	15
Cabbage	Diamond back moth	75	93.75	500	15
Chilli	Thrips	40 – 50	50 – 62.5	500	5

Cotton	Thrips (*Thrips tabaci*)	60	75	375-500	14
Flocoumafen 0.005% Block Bait (Strom)					
Usage	**Common pest**	**a.i. (mg)**	**Formulation (g)**	**How to apply**	**Waiting Period**
For rodent control in field , storage andcrops like rice, soybean and coconut)	*Rattus rattus, Bandicota bengalensis, Tatera indica, Mus musculus*	0.75-1.0	15-20	At an interval of 5-10m in bait station or active burrow. Repeat the application after 14days if problem persists.	NA
Flonicamid 50 % WG					
Rice	Brown plant hopper,White backed plant hopper, Green leaf hopper	75	150	500	36
Cotton	Aphids, Jassids, Thrips & Whiteflies	75	150	500	25
Okra	Aphids (Aphis gossipy), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	10
Brinjal	Aphids (Aphis gossipy), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	15
Mango	Mango Hoppers (*Amritodus atkinsoni*)	200	400	1000	15
Tomato	Aphids (*Aphis gossipy*), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	05
Chilli	Aphids (*Aphis gossipy*), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	05
Soybean	Aphids (*Aphis gossipy*), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	35
Potato	Aphids (*Aphis gossipy*), Jassids (*Amrasca bigutulla bigutulla*), White fly (*Bemisia tabaci*)	100	200	500	20
Flometoquin 10% SC					
Cabbage	Diamond Back Moth	75-100	750-1000	500	10
Chilli	Thrips (*Scirtothrips dorsalis*)	75-100	750-1000	500	10
Flubendiamide 20 % WG					

Rice	Stem borer, Leaffolder	25	125	500	30
Cotton	American bollworm	50	250	500	30
Pigeon pea (Tur/Arhar)	Pod borer	50	250	500	30
Cabbage	Diamond back moth	18.24	37.5 – 50	375 – 500	07
	Diamond back moth	12.5	62.5	500	07
Tomato	Fruit borer	48	100	375 – 500	05
	Fruit borer	50	250	500	05
Tea	Semilooper	30	150	400	07
Chilli	Fruit borer	50 – 60	250 – 300	500	05
Soybean	*Spodoptera litura*, Semilooper	50 – 60	250 – 300	500	29
Groundnut	*Spodoptera litura*	60	300	500	31
Black gram	*Spodoptera litura,Maruca* spp.	60	300	500	23
Bengal gram	Pod borer	50	250	500	15
Sugarcane	Early shoot borer	75	375	500 – 750	204
Maize	Fall Armyworm (*Spodoptera frugiperda*)	50	250	500	55
Flubendiamide 39.35 % w/w SC					
Rice	Stem borer, Leaffolder	24.0	50.0	375 – 500	40
Cotton	Bollworms (American & Spotted bollworm)	48 – 60	100 – 125	375 – 500	25
Pigeon pea	Pod borer	48	100	500	10
Black gram	Fruit borer	48	100	500	11
Chilli	Fruit borer	48 – 60	100 – 125	500	07
Tomato	Fruit borer	48	100	375 – 500	05
Cabbage	Diamond moth back	18.24	37.5 – 50	375 – 500	07
Brinjal	Shoot and fruit borer	72 – 90	150 – 187.5	500	05
Bengal gram	Pod Borer (*Helicoverpa armigera* & *Spodoptera* spp.)	48	100	500	05
Okra	Shoot & fruit borer	48 – 60	100 – 125	500	03

Soybean	Defoliators (*Helicoverpa armigera, Spodoptera litura*and Semilooper)	72	150	500	17
Gherkin	Fruit borer	36-48	75-100	500	5
Cardamom	Capsule borer	72	15ml/100 ltr water	500	15
Flubendiamide 00.70 % GR					
Paddy (Rice)	Stem borer	85 – 100	12.14-14.28	NA	25
Fluensulfone 2% GR					
Tomato	Root knot nematode (*Meloidogyne incognita*)	0.02g/plant & 444 to 512 g/ha	1.0 g/plant & 22.2 to 25.6 Kg/ha	-	7
Cucumber		0.02 g/plant & 160 to 200 g/ha	1.0 g/plant & 8.0 to 10 Kg/ha	-	55
Okra		0.02 g/plant & 800 to 910 g/ha	1.0 g/plant & 40.0 to 45.5 Kg/ha	-	7
Pomegranate		0.20g/ dripper & 0.80 g/ plant	10.0 g/ dripper & 40.0 g/ plant	-	91
Flufenoxuron 10 % DC					
Rose	Mites	50	500	500 –1000	06
Flumite 20 % SC / Flufenzine 20 % SC					
Brinjal	Mite	80 – 100	400-500	500 – 1000	05
Tea	Pink mite, Purplemite	80 – 100	400-500	500 – 1000	07
	Red spider mite	100 – 120	500-600	500 – 1000	07
Fluopyram 34.48 % w/w SC					
Tomato	Root knot nematode (*Meloidogyne incognita*)	250 (2 application) or 500 (Single application)	625 (2 application) or 1250 (Single application)	1000	05
Cucumber	Root knot nematode (*Meloidogyne incognita*)	250 (2 application) or 500 (Single application)	625 (2 application) or 1250 (Single application)	1000	05
Flupyradifurone 17.09 % w/w SL					
Okra (Bhindi)	Jassids, Whitefly	250	1250	500	03

Tea	Mosquito bug	150	750	400-500	7
Flupyrimin 2% GR					
Rice	Stem Borer, Brown Plant Hopper	100-150	5000-7500	NA	77
Flupyrimin 10 % SC					
Paddy	Brown Plant Hopper (*Nilparvata lugens*)	75-100	750-1000	500	7
Fluvalinate 25 % EC					
Cotton	Aphids, Jassids, Redcotton bug	50 – 100	200 -400	500 –1000	07
	Bollworm	50 – 100	200 -400	500 – 1000	07
Fluxametamide 10% w/w EC					
Brinjal	Leaf hopper, Thrips, Fruit and Shoot Borer	40	400	500	5
Cabbage	Diamond Back Moth, Tobacco caterpillar, Semi looper	40	400	500	5
Chilli	Thrips, Fruit Borer, Tobacco caterpillar	40	400	500	5
Okra	Leaf hopper, thrips, Fruit borer	40	400	500	5
Redgram	Spotted pod Borer, Pod Borer	40	400	500	5
Tomato	Thrips, Fruit borer	40	400	500	5
Gossyplure 4% RTU					
Cotton	Pink Bollworm	12.5	312.5	Ready to use	-
Hexythiazox 05.45 % w/w EC					
Tea	Scarlet mite (*Brevipalpus phoenicis*)Red spider mite (*Oligonychus cofeeaea*)	15-25	300 – 500	400/ha	05
Chilli	Yellow mites (*Polyphagotarsonemus latus*)	15 – 25	300 – 500	625/ha	03
Apple	European Red Mite (*Panonychus ulmi*)	0.002%	0.04%	10 ltr./tree	15
Grapes	Red spider mite	25	500	1000	05
Rose	Red spider mite	20-25	400 – 500	500	05
Brinjal	Red spider mite	25	500	500	07
Okra	Red spider mite	25	500	500	07
Imidacloprid 70 % WG					

Cotton	Jassids, Aphids,Thrips	21 – 24.5	30 – 35	375 – 500	07
Rice (Paddy)	Brown plant hoppers,White backed plant hoppers	21 – 24.5	30 – 35	300 – 375	07
Okra (Bhindi)	Jassids, Aphids,Thrips	21 – 24.5	30 – 35	300 – 375	03
Cucumber	Aphids & Jassids	24.5	35.0	500	05
Tomato	Thrips & White fly	35	50	500	05
Potato	Aphids & White fly	63	90	500	30
Imidacloprid 48 % FS (dosage per 100 kg of seeds)					
Cotton	Aphids, Whitefly, Jassids, Thrips	300 - 540	500 - 900	-	NR
Okra (Bhindi)	Jassid, Aphid	300 - 540	500 - 900	-	-
Sunflower	Jassid, Whitefly	300 - 540	500 - 900	-	-
Sorghum	Shoot fly	720	1200	-	-
Pearl millet	Shoot fly and termites	720	1200	-	-
Soybean	Jassids	75	125	-	-
Maize	Shoot fly	0.6	1.0	-	-
Potato	Aphid & Jassids	0.0105	0.0175	-	-
Rice	Thrips	0.15	0.25	-	-
Wheat	Aphids, Termite	0.21	0.35	-	-
Green Pea	Aphids, Jassids	0.12-0.18	0.2-0.3	-	-
Bengal gram	Termite	0.15	0.25	-	-
Black gram	Aphids, Jassids, Whitefly	0.42	0.7	-	-
Cumin	Aphids, Jassids, Thrips	0.36-0.42	0.6-0.7	-	-
Groundnut	Termite	0.12	0.2	-	-
Chilli	Aphids, Jassids, Thrips	0.72-0.90	1.2-1.5	-	-
Imidacloprid 70 % WS					
Cotton	Aphids, Whitefly,Jassids, Thrips	350 - 700	500 - 1000	-	NR
Okra (Bhindi)	Jassid, Aphid	350 - 700	500 - 1000	-	-
Chilli	Jassid, Aphid, Thrips	700 - 1050	1000 - 1500	-	-
Sunflower	Jassid, Whitefly	490	700	-	-
Sugarcane	Termite	70 - 105	100 - 150	-	-

Sorghum	Shoot fly	700	1000	-	-
Pearl millet	Termites and shootfly	700	1000	-	-
Mustard	Mustard sawfly,Painted bug	490	700	-	-
Imidacloprid 30.50 % m/m SC					
Cotton	Aphid, Jassids,Thrips	21 - 26.25	60 - 75	500 - 750	26
Rice (Paddy)	Brown plant hopper,White backed plant hopper	21 - 26.25	60 - 75	500 - 750	37
Chilli	Aphids, Thrips	43.75 – 52.5	125 - 150	500	5
For non- agricultural use:- For protecting building from termite attack at pre and post construction stages, apply Imidacloprid 30.5% m/m SC @ 0.075% a.i. concentration.					
Imidacloprid 17.80 % SL					
Cotton	Aphid, Whitefly,Jassid, Thrips	20 - 25	100 - 125	500 - 700	40
Paddy (Rice)	Brown plant hopper,White backed plant hopper, Green leaf hopper	20 - 25	100 - 125	500 - 700	40
Chilli	Jassid, Aphid, Thrips	25 - 50	125 - 250	500 - 700	40
Sugarcane	Termite	70	350	1875	45
Mango	Hopper	0.40-0.80g/tree	2-4ml/tree	10 litre	45
Sunflower	Whitefly, Jassid,Thrips,	20	100	500	30
Okra (Bhindi)	Aphid, Jassid, Thrips	20	100	500	03
Citrus	Leaf miner, Psylla	10	50	Depending on size of tree & Protection equipment used	15
Groundnut	Aphid , Jassid	20 - 25	100 - 125	500	40
Tomato	Whitefly	30 - 35	150 - 175	500	03
Grapes	Flea beetle	0.06 - 0.08	300 - 400	1000	32
Tea	Tea mosquito bug (*Helopeltis theiovora*)	24	120	400	07
Imidacloprid 00.30 % GR					
Paddy(Rice)	Stem borer	0.045	15.0 kg	-	26
Imidacloprid 17.1 % w/w SL					

Cotton	Aphid, Whitefly, Jassids, Thrips	50	250	500	50
Rice	Brown Plant Hopper, White Backed Plant Hopper and Green Leaf Hopper	60	300	500	39
Indoxacarb 14.50 % SC					
Cotton	Bollworm	75	500.0	600 - 1000	16
Cabbage	Diamond back moth	30 - 40	200 - 266	400 - 750	07
Chilli	Fruit borer	50-60	333 - 400	300 - 600	05
Tomato	Fruit borer	60-75	400 - 500	300 - 600	05
Pigeon pea	Pod borer complex	50-60	333 - 400	500 - 1000	15
Rice	Yellow Stem Borer (*Scirpophaga incertulas*), Leaf Folder (*Cnaphalocrosis medinalis*)	30-50	200-333	500	37
Soybean	Tobacco caterpillar (*Spodoptera litura*)	50	333	500	36
Chickpea	Pod Borer (*H. armigera*)	60-75	400-500	500	19
Indoxacarb 15.80 % EC					
Cotton	Bollworms (*Helicoverpa armigera*)	75	500	500 – 1000	14
Cabbage	Diamond back moth (*Plutella xylostella*)	40	266	500 – 1000	05
Pigeon pea	Pod borer complex (*Helicoverpa armigera*), Pod fly	50	333	500 – 700	12
Rice	Leaf folder (*Cnephalocrosis medinalis*), Green Semilooper, Stem fly	30	200	500	14
Soybean	Tobacco caterpillar (*Spodoptera litura*),Pod borer (*Heliothis armigera*), Green Semilooper (*Chrysodexis acuta*), stem fly (*Melanogromyza spp*.)	30	333	500	31
Chickpea	Pod borer *(Helicoverpa armigera)*	50	333	500	18
Isocycloseram 9.2% W/W Dc (10% W/V) DC					
Brinjal	Jassids and Red Spider mite	20	200	500	5
	Shoot and Fruit Borer	60	600		
Cabbage	Leaf Feeder, DBM	20-30	200-300	500	10
Chilli	Yellow Mites	20	200	500	7
	Thrips and Fruit borer	60	600		

Cotton	Jassids	20	200	500	37
	Thrips and Ball worm	60	600		
Red Gram	Gram Pod Borer and Spotted Pod borer	50-60	500-600	500	58
Groundnut	Leaf miner, Leaf feeder, thrips and Jassids	50-60	500-600	500	48
Soybean	Leaf worm, semi-looper, Gridle Beetle, Stem fly	60	600	500	35
Isocycloseram 18.1% W/W SC (20 % W/V Sc)					
Rice	Leaf Folder	20	100	500	29
	Stem Borer	60	300		
Maize	Fall Armyworm, Stem Borer	60	300	500	48
Lambda-cyhalothrin 04.90 % CS					
Cotton	Bollworms	25	500	500	21
Paddy (Rice)	Stem borer, Leaffolder	12.50	250	500	15
Brinjal	Shoot & fruit borer	15	300	500	05
Okra (Bhindi)	Fruit borer	15	300	500	05
Tomato	Fruit borer	15	300	500	05
Grapes	Thrips & Flea beetle	12.50	250	500 – 1000	07
Chilli	Thrips, Pod borer	25	500	500	05
Soybean	Stem fly, Semilooper	15	300	500	31
Pomegranate	Thrips & fruit borer	0.002	0.04	500-1000/ as per age of tree	5
Cardamom	Shoot and CapsuleBorer and Thrips	20	400	1000	34
Lambda-cyhalothrin 02.50 % EC					
Cotton	Bollworms, Jassids,Thrips	15 – 25	600 – 1000	400 – 600	21
Rice (Paddy)	Leaf folder, Stem borer, Green leaf hopper, Gall midge,Hispa, Thrips	12.50	500	400 – 600	15
Lambda-cyhalothrin 05 % EC					
Cotton	Bollworms, Jassids,Thrips	15 – 25	300 – 500	400 – 600	21
Rice (Paddy)	Leaf folder, Stem borer, Green leaf hopper, Gall Midge,Rice 38igut, Thrips	12.50	250	400 -600	15

Brinjal	Shoot & fruit borer	15	300	400 – 600	04
Tomato	Fruit borer	15	300	400 – 600	04
Chilli	Thrips , mite, podborer	15	300	400 – 600	05
Pigeon pea	Pod borer, Pod fly	20 – 25	400-500	400 – 600	15
Onion	Thrips	15	300	300 – 400	05
Bhindi (Okra)	Jassids , Shoot borer	15	300	300 – 400	04
Chickpea	Pod borer	25	500	300 – 400	06
Groundnut	Thrips, Leaf hopper,Leaf miner	10 – 15	200 – 300	400 – 500	10
Mango	Hoppers	0.0025-0.005%	0.5-1.0 ml/lof water	-	07
Lambda cyhalothrin 24% w/w WDG					
Rice	Yellow stem borer (*Scirpophaga incertulas*), Leaf folder (*Cnaphalocrocis medinalis*) Brown planthopper (*Nilaparvata lugens*)	12.5	52.0	500	29
Mango	Mango Hopper (*Idioscopus* clypealis), *I. niveosparsus* & I. *nitidulus*)	72.0	300.0	1500	7
LAMBDACYHALOTHRIN 22.8% CS					
Lambda-cyhalothrin 22.8% CS is a manufacturing concentrate that will be used for the manufacture of Lambda-cyhalothrin 4.9% CS and is not intended to be used directly on crops.					
Lufenuron 05.40 % EC					
Cabbage	Diamond back moth	30	600	500	14
Cauliflower	Diamond back moth	30	600	500	05
Pigeon pea	Pod borer, Pod fly	30	600	500 – 1000	65
Cotton	American bollworm	30	600	500 – 750	48
Black gram	Pod borer	30	600	500	10
Chilli	Fruit borer	30	600	500	05
Magnesium Phosphide Degesch plates recommended for fumigation of un-manufactured tobacco forexport, as per importing country requirement.					

Malathion 05 % DP					
Paddy (Rice)	Rice Hispa	1250	25000	-	-
Malathion 50.00% EC					
Paddy (Rice)	Rice Hispa	575	1150	500- 1000	-
Cabbage	Mustard aphid	750	1500	500 – 1000	-
Metaflumizone 22 % SC					
Cabbage	Diamond back moth	165 – 220	750 – 1000	500	03
Metaldehyde 2.5% DP					
Citrus, Rubber, Paddy (Rice), Tea,Vegetables	Snails, Slugs, Giant, African snails	Available in ready to use 2.5% Dust			
Methoxyfenozide 21.8 % w/w SC					
Groundnut	Leaf eating caterpillar (*Spodopotera litura*)	210	875	500	26
	Groundnut leaf minor (*Aproaerema modicella*)	210	875	500	
	Pod borer (*Helicoverpa armigera*)	210	875	500	
Sugarcane	Early shoot borer (*Chilo infuscatellus*)	120-150	500-625 (depending upon stage of crop)	500	161
Methyl Bromide 98 % w/w					
Stored Whole Cereals and Seed,Millet, Pulses	Rice Weevil, LesserGrain Bore, KhapraBeetle, Rust Red Flour Beetle, Saw Drug Store Beetle	Air tight cover	24 gm/m3	6-8 hours waiting Period 24 hrs.	As when residues not to exceed 25 ppm
Milled Products:Flour	Khapra Beetle, RustRed Flour Beetle, Lesser grain borer	Air tight cover	24 -32 gm/m3	12-24 hrs waiting Period 72 hrs	As when residues not to exceed 25 ppm
Dry Fruits, Nuts Spices & Oil Seeds	Rust Red Flour Beetle	Air tight cover	24 -32 gm/m3	24 hrs waiting Period 72hrs	As when residues not to exceed 25 ppm
Milbemectin 01 % EC					
Rose	Two spotted, Spidermite	04.50	450	1000	05

Chilli	Yellow , White mite	03.25	325	500	07
Monocrotophos 15 % SG					
Cotton	Aphids, Jassids, Thrips, Whiteflies	200	1333	500 – 1000	58
Nitenpyram 25% w/w SP					
Cotton	Jassids and Whiteflies	80	320	500	92
Paddy	Brown Plant hoppers & White backed plant hoppers	150	600	500	37
Novaluron 10 % EC					
Cotton	American Bollworm	100	1000	500 – 1000	40
Cabbage	Diamond back moth	75	750	500 – 1000	05
Tomato	Fruit borer	75	750	500 – 1000	1-3
Chilli	Fruit borer, Tobacco Caterpillar	33.50	375	500	03
Bengal gram	Pod borer	75	750	500	07
Novaluron 8.8 w/w % SC					
Cotton	American boll worm (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura*)	100	1000	500 -1000	20
Oxydemeton-methyl 25 % EC					
Paddy (Rice)	Blue leaf hopper	125	500	500 – 1000	-
	White leaf hopper	250	1000	500 – 1000	-
Maize	Shoot fly	250	1000	500 – 1000	-
Sorghum	Shoot fly	250	1000	500 – 1000	-
Cotton	Aphid, Jassid (leaf hopper)	300	1200	500 – 1000	-
Ground nut	Aphid, Leaf minor	250	1000	500 – 1000	-
Mustard	Aphid	250	1000	500 – 1000	-
Sesamum	Leaf hopper	300	1200	500 – 1000	-
Bhindi (Okra)	White fly	250	1000	500 – 1000	-
	Jassid, Leaf beetle	400	1600	500 – 1000	-

Chilli	Aphid	400	1600	500 – 1000	-
	Mites	500	2000	500 – 1000	-
	Thrips	250	1000	500 – 1000	-
Onion	Thrips	300	1200	500 – 1000	-
Tomato	White fly	250	1000	500 – 1000	-
Potato	Aphids	250	1000	500 – 1000	-
Apple	Sanjose scale	0.07%	4200- 5600	1500 – 2000	-
	Wooly Aphid	0.025%	1500-2000	1500- 2000	-
Banana	Tingid bug	0.025%	1500-2000	1500- 2000	-
	Aphids	0.05%	3000-4000	1500- 2000	-
Mango	Hoppers	0.025%	1500-2000	1500- 2000	-
Peaches	Leaf curl aphids	0.025%	1500-2000	1500- 2000	-
Coffee	Green bug	625	2500	500-1000	-
	Leaf minor	1000	4000	500-1000	-
Tobacco	White fly, Aphids	250	1000	500-1000	-
Permethrin 25 % EC					
Cotton	Bollworms	100 – 125	400 – 500	500 – 1000	-
Phenthoate 02 % DP					
Sorghum	Red spider mite, Pinkmite, Purple mite, Scarlet mite	400	20000	-	90 % Emergence of earhead
Safflower	Aphid	400	20000	-	-
Phenthoate 50 % EC					
Paddy (Rice)	Rice case worm	500	1000	500 -1000	-
Ground nut	Leaf Webber	500	1000	500 -1000	-
Phosalone 35 % EC					
Barely	Aphid	500	1428	500 – 1000	-
Sorghum	Ear head midge	400	1143	500 – 1000	-
Jute	Red spider mite	350	1000	500 – 1000	-

Brinjal	Fruit borer	500	1428	500 – 1000	-
Cabbage	Aphid	500	1428	500 – 1000	-
Tomato	Fruit borer	450	1285	500 – 1000	-
Tea	Aphid, Pink mite,Purple mite	360	1028	500 – 1000	-
Phosalone 04 % DP					
Sorghum	Earhead midge	1000	25000	-	-
Phosmet 50% WP					
Chilli	Aphids, Thrips & Fruit borer	500	1000	500	10
Rice	Yellow stem borer & Leaf folder	600	1200	500	45
Cotton	Jassid, Aphids, Whitefly & Bollworms	600	1200	500	47
Apple	Wolly apple aphid (*Eriosoma lanigerum*), Green Aphid (*Aphis pomi*), Tortricid moth	0.5ml/l	1	1	31
Profenofos 50 % EC					
Cotton	Bollworm	750 – 1000	1500 – 2000	500 – 1000	15
	Jassids, Aphids, Thrips, Whiteflies	500	1000	500 – 1000	15
Soybean	Semi looper & Girdle beetle	500	1000	500	40
Chilli	Fruit borer, Tobacco caterpillar, Thrips, Aphids	500	1000	500	3
Onion	Thrips	500	1000	500	80
Propargite 57 % EC					
Tea	Red spider mite, Pinkmite, Purple mite, Scarlet mite	430-612	750-1250	400	07
Chilli	Mite	850	1500	500 – 625	07
Apple	European red mite,Two spotted mite	2.85-5.7/tree	5-10 ml/tree	10 lit/tree	09
Brinjal	Two spotted spidermite	570	1000	400	06
Pymetrozine 50 % WG					
Paddy (Rice)	Brown plant hopper	150	300	500	19

Mango	Hoppers	150	300	1000	36
Pyrifluinazon 20% WG					
Cotton	White fly (Bemisia tabaci)	100	375-500	500	30
Pyriproxyfen 10 % EC					
Brinjal	White fly & Jassids	50	500	300	07
Cotton	Whitefly	100	1000	500	31
Cotton	Whitefly	50-60	500-700	500	50
Chilli	Whitefly, Aphids	50	500	300	07
Okra	White fly & Jassids	50	500	300	07
Pyriproxyfen 10% EW					
Cotton	Whitefly (*Bemisia tabaci*), Jassid (*Amrasca biguttula*) and thrips (*Thrips tabaci*)	100-125	1000-2500	500	38
Pyridaben 20 % w/w WP					
Tea	Red spider mite	100	500	500	07
Cotton	White fly	100	500	500	28
Chilli	Yellow mite	75 – 100	375 – 500	500	5
Brinjal	Red Spider Mite	100	500	500	3
Apple	European Red Mite	1	5	10 litre water/ tree	5
Pyridalyl 10 % EC					
Cotton	Bollworms	75 – 100	750 – 1000	500 – 750	07
Okra	Fruit & shoot borer	50 – 75	500 – 750	500 – 750	03
Cabbage	Diamond back moth	50 – 75	500 – 750	500 – 750	03
Maize	Fall armyworm	100	1000	750	27
Quinalphos 25 % Gel					
Chilli	Aphid	250	1000	500 – 1000	-
Paddy (Rice)	Brown plant hopper,Leaf folder, Stem borer, Hispa	250	1000	500 – 1000	-
Quinalphos 05 % Granules					
Paddy (Rice)	Gall midge, Stemborer	250	5000	-	-

Quinalphos 20 % AF					
Rice (Paddy)	Brown plant hopper,Green leaf hopper, Leaf folder, Stem borer	250 – 300	1250 – 1500	750 – 1000	40
Okra (Bhindi)	Shoot /Fruit borer	250 – 300	1250 – 1500	750 – 1000	07
Cotton	American bollworm,Pink Bollworm, Spotted bollworm	350 – 500	1750 – 2500	750 – 1000	07
Tomato	Fruit borer	300 – 350	1500 – 1750	750 – 1000	07
Tea	Hopper caterpillar	0.05%	1000	400	07
Pigeon pea	Pod borer	500.	2500	750 – 1000	30
Groundnut	Spodoptera	250 – 375	1250 – 1775	750 – 1000	30
Quinalphos 25 % EC					
Paddy (Rice)	Brown plant hopper	375	1500	500 – 1000	40
	Hispa/blue beetle	500	2000	500 – 1000	40
	Leaf folder	250	1000	500 – 1000	40
	Stem borer	325	1300	500 – 1000	40
Wheat	Aphid	250	1000	500 – 1000	-
	Ear head Caterpillar,Mite	400	1600	500 – 1000	-
Bengal gram	Pod borer	250	1000	500 – 1000	-
Black gram	Bihar hairy caterpillar	375	1500	500 – 1000	-
French bean	Stem fly	250	1000	500 – 1000	-
Red gram	Pod borer, Pod fly	350	1400	500 – 1000	30
Soybean	Leaf weevil	250	1000	500 – 1000	-
Groundnut	Leaf Hopper, Thrips	350	1400	500 – 1000	30
	Leaf miner	250	1000	500 – 1000	30
Mustard	Sawfly	300	1200	500 – 1000	-
Sesamum	Leaf Webber, Jassids	500	2000	500 – 1000	-
Bhindi (Okra)	Fruit borer	200	800	500-1000	-

	Leaf hopper, Mite	250	1000	500-1000	-
Cauliflower	Stem borer	500	2000	500 – 1000	-
Chilli	Aphid	250	1000	500 – 1000	-
	Mite	375	1500	500 – 1000	-
Tomato	Fruit borer	250	1000	500 – 1000	-
Apple	Wooly Aphid	0.05%	3000 – 4000	500 – 1000	-
Banana	Tingid bug	0.05%	3000 – 4000	500 – 1000	-
Citrus	Scale	0.07%	4200 – 5600	500 – 1000	-
	Citrus butterfly	0.025%	1500 – 2000	500 – 1000	-
Pomegranate	Scales	0.08%	4800 – 6400	500 – 1000	-
Tea	Thrips	190	760	500 – 1000	07
Quinalphos 01.50 % DP					
Paddy (Rice)	Brown plant hopper	300	20000	-	40
Gram	Pod borer	350	23300	At pod formation	
Red gram	Pod borer	350	23300	-	30
Soybean	Leaf weevil	250	16600	-	-
French bean	Stem fly	30	20000	-	-
Cotton	Aphid, Jassids, Thrips	300	20000	From square formation onwards	
	Bollworms	450	30000	From square formation onwards	
Ground nut	Thrips, Jassids	350	23300	-	30
	Red hairy caterpillar	375	25000	-	30
Safflower	Aphid	300	20000	-	-
Chilli	Aphid	300	20000	-	-
Spinetoram 11.70 % SC					
Cotton	Thrips	50	420	500 – 1000	30
	Tobacco caterpillar	50-56	420 – 470	500 – 1000	
	Spotted boll worm				
	Pink boll worm	54	450	500-1000	

Soybean	Tobacco caterpillar	54	450	500 – 625	30
Chilli	Thrips, Fruit borer, Tobacco caterpillar	56-60	470-500	400 – 500	07
Okra	Fruit borer (*Helicoverpa armigera*), Shoot and Fruit borer (*Earias 47iguttul*)	45-54	375-450	500-1000	3
Brinjal	Shoot and Fruit borer	45-54	375-450	500-1000	3
Chickpea	Pod Borer (*Helicoverpa armigera)*	45-54	375-450	500-1000	20
Red gram	Pod Borer (*Helicoverpa armigera, Maruca vitrata)*	45-54	375-450	500-1000	23
Grapes	Thrips (*Scirtothrips dorsalis*)	36	300	500-1000	5
Rice	Yellow Stem Borer (*Scirpophaga incertulas*),	42-45	350-375	500	20
	Leaf Folder (*Cnaphalocrosis medinalis*)	30	250	500	
Tomato	Fruit borer (*Helicoverpa armigera),* Tobacco caterpillar (*Spodoptera litura*), Leaf miner (*Liriomyza trifolii*), tomato pinworm (*Tuta absoluta*)	45-54	375-450	500	3
Mango	Thrips	0.05ml/L	0.4 ml/L	-	21
Maize	Fall Armyworm (*Spodoptera frugiperda*)	30	250	500	32
Spinetoram 0.1% w/w RB					
Maize	Fall Army Worm (*Spodoptera frugiperda*)	5 -7.5	5000 - 7500	NA	21
Spinosad 45 % SC					
Cotton	American bollworm	75-100	165-220	500	10
Chilli	Fruit borer, Thrips	73	160	500	03
Chilli	Fruit borer (*H.armigera*) (*Scirtothrips dorsalis*)	56-73	124-162	500	03
Red gram	Pod borer	56 – 73	125 – 162	800 – 1000	47

Brinjal	Fruit & Shoot borer	73 – 84	162 – 187	500	03
Grapes	Thrips	25 ml/100 lit	250	1000	15
Spinosad 02.50 % SC					
Cabbage & Cauliflower	Diamond back moth	15 – 17.50	600 – 700	500	03
Spiromesifen 22.90 % SC					
Brinjal	Red spider mite	96	400	500	05
Cotton	White fly & mite	144	600	500	10
Apple	European Red Mite& Red spider mite	72 (0.03%)	300	1000	30
Chilli	Chilli Yellow Mite	96	400	500 – 750	07
Tea	Red spider mite	96	400	400	07
Okra (Bhindi)	Red spider mite	96 – 120	400 – 500	500	03
Tomato	Whiteflies & Mites	150	625	500	03
Spirotetramat 15.31 % w/w OD					
Chilli	Thrips & Aphids	60	400	500	05
Okra	Aphid, Whitefly, Mites	90	600	500	03
Grapes	Mealy bug, Mites	105	700	500 – 1000	60
Cotton	Aphids, Whiteflies and Thrips	105	700	500	52
Citrus	Psylla and Mites	0.90/10 Lit water	6 ml/10 litres water	500	30
Cabbage	Diamond back moth (*Plutella xylostella*)	90	600	500	7

Sulfoxaflor 21.8 % w/w SC

Crop	**Target pest**	**Dosage/ha**		**Water (l/ha)**	**Waiting period**
		a.i. (gm/ha)	**Formulation (ml/ha)**		
Rice	Brown Plant Hopper (*Nilaparvatha Lugens*)	90	375	500	14
	White backed planthopper (*Sogatella furcifera*)	90	375	500	

Cotton	Jassids (*Amarasca bigutella*),	75	313	500	
	Aphid (*Aphis gossypi*)	75	313	500	
	Whitefly (*Bemmissia tabaci*)	90	375	500	
	Cotton mealy bug (*Phenococcus spp.*)	90	375	500	
Tebufenpyrad 20% WP					
Tea	Red Spider Mites	75-100	375-500	500	7
Tomato	Red Spider Mites	75-100	375-500	500	20
Tetraniliprole 18.18% w/w SC					
Rice	Yellow stem borer (*Scripophaga 49iguttula49*) Leaf folder (*Cnaphalocrocis medinalis*)	50 – 60	250 – 300	500	43
Soybean	Girdle beetle (*Oberea brevis*) *Spodoptera spp.* Semilooper (*Chrysodeixis acuta*)	50 – 60	250 – 300	500	35
Tomato	Fruit borer (*Helicoverpa armigera*); Leaf miner (*Liriomyza trifolii*)	60	300	500	5
Tetraniliprole 40.34% FS					
Rice	Stem borer and leaf folder	4.8-6.0	10.0-12.5	NA	NA
Maize	Stem borer	2.4-3.6	5.0-7.5	NA	NA
Thiacloprid 21.70 % SC					
Cotton	Aphid, Thrips, Jassid	24 – 30	100 – 125	500	52
	Whitefly	120-144	500 – 600	500	52
Paddy (Rice)	Stem borer	120	500	500	30
Chilli	Thrips	54 – 72	225 – 300	500	05

Tea	Mosquito bug	90	375	400	07
Brinjal	Shoot & fruit borer	180	750	500	05
Soybean	Girdle beetle	180	750	500	17
Apple	Thrips	0.01-0.012%	0.04-0.05%	As persize of tree	30
Thiocyclam Hydrogen Oxalate 50% SP					
Rice	Stem borer, Leaffolder	500	1000	500	30
Thiodicarb 75 % WP					
Cabbage	Diamond back moth	750 – 1000	1000 – 1330	500	07
Cotton	Bollworms	750	1000	500	30
Brinjal	Shoot & Fruit borer	470 – 750	625 – 1000	500	06
Chilli	Fruit borer	470 – 750	626 – 1000	500	06
Black gram	Pod borer (*Maruca* spp.) & (*Helicoverpa*spp.)	468 – 562	625 – 750	375 – 500	17
Pigeon Pea	Pod Borer	470 – 750	625 – 1000	500	30
Thiamethoxam 1.2% W/W G.R					
Rice	Yellow stem borer , Leaf folder, Brown plant hopper	120	10000	Broadcasting	52
Groundnut	White grub ,Termites	120	10000	Broadcasting	53
Sugarcane	Early shoot borer ,White grub, Termites	120	1000	Broadcasting	237
Thiamethoxam 30 % FS					
Cotton	Aphid, whiteflies,Jassids	03	10	This is used as seed dresser	
Sorghum	Shoot fly	03	10	This is used as seed dresser	
Wheat	Termites	01	3.3	This is used as seed dresser	
Soybean	Shoot fly	03	10	This is used as seed dresser	
Chilli	Thrips	02.1	7.0	This is used as seed dresser	

Okra (Bhindi)	Jassids	01.7	5.7	This is used as seed dresser
Maize	Stem Fly	02.4	8	This is used as seed dresser
Sunflower	Jassids, Thrips	03	10	This is used as seed dresser
Maize	Aphids	1.8 g a i/kg seed	6 ml/kg seed	10-12 ml/kg seed
	Shoofly	2.4 g a i/kg seed	8 ml/kg seed	10-12 ml/kg seed
Cotton	Aphids Jassids White fly and Thrips	2.4 - 3 g a i/kg seed	8-10 ml/kg seed	10-15 ml/kg seed
Soybean	Jassids	1.2 g a i/kg seed	4 ml/kg seed	6-8 ml/kg seed
	Stem fly Girdle beetle Whitefly	1.8 g a i/kg seed	6 ml/kg seed	6-8 ml/kg seed
Groundnut	Jassids	0.9 g a i/kg seed	3 ml/kg seed	6-8 ml/kg seed
	Aphids and Thrips	1.5 g a i/kg seed	5 ml/kg seed	6-8 ml/kg seed

Thiamethoxam 30 % WS (Seed treatment)

Name of crop	Name of Pest	a.i	Formulation	Dilution
Maize	Aphids	1.8 g ai/kg seed	6 ml/kg seed	10-12 ml/kg seed
	Shoot fly	2.4 g ai/kg seed	8 ml/kg seed	10-12 ml/kg seed
Cotton	Aphids, Jassids, Whitefly & Thrips	2.4 - 3g ai/kg seed	8-10 ml/kg seed	10-15 ml/kg seed
Soybean	Jassids	1.2 g ai/kg seed	4 ml/kg seed	6-8 ml/kg seed
	Stem fly Girdle beetle, Whitefly	1.8 g ai/kg seed	6 ml/kg seed	6-8 ml/kg seed
Groundnut	Jassids	0.9 g ai/kg seed	3 ml/kg seed	6-8 ml/kg seed
	Aphids and Thrips	1.5 g ai/kg seed	5 ml/kg seed	6-8 ml/kg seed

Thiamethoxam 70 % WS

Cotton	Aphid, Thrips, Whitefly, Jassids	300	430	Use as seed dresser at the time of sowing
Okra (Bhindi)	Aphids, Jassids	200	286	Use as seed dresser at the time of sowing

Tomato	Aphids, Thrips	420	600	Use as seed dresser at the time of sowing	
Sunflower	Jassids, Thrips	280	400	Use as seed dresser at the time of sowing	
Wheat	Termite, Aphids	121	175	Use as seed dresser at the time of sowing	
Maize	Shoot fly, Aphids	245	350	Use as seed dresser at the time of sowing	
Rice (Paddy)	Thrips, Green leafhopper	105	150	Use as seed dresser at the time of sowing	
Thiamethoxam 75 % w/w SG					
Groundnut	Termite	94	125	500 – 1000	57
Sugarcane	Termite, Early shootborer	120	160	500 – 1000	230
Rice (Paddy)	Green leaf hopper, Brown plant hopper	113	150	Dissolve in 500 ml water and mix with 20 kg sand/ha.	60
Cotton	Jassids & Thrips	94	125	50 – 100 ml/plant	10 9
Thiamethoxam 25 % WG					
Rice (Paddy)	Stem borer, Gall midge, Leaf folder, White backed plant hopper, Brown planthopper, Green leaf hopper, Thrips	25	100	500 – 750	14
Cotton	Jassid, Aphid,Thrips	25	100	500 – 750	21
	Whitefly	50	200	500 – 750	21
Okra (Bhindi)	Jassid, Aphid,Whitefly	25	100	500 – 1000	05
Mango	Hoppers	25	100	1000	30
Wheat	Aphid	12.5	50	500	21
Mustard	Aphid	12.5 – 25	50-100	500 – 1000	21
Tomato	Whitefly	50	200	500	05
Foliar application	(Apply first spray during initial appearance of pest and repeat 2 – 3 sprays at 10 – 15 days interval depending on the level of pest intensity)				
Brinjal	Whitefly	50	200	500	03
Foliar application	(Apply first spray during initial appearance of pest and repeat 2 – 3 sprays at 15 – 21 days				

	interval depending on the level of pest intensity)				
Tea	Mosquito bug	25	100	400 – 500	07
Potato	Aphids: Foliar Application	25	100	500	77
	Soil drench	50	200	400 – 500	77
Citrus	Psylla	25	100	1000	20
	(Apply first spray during initial appearance of pest and repeat 2 – 3 sprays at 15 – 21 days interval depending on the level of pest intensity)				
Rice-Nursery (SoilDrenching)	Green leaf hopper,Thrips, Whorl Maggot	500	2000	250 ml/sq.mtr	86
Tomato Soil drench	White flies	100	400	500	05
	(Apply root zone after transplanting as soil drench once during crop season.)				
Cumin	Aphids	25	100	500	05
Grapes –Soil drench	Mealy bug, Thrips, Flea beetle	100	400	200ml/vine	10
	(Apply first soil drench immediately after pruning followed by another drenching at 30-40 days interval coinciding with berry development stage. Apply the product near root zone of vines.)				
Grapes – Foliar application	Mealy bug, Thrips, Flea beetle	25	100	1000	15
	(1st spray when insect pest reaches ETL. Repeat one spray at 10-15 days interval depending upon pest severity)				
Tolfenpyrad 15 % EC					
Cabbage	Diamond back moth,Aphids	150	1000	500	05
Okra (Bhindi)	Aphids, Jassids,Thrips, Whitefly	150	1000	500	03
Cotton	Aphids, Jassid s,Thrips, Whitefly	150	1000	500	26
Cumin	Aphids, Thrips	150	1000	500	29
Chilli	Aphids, Thrips	150	1000	500	7
Mango	Hoppers, Thrips	150.0	1000	500	7
Onion	Thrips	150.0	1000	500	10
Pomegranate	Thrips	150.0	1000	500	7
Triflumezopyrim 10% w/w SC					
Paddy	Brown plant hopper & White backed plant hopper	25	236	500	21

Mango	Hoppers (*Idioscopus spp.*)	25	23.6 ml/100 L water	1000	30
Triflumezopyrim 20% w/w WG					
Rice	Brown Plant Hopper (*Nilaparvata lugens*), White Backed Plant Hopper (*Sogatella furcifera*)	25	125	500	21
Zinc Phosphide 80 % Powder					
Crop	**Pest organism**	**Dosage**		**Technical**	
For rodent control in field and residential premises(to be used under the supervision of trained personal)	*Rattus rattus, Bandicota bengalensis, Rattus meltade , Tatera indica, Meriones hurrianae, Mus platythrix, Mus musculus, Rattus norvegicus, Musbooduga, Suncus caeruleus*	1.5-2.5% active ingredient inbait		Mix 10 g of Zinc phosphide with 10g of edible oil and then mixwith 380g of food material. Keep 10g of poisoned bait at each point.	
Combination Product					
Acephate 50 % + Bifenthrin 10 % WDG					
Cotton	Leaf hopper, Thrips, Bollworms	400 + 80.	800	500 – 750	20
Acephate 45 % + Cypermethrin 5 % DF					
Cotton	Aphid, Jassids, Thrips& White fly.	425	850	500 – 600	22
Maize	Stem borer (Chilo partellus) Fall armyworm (Spodoptera frugiperda)	625	1250	500	52
Acephate 25 % + Fenvalerate 03 % w/w EC					
Cotton	American bollworm, Sucking pest	500 + 60.	2000	500	15
Acephate 50 % + Fipronil 5% WDG					
Paddy	Stem borer, Leaf folder,	500 + 50	1000	500	27

	Brown Plant Hopper				
Acephate 50 % + Imidacloprid 01.80 % SP					
Cotton	Aphid, Jassids, Thrips, Whitefly, Bollworms	518	1000	500	40
Paddy (Rice)	Brown Plant Hopper,Green Leaf Hopper, Stem borer & Leaf folder	518	1000	500	--
Sugarcane	Termite, White grubs, Early Shoot borer, Black bug, Pyrilla	1250 + 45	2500	500	123
Chilli	Jassids (*Amrasca biguttula biguttula*), Aphid (*Aphis gossypii*), Whitefly (*Bemisia tabaci*) Thrips (*Scirtothrips dorsalis*), Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura*)	518	1000	500	03
Acephate 50 % + Imidacloprid 5 % WG					
Cotton	Jassids, Aphids, Thrips, White fly, Bollworms (*Pectinophora gossypiella*, *Helicoverpa armigera*, *Earias vitelli*)	500 + 50	1000	500	53
Acetamiprid 20% + Chlorantraniliprole 20% WG					

Cabbage	Diamond back moth (*Plutella xylostella*), Tobacco caterpillar (*Spodoptera litura*) and Aphid (*Lipaphis erysimi*)	15+15	75	500	3
Tomato	Tomato fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura),* Aphid (*Aphis gossypii)* Leafhopper (*Amrasca biguttulla biguttulla*) and Whitefly (*Bemisia tabaci*).	30+30	150	500	7
Acetamiprid 19.50 % + Chlorantraniliprole 19.50% w/w FS					
Maize	Shoot fly (Atherigona soccatta), Fall armyworm (Spodoptera frugiperda) Stem borer (Chilo partellus) and Aphid (Rhopalosiphum maidis)	22+22 g per 10 kg seed	100 ml per10 kg seed	NA	-
Acetamiprid 00.40 % + Chlorpyriphos 20 % EC					
Paddy (Rice)	Stem Borer, Brownplant hopper & White backed plant hopper	10 + 500	2.5	500 - 800	10
Acetamiprid 00.40 % + Chlorpyriphos 20 % EC					
Paddy (Rice)	Stem borer, Brown plant hopper,	10 + 500	2.50	500 – 800	10

	Whitebacked plant hopper				
Acetamiprid 01.10 % + Cypermethrin 05.50 % EC					
Cotton	Aphids, Jassids, Thrips, Bollworms	10 + 50	1000	400 – 1000	30
Azoxystrobin 10.0% + Fipronil 5% SC					
Chilli	Thrips and fruit borer	50+100	1000	500	5
Rice	Yellow stem borer, leaf folder, brown plant hopper	62.5+ 125	1250	500	53
Azoxystrobin 1.3% + Tebuconazole 0.22% + Thiamethoxam 25.9% FS					
Okra (seed treatment)	Aphids and Jassids	0.9+0.15+18.0	60	NA	NA
Benzpyrimoxan 10% + Pymetrozine 20% WG					
Rice	Brown Plant Hopper, White Backed Plant Hopper and Green Leaf Hopper	50+100 – 70+140	500-700	500	36
Benzpyrimoxan 10% + Thiamethoxam 3.3% SC					
Rice	Brown Plant Hopper, White Backed Plant Hopper	75 + 25	750	500	30
Benzpyrimoxan 15% + Fipronil 10% SC					
Rice	Brown Plant Hopper, White Backed Plant Hopper and leaf folder	60-75 + 40-50	400-500	500	22
Beta-cyfluthrin 08.49 % + Imidacloprid 19.81 % w/w OD					
Brinjal	Aphids, Jassids, Shoot & fruit borer	15.75 + 36.75- 18 + 42	175 – 200	500	07
Soybean	Girdle beetle Semilooper	31.5 + 73.5	350	500	17

Cotton	Jassid Whitefly	18 + 42	200	500	21
Chilli	Thrips, Aphids, Whitefly	27.9 + 65.1	310	500	5
Cucumber	Whitefly	45 + 105	500	500	5
Wheat	Aphids	36 + 84	400	500	31
Cashew	Tea mosquito bug and Thrips	22.5 + 52.5	250	500 - 1000	45
Tea	Mosquito bug and Looper	27 + 63	300	400	7
,					
Paddy (Rice)	Stem borer, Leaf folder	24 + 240- 30 + 300	800 – 1000	500	21
Bifenthrin 8% + Clothianidin 10% SC					
Ground nut	Whitegrub, thrips and aphids	80+100	1000	1000	83
Cotton	Grey weevil, mealy bug, jassids, whitefly, aphids and thrips	80+100	1000	1000	73
Sugarcane	Termites and early shoot borer	80+100	1000	1000	300
Bifenthrin 32 % + Chlorantraniliprole 12% WG					
Soybean	White fly (*Bemisia tabaci*), Girdle beetle (*Obereopsis brevis*) Green semilooper (*Chrysodeixis acuta*) and Stem fly (*Melanagromyza sojae*)	80 + 30	250	500	40
Bifenthrin 10% + Thiamethoxam 5% SE					

Paddy	Stem Borer , Leaf Folder, Gall Midge & Brown Plant Hopper	50 + 25	500	500	39
Bifenthrin 25% + Thiamethoxam 25% WG					
Cotton	Jassids, Aphids, Thrips White fly, American Bollworm, Pink Bollworm, Spotted Bollworm.	50 + 50	200	500	42
Paddy	Stem borer, Leaf folder, Hispa, Brown plant hopper, Green leaf hopper.	50 + 50	200	500	25
Buprofezin 09 % + Acephate 24 % w/w WP					
Rice (Paddy)	Brown plant hopper	54 + 144	600	500	20
Buprofezin 15 % + Acephate 35 % w/w WP					
Cotton	Jassids, Thrips & White fly	187.5 + 437.5	1250	500	48
Okra	Jassids & White fly	112.5 + 262.5	750	500	07
Paddy (Rice)	Brown plant hopper,White backed plant hopper	187.5 + 437.5	1250	500	20
Buprofezin 20 % + Acephate 50 % w/w WP					
Paddy (Rice)	Stem Borer, Leaf folder, Brown plant hopper	200 + 500	1000	500	20
Cotton	Thrips, Jassids, Mealy bug	250 + 625	1250	500	15
Chilli	Thrips, Yellow mites Aphids	250 + 625	1250	500	7
Buprofezin 20 % + Acetamiprid 2% w/w WP					

Rice	Brown plant hopper, White backed plant hopper, Leaf Folder, Green Leaf Hopper, Stem Borer	176	800	400	15
Buprofezin 22. 0% + Fipronil 3 % SC					
Rice (Paddy)	Brown plant hopper	110 + 15	500	400 – 500	32
Cotton	Mealy bugs & Thrips	220+30	1000	500	17
Buprofezin 23.10 % + Fipronil 3.85 % w/w SC					
Rice	Brown plant hopper (*Nilparvata lugens)*	173.25 + 28.88	750	500	30
CARTAP HYDROCHLORIDE 7.5% + CHLORANTRANILIPROLE 0.4 % GR					
Paddy	Yellow stem borer Leaf folder and Whorl maggot	691.25	8750	NA	59
Cartap Hydrofipronil ide 50 % + Buprofezin 10 % w/w WP					
Rice	Yellow stem borer, Brown plant hopper,Leaf folder, Green leaf hopper, White backed plant hopper	480	800	500	20
Cartap Hydrochloride 7.5 % w/w + Emamectin benzoate 0.25 %w/w GR					
Rice	Yellow stem borer (*Scirpophaga incertulas*)	18.75 + 562.5	7.5	-	35
Cartap Hydrochloride 04 % + Fipronil 00.50 % CG					
Paddy (Rice)	Stem borer, Leaffolder	675 – 900	15 – 20	-	27

Soybean	Leaf worm, Girdle beetle, Semilooper, Stem fly	28	200	500	41
Chlorantraniliprole 4.3% +Abamectin 1.7% SC					
Chilli	Thrips, Mites and Fruit Borer	26.875 + 10.625	625	500	5
Tomato	Fruit Borer, Leaf Miner, mites	21.5 + 8.5	500	500	5
Chlorantraniliprole 0.60%+Thiamethoxam 1.25%+Fipronil 1.25% GR					
Rice	Rice stem borer (*Scirpophaga incertulas*), Leaf folder (*Cnaphalocrocis medinalis*)	30 + 62.5+62.5	5000	NA	91
Chlorantraniliprole 24% + Emamectin Benzoate 8.8% WG					
Chilli	Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura*), Thrips (*Scirtothrips dorsalis*)	30+11	125.0	500	5
Maize	Stem borer (*Chilo partellus*), Fall armyworm (*Spodoptera frugiperda*)	36+13.20	150	500	27
Chlorantraniliprole 15.75% w/w + Clothianidin 26.25% w/w SC					
Sugarcane	Early shoot borer, Top borer, Termites and White grub	200	400	1000	258
Chlorantraniliprole 0.53% w/w + Fipronil 0.8% w/w GR					

Rice	Rice yellow stem borer (*Scirpophaga incertulas*)	39.75 + 60	7.5 kg	Broadcasting mixed with sand	53
Chlorantraniliprole 0.4% + Fipronil 0.6% W/W GR					
Paddy	Stem borer (*Scirpophaga incertulas*) and Leaf folder (*Cnaphalocrocis medinalis*)	40+60 (gm)	10 ml	NA	34
Chlorantraniliprole 0.35% + Fipronil 0.35% GR					
Paddy	Stem Borer (*Scirpophaga incertulas*) and Leaf Folder (*Cnaphalocrocis medinalis*)	35 + 35	10 Kg	Braoadcasting mixed with sand	65
Chlorantraniliprole 12% + Fipronil 30% GR					
Chilli	Thrips (*Scirtothrips dorsalis*), Aphids (*Aphis gossypii*), Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (*Spodoptera litura*)	30 + 75	250	500	10
Cotton	Thrips (*Thrips tabaci*), Aphids (*Aphis gosypii*), American bollworm (*Helicoverpa armigera*), Pink bollworm (*Pectinophora gossypiella*)	30 + 75	250	500	41
Chlorantraniliprole 0.83% + Fipronil 1.33% GR					
Paddy	Stem Borer (*Scirpophaga incertulas*), Leaf Folder (*Cnaphalocrocis*	108.3	5000 gm	NA	49

	medinalis) and Hispa (*Dicladispa armigera*)				
Chlorantraniliprole 0.45% + Fipronil 0.50% + Emamectin Benzoate 0.18% GR					
Tomato	Whitefly & Fruit borer	45 + 50 + 18	10 Kg/Ha	NA	3
Paddy	Yellow stem borer	45 + 50 + 18	10 kg/ha	N.A.	31 days
Chlorantraniliprole 09.30 % + Lambda-cyhalothrin 04.60 % ZC					
Pigeon pea	Pod borer	30	200	500	18
Cotton	Bollworms complex	37.50	250	500	20
Brinjal	Shoot and fruit borer,Jassids	28	200	500	05
Okra	Shoot and fruit borer,Jassids	28	200	500	03
Rice	Stem borer, Leaf folder & Green leaf hopper	28 – 35	200 – 250	500	53
Soybean	Leaf Worm, Girdle Beetle, Semi looper, Stem Fly	28	200	500	41
Okra	Shoot & Fruit Borer, Jassids	28	200	500	3
Cabbage	Spodoptera, Diamond Back Moth	30	200	500	5
Groundnut	Leaf miner, Leaf feeder, Thrips	28	200	500	20
Blackgram	Pod Borers (*Spodoptera litura, Maruca vitrata, Helicoverpa armigera*)	28 (18.6+9.20)	200	500	24
Pomegranate	Fruit Borer & Thrips	28	200	1000	7
Maize	Fall Army worm and Stem borer	35	250	500	36
Grapes	Flea beetle and Thrips	28	200	1000	10

Tomato	*Helicoverpa armigera*, *Spodoptera litura*, and Leaf Miner	37.5	250	500	5
Chlorantraniliprole 10% w/w + Novaluron 20% w/w SC					
Cabbage	Diamond back Moth	30 + 60	300	500	10
Chilli	Fruit borer and Tobacco Caterpillar	30 + 60	300	500	10
Maize	Fall armyworm	30 + 60	300	500	21
CHLORANTRANILIPROLE 5.2% + NOVALURON 8.5% SC					
Chickpea	Pod borer (*Helicoverpa armigera*),	20.8 + 34	400	500	7
Chilli	Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (Spodoptera litura)	26+42.5	500	500	7
Tomato	Fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (Spodoptera litura)	26+42.5	500	500	7
Chlorantraniliprole 4.5 % + Novaluron 11.5 + Emamectin Benzoate 1.5 % SE					
Brinjal	Shoot & Fruit borer	28.12 + 71.87 + 9.37	625	500	16
Chlorantraniliprole 0.64% + Tebuconazole 3% + Clothianidin 1.92% GR					
Paddy	Yellow Stem Borer and Sheath Blight	40 + 187.5 + 120	6250	NA	46
Chlorantraniliprole 00.50 % + Thiamethoxam 01 % w/w GR					
Rice	Stem borer, Leaf folder, Brown plant hopper, Green leaf hopper	30.0 + 60.0	6 kg/ha	-	60
Chlorantraniliprole 08.80 % + Thiamethoxam 17.50 % w/w SC					

Tomato	Leaf Miner, Whitefly, Fruit borer	150 (50+100)	500 **Application method**-Soil drench (Single application), **Application time-**8-10 days after transplanting	50-100 ml/plant	36
Rice Nursery	Stem borer, Leaf folder, Green leaf Hopper	180 (60+ 120)	600 **Application method**-Soil drench (Single application), **Application time-**At the time of sowing to before transplanting	100L/ha	116
Chlorantraniliprole 7.50 % + Thiamethoxam 7.50 % w/w OD					
Transplanted Rice	Stem borer (*Scirpophaga incertulas*) Whorl maggot (*Hydrellia sasakii*)	36 + 36	480	Splash into water at 15 - 20 days after transplanting	95
Chlorpyrifos 50 % + Cypermethrin 05 % EC					
Cotton	Aphid, Jassids, Thrips,Whitefly, *Spodoptera litura*, Spotted bollworm, Pink bollworm, American bollworm	500 + 50	1000	500 - 1000	15
Rice (Paddy)	Stem borer, Leaf folder	312 + 32	625 – 750	500 – 700	15
Brinjal	Shoot & Fruit Borer	500+50	1000	500	7
Cabbage	Diamond Back Moth	375+37.5	750	500	5
Okra	Fruit borer (*Earias vitella*)	500+50	1000	500	10
Chlorpyriphos 16 % + Alphacypermethrin 01 % EC					

Cotton	Spotted bollworm, Pink bollworm, American bollworm	425	2500	500 – 750	15
Clothianidin 3.5%+Pyriproxyfen 8% SE					
Brinjal	Whitefly, Jassid, Thrips and Aphids	44+100	1250	500	3
Cotton	Whitefly, Jassid, Thrips and Aphids	44+100 – 52.5+120	1250-1500	500	60
Cyantraniliprole 7.3% w/w + Diafenthiuron 36.4% w/w SC (Cyantraniliprole 8% w/v + Diafenthiuron 40% w/v SC) (Cyantraniliprole 80 g/L + Diafenthiuron 400 g/L – 480 SC)					
Cotton	Jassids, Whitefly, Thrips, Aphids, Pink bollworm	300 (50 + 250)	625	500	29
Chilli	Thrips, Mites, Whitefly, Fruit borer	300 (50 + 250)	625	500	5
Cyantraniliprole 10%w/w + Pymetrozine 50%w/w WG					
Rice	Stem borer, Leaf Folder, Brown Plant Hopper	180	300	500	43
Cyantraniliprole 16.9% + Lufenuron 16.9% w/w SC					
Rice	Stem borer, Leaf folder	20 (10+10)	50	500	39
Tomato	Fruit borer: Helicoverpa armigera Leaf miner: Liriomyza trifolii Thrips: Thrips tabaci	60 (30+30)	150	500	5
Maize	Fall Army Worm: Spodoptera frugiperda Stem borer: Chilo partellus / Sesamia inferans	60 (30+30)	150	500	44
Cotton	Pink boll Worm: Pectinophora gossypiella Leaf worm: Spodoptera litura American Bollworm: Helicoverpa armigera Thrips: Thrips tabaci	100 (50+50)	250	500	25

CYANTRANILIPROLE 19.8%w/w + THIAMETHOXAM 19.8%w/w FS					
Maize	Shoot fly, Stem borer, Cut worms, Aphids	1.58 (Cyantranilip role 0.79 + Thiamethoxa m 0.79)	4	10-20	Single application by seed treatment before sowing.
	Fall Army Worm (Spodoptera frugiperda)	2.38 (Cyantranilip role 1.19 + Thiamethoxa m 1.19)	6		
Cypermethrin 10 % + Indoxacarb 10 % w/w SC					
Cotton	Jassids, Thrips, Bollworms	50 +50	500	400 – 1000	07
Rice	Yellow stem Borer	25 + 25	250	500	37
	Leaf Folder	37.5+37.5	375		
Cypermethrin 3 % + Quinalphos 20 % EC					
Brinjal	Shoot & Fruit borer	-	350 – 400	500 – 600	07
Cotton	American bollworm, Spotted bollworm, Jassids	-	1000 – 1250	500 – 600	15
Cyantraniliprole 7.20% + Tolfenpyrad 12.60% + Abamectin 1.35% w/w SC					
Chilli	Thrips (*Scirtothrips dorsalis*), Black thrips (*Thrips parvispinus*), Fruit borer (Helicoverpa armigera) and yellow mites (*Polyphagotarsonemus latus*)	50 + 87.5 + 9.38	625	500	10
Deltamethrin 00.72 % + Buprofezin 05.65 % w/w EC					
Rice (Paddy)	Brown plant hopper, Leaf folder	0.78 + 62.50- 0.94 + 75.00	1250 + 1500	500	30
Diafenthiuron 40.1 % + Acetamiprid 3.9 % w/w WP					
Cotton	Thrips, Mites and Whitefly	(200.5 + 19.5 –	500 - 600	500	21

		(240.6 + 23.4)			
Chilli	Thrips, Mites and Whitefly	(200.5 + 19.5 – (240.6 + 23.4)	500 - 600	500	21
Diafenthiuron 47 % + Bifenthrin 9.4 % w/w SC					
Cotton	Thrips (*Thrips tabaci*), Leaf hopper (*Amrasca devastans*),Whitefly (*Bemisia tabaci*), Aphid (*Aphisgossypii*)	293.75 + 58.7	625	500	30
Chilli	Thrips (*Scirtothrips dorsalis*), Aphids (*Aphis gossypii*)	293.75 + 58.7	625	500	07
Diafenthiuron 48 % + Dinotefuran 8% WG					
Cotton	Thrips (*Thrips palmi*), Jassids (*Amrasca bigutulla bigutulla*), Whiteflies (*Bemisia tabaci*), Mites (*Tetranychus urticae)*	300 + 50	625	500	43
Brinjal	Thrips (Thrips palmi), Jassids (*Amrasca bigutulla bigutulla*), Whiteflies (*Bemisia tabaci*), Mites (*Tetranychus urticae)*	300 + 50	625	500	03
Diafenthiuron 18% + Fipronil 8% + Dinotefuran 5% w/w SC					
Chilli	Thrips (*Scirtothrips dorsalis*) Fruit borers (*Helicoverpa armigera* and *Spodoptera litura*)	157.50 + 70.00 + 43.75	875	500	28
Diafenthiuron 30 % + Pyriproxyfen 8% w/w SC					
Chilli	Mites & Whitefly	300+80	1000	500	3
Cotton	Whitefly	300+80	1000	500	35
Dimethoate 20% (w/w) + Cypermethrin 3% (w/w) EC					
Brinjal	shoot and fruit borer (*Leucinodes orbonalis*),	122.9 + 18.4	800	500	7

	Jassids (*Amrasca biguttula biguttula*) and Epilachana Beetle (*Henosepilachna vigintioctopunctata*)	(141.3)			
Dinotefuran 4 % + Acephate 50% w/w/ SG					
Rice	Brown Plant Hopper & white Backed Plant Hopper	35 + 400	500	500	28
Cotton	Aphids, Jassids, Thrips & White flies.	22 + 275	880	500	10
Dinotefuran 5% + Ethion 50% EC					
Rice	Brown plant hopper, White backed Plant hopper, Green leaf hopper and Leaf folder	50 + 500	1000	500	33
Dinotefuran 11% + Pymetrozine 36% WG					
Paddy	Brown plant hopper *(Nilparvata lugens*), White backed plant hoppers (*Sogatella furcifera*)	164.5	350	500	36
Dinotefuran 15 % + Pymetrozine 45% WG					
Rice	Brown Plant Hopper, whiteBacked Plant Hopper, Green Leaf Hopper, Rice Ear Head Bug	200	333	500	24
Dinotefuran 12%+Thiamethoxam 22.5%+Pymetrozine 30% w/w WG					
Paddy (Rice)	Brown planthopper, Nilaparvata lugens (Stal) White backed planthopper, Sogatella furcifera (Horvath)	42.00 + 78.75 +105.00	350	500	18
Emamectin benzoate 1.9% + Chlorantraniliprole 5 % + Deltamethrin 1 % w/w EC					
Chilli	Thrips (*Scirtothrips dorsalis*) Mites (*Polyphagotarsonemus latus*) Fruit borers (*Helicoverpa armigera*	7.13 + 18.75 + 3.75	375	500	28

	and *Spodoptera litura*)				
Emamectin Benzoate 1.1% + Diafenthiuron 30 % SC					
Cotton	Aphids , Jassids, thrips, whitefly & Pink Bollworm	11 + 300	1000	500	38
Brinjal	Aphids, Thrips, Whitefly, red Spider Mites, Shoot borer and fruit borer	11 + 300	1000	500	08
Emamectin Benzoate 01.50 % + Fipronil 03.50 % SC					
Chilli	Thrips, Fruit borer	07.5 + 17.5-11.25 + 26.25	500 – 750	500	03 (day) or 48 (Hrs) Re-entry period after each application
Emamectin Benzoate 3.5% (w/w) + Lambda Cyhalothrin 5% (w/w) WP					
Chilli	Fruit borer (*Helicoverpa armigera*), Thrips (*Thrips tabaci*), Mites (*Tetranychus urticae*)	8.75 + 12.5	250	500-700	7
Emamectin Benzoate 1.25% + Lambda Cyhalothrin 1.5% + Bifenthrin 4.5% EC					
Paddy (Rice)	Stem borer and leaf folder	(9.375 + 11.25 + 33.75) = **54.375**	750	500	17
Emamectin benzoate 5 % w/w + Lufenuron 40 % w/w WG					
Cauliflower	Diamond Back Moth (*Plutella xylostella*) Fruit borer (*Spodoptera litura & Helicoverpa armigera*)	27 (Emamectin benzoate 3.0 + Lufenuron 24.0)	60	500	03
Chilli	Fruit borer (*Spodoptera litura & Helicoverpa armigera*), Thrips (*Scritothrips dorsalis*), Mites (*Polypagotarsonemus latus*)	27 (Emamectin benzoate 3.0 + Lufenuron 24.0)	60	500	03
Maize	Fall Army Worm and Stem borer	36 (Emamectin benzoate 4.0 + Lufenuron 32.0)	80	500	36

Tea	Looper, Thrips & Leaf roller	27 (Emamectin benzoate 3.0 + Lufenuron 24.0)	60	450	7
Tomato	Fruit Borer and Thrips	27 (Emamectin benzoate 3.0 + Lufenuron 24.0)	60	500	5
Brinjal	Shoot & Fruit borer	27 (Emamectin benzoate 3.0 + Lufenuron 24.0)	60	500	5
Emamectin Benzoate 2.2% + Permethrin 15.3 % EC					
Maize	*Spodoptera frugiperda* (FAW)	16.5 + 114.75	750	500	-
Emamectin Benzoate 1.5% + Profenofos 35% w/w WDG					
Cotton	Whiteflies, Jassids, Thrips, Aphids and Pink Boll Worm	10.5 + 245	700	500	15
Chilli	Yellow Mites, Thrips and Fruit Borer	10.5+245	700	500	7
Maize	Fall armyworm	11.25+262.5	750	500	35
Emamectin Benzoate 3.8% (w/w) + Thiamethoxam 20% (w/w) WDG					
Okra	Aphid (*Aphis gossypii*), Jassid (*Amrasca devastans*), Whitefly (*Bemisia tabaci),* Shoot and fruit borer (*Earias vittela).*	3.8 + 20	100	500-700	5
Emamectin Benzoate 3% + Thiamethoxam 12% WG					
Chilli	Fruit borer (*Helicoverpa armigera*) and Thrips (*Scirtothrips dorsalis*)	6 + 24	200	500	3
Green gram	Pod borer (*Helicoverpa armigera*) and Leaf hopper (*Empoasca kerri*)	6 + 24	200	500	15
Tea	Tea Mosquito bug, Tea Looper	(4.5 + 18.0) – (6 + 24)	150 - 200	400 - 500	7
Ethion 40 % + Cypermethrin 05 % w/w EC					
Cotton	American bollworm	400 + 50	1000	500	15
Chilli	Mites and Thrips	600 + 75	1500	500	10
Ethiprole 40% + Imidacloprid 40 % WG					

Rice (Paddy)	Brown plant hopper	37.50 + 37.50	93.75	375	15
	White backed plant hopper	50 + 50	125	375	15
Ethiprole 10.7% + Pymetrozine 40% WG					
Rice	Brown plant hopper & White backed plant hopper	40.12+150 to 45.48+170	375-425	375	27
Etoxazole 6% + Abamectin 1.5% SC					
Tea	Red Spider Mite, Purple Mite, Scarlet Mite	33.75 (27 + 6.75)	450	500	3
Brinjal	Red Spider Mite	33.75 (27 + 6.75)	450	500	5
Fenazaquin 10% + Bifenthrin 4% EC					
Tomato	Red Spider mite (*Tetranychus urticae*) Whitefly (*Bemisia tabaci*)	125+50	1250	500	5
Tea	Red Spider mite (*Oligonychus coffeae*) Mosquito Bug (*Helopeltis theivora*)	100+40	1000	400	7
Cotton	Whitefly and Mites	125 + 50	1250	500	74
Fenobucarb 20 % + Buprofezin 05 % w/w SE					
Paddy (Rice)	Brown plant hopper, Green leaf hopper	400 + 100	2000	500	30
Fenobucarb 22.5% + Buprofezin 11.25% + Acephate 2.5% ME					
Paddy	Brown plant hopper, Green leaf hopper & White Backed Plant Hopper	393.75 + 196.875 + 43.75	1750	500	25
Fipronil 5% + Buprofezin 20% SC					
Chilli	Thrips	37.5+150	750	500	5
	Fruit Borer	50+200	1000		
Cotton	Jassids, Thrips. Aphid	50+200	1000	500	6

	& Whitefly				
Rice	Brown Plant Hopper	25 + 100	500	500	20
	Yellow Stem Borer, Leaf Folder	50 + 200	1000	500	
Fipronil 15% + Chlorantraniliprole 5% w/v SC					
Rice	Stem borer, Leaf folder	75 + 25	500	500	49
Fipronil 15% + Deltamethrin 2.5% w/v SC					
Tomato	Tomato fruit borer	75 + 12.5	500	500	7
Fipronil 10% + Diafenthiuron 30% w/w WG					
Cotton	Jassids (*Amrasca bigutulla bigutulla*), thrips (*Thrips tabaci*), Pink bollworm (*Pectinophora gossipiella*)	75 + 225	750	500	21
FIPRONIL 12.5% + LAMBDA-CYHALOTHRIN 5% ZC					
Cotton	Thrips, Leaf hoppers, Aphids & Pink bollworm	50 + 20	400	500	35
Chilli	Thrips, Aphids & Fruit borer (*Helicoverpa armigera*)	50 +20	400	500	7
Onion	Thrips	50 + 20	400	500	7
Tomato	Thrips, Whitefly & Fruit borer (*Helicoverpa armigera*)	50 +20	400	500	7
Fipronil 2.50% + Propargite 35.00% SE					
Chilli	Aphids, thrips, mites & Fruit Borer	50 + 700	2000	500	20
Flubendiamide 04 % + Buprofezin 20 % w/w SC					
Paddy (Rice)	Yellow stem borer, Leaf folder, Brown plant hopper	35 + 175	175 + 700	500	30
Flubendiamide 8.33 % + Deltamethrin 5.56 % w/w SC					
Chickpea	Pod borer	22.50 + 15	250	500	07

Cucumber	Cucumber beetle, Fruit fly	18 + 12- 22.50 + 15	200 – 250	500	05
Rice	Leaf folder (*Cnaphalocrocis medinalis)* and Hispa (*Dicladispa armigera)*	27+18	300	500	35
Chilli`	Thrips (*Scirtothrips dorsalis*) and Fruit borer (*Helicoverpa armigera* and *Spodoptera litura*)	36+24	400	500	5
Tomato	Thrips and Fruit borer	36+24	400	500	5
Flubendiamide 03.50 % + Hexaconazole 05 % w/w WG					
Paddy (Rice)	Stem borer, Leaf folder	35 + 50	1000	500	20
Groundnut	*Spodoptera litura*	52.5 + 75	1500	500	31
Chilli	*Spodoptera litura, Helicoverpa armigera*	52.5 + 75	1500	500	10
Tomato	Spodoptera litura Helicoverpa armigera Early blight Leaf spot	Flubendiamide – 43.75 & Hexaconazole – 62.5	1250	500	5
Flubendiamide 07.5 % + Kresoxim Methyl 37.5 % w/w SC					
Rice	Stem borer & Leaf folder	50 + 250	667	500	30
Tomato	Fruit borer (*Helicoverpa armigera*) Leaf eating caterpillar/ Fruit borer (*Spodoptera litura*)	50 + 250	667	500	07
Flubendiamide 19.92 % + Thiacloprid 19.92 % w/w SC					
Chilli	Thrips, Fruit borer	48 + 48 – 60 + 60	200 – 250	500	05
Rice	Yellow stem borer, Leaf folder	60 + 60	250	500	33

Tea	Mosquito Bug and Semi looper	60 + 60	250	400	7
Brinjal	Fruit & shoot borer, Aphids, Thrips, Jassids	84 + 84	350	500	5
Fluxametamide 5% +Flonicamid 15% + Abamectin 1.5% ME					
Chilli	aphids, whitefly, thrips, mites, tobacco caterpillar and fruit borer	31.25 + 93.75 + 9. 375	600	500	5
Fipronil 04 % + Acetamiprid 04 % w/w SC					
Cotton	Aphid, Jassids, Whitefly	40 + 40	1000	500	30
Fipronil 15% + Imidacloprid 5% w/v SC					
Cotton	Aphids, Jassids, Thrips	75 + 25	500	500	6
	Pink bollworm	90 + 30	600		
Fipronil 40 % + Imidacloprid 40 % WG					
Sugarcane	White grub (*Holotrichia consanguinea*)	175 + 175- 200 + 200	437.5-500	1000 – 1250	296
Groundnut	White Grubs (*Holotrichia serrata*)	100 + 100 to 120 + 120	250 – 300	1000	106
Cotton	Jassids and Thrips	50 + 50	125	500	36
Rice	Yellow Stem Borer (*Scirpophaga incertulas*), Leaf Folder (*Cnaphalocrocis medinalis*), Brown Plant Hopper (*Nilparvata lugens*)	40+40 – 50+50	100-125	500	7
Chilli	Thrips (*Scirtothrips dorsalis*), Fruit Borer (*Helicoverpa armigera*)	30+30 – 40+40	75 – 100	500	7
Fipronil 07 % + Hexythiazox 02 % w/w SC					
Chilli	Mites and Thrips	70 + 20	1000	500	07
Fipronil 04 % + Thiamethoxam 04 % w/w SC					
Rice	Brown Plant Hopper, Green Leaf Hopper & White Backed Plant Hopper	44 + 44	1100	500	45

Fipronil 7 % + Tolfenpyrad 15 % + Abamectin 1.25 % SC					
Okra	Aphids, Jassids, Thrips, Whitefly, Mites and Fruit borer	52.5 + 112.5 + 9.37	750	500	3
Chilli	Aphids, Thrips, Mites and Fruit borer	52.5 + 112.5 + 9.37	750	500	10
Fipronil 15% + Flonicamid 15% WDG					
Cotton	Aphid, Jassid, thrips, whitefly, mealy bug and bollworm	60 + 60	400	500	33
Paddy	Brown plant hopper, green leaf hopper, stem borer and leaf folder	60 + 60	400	500	30
Flonicamid 11.7% + Diafenthiuron 36% WG					
Cotton	White fly, Aphids, Jassids	73.13+225	625	500	80
Tomato	White fly, Aphids, Jassids	73.13+225	625	500	5
Flonicamid -20% + Fipronil- 8% SC (W/W)					
Cotton	Aphids, Jassids, Thrips & White fly	75 + 30	375	500	35
Fluxametamide 5.81% + Bifenthrin 5.81 % EC (11.62% EC w/w) (12% EC w/v)					
Chilli	Fruit borer (*Helicoverpa armigera*)	37.5 + 37.5	625	375	7
	Thrips (*Scirtothrips dorsalis*) Whitefly (*Bemisia tabaci*)	30 + 30	500	375	7
Tomato	Fruit borer (*Helicoverpa armigera*)	37.5 + 37.5	625	375	7
	Leaf miner (*Liriomyza trifolii*)	30 + 30	500	375	7
Brinjal	Shoot and Fruit Borer (*Leucinodes orbonalis*)	37.5 + 37.5	625	375	7
	Whitefly (*Bemisia tabaci*), Jassid (*Amrasca bigutulla bigutulla*)	30 + 30	500	375	7
Flubendiamide 5.75% + Emamectin Benzoate 1.92%					
Maize	Fall army worm (*Spodoptera frugiperda*), stem borer (*Chilo partellus*)	37.5+12.5	625	500	46
Soybean	Stem fly (*Melanagromyza sojae*), pod borer (*Chrysodeixis acuta*) and Semi looper (*Helicoverpa armigera*)	37.5+12.5	625	500	42

Fluxametamide 3.8% w/w + Pyridaben 9.5 % w/w SC					
Brinjal	Leaf hopper – *Amrasca biguttula biguttula* Thrips – *Thrips palmi* Fruit & shoot Borer – *Leucinodes orbonalis* Red spider mite – *Tetranychus urticae* Whitefly – *Bemisia tabaci*	30 + 30	500	375	5
Chilli	Thrips- *Scirtothrips dorsalis,* Fruit borer- *Helicovepra armigera,* Tobacco caterpillar- *Spodoptera litura*. Yellow mite – *Polyphagotarsonemus latus*	38 + 95	1000	500	5
Hexythiazox 3.5% + Diafenthiuron 42% WDG					
Chilli	Mites, Thrips, Jassids, Aphids & White fly	22.75 + 273	650	500	07
Imidacloprid 18 + Carboxin 22.5% + Thiram 22.5					
Wheat	Aphids, Termites	1.89-2.52	3.4	-	-
Imidacloprid 18.50 % + Hexaconazole 01.50 % FS					
Groundnut	Termites, Thrips, Jassids, Root grubs, Collar rot, Stem rot, Tikka leaf spot & Rust	37 + 3	200	Not applicable	This is usedas seed dresser
Wheat	Termites, Aphids, Smut, Rust	37 + 3	200	Not applicable	This is usedas seed dresser
Imidacloprid 06 % + Lambda-cyhalothrin 04 % SL					
Paddy (Rice)	Stem borer, Hispa, Plant hopper, Gundhi bug	18 + 12	300	500	10
Indoxacarb 5% + Fipronil 5 % W/w SC					
Chilli	Thrips	37.5+37.5 – 50+50	750-1000	500	5
Cabbage	Diamond back Moth	37.5+37.5 – 50+50	750-1000	500	7

Indoxacarb 14.50 % + Acetamiprid 07.70 % w/w SC					
Cotton	Jassids, Whitefly, Bollworms	88.8 – 111	400 – 500	500	30
Chilli	Thrips, Fruit borer	88.8 – 111	400 – 500	500	05
Indoxacarb 10.0% + Thiamethoxam 10.0% WG					
Tomato	Whitefly and fruit borer	75 + 75	750	500	5
Paddy	Brown plant hopper, yellow stem borer and leaf folder	50 + 50	500	50	14
Isoprothiolane 28% + Fipronil 5% EC					
Rice	Stem borer, Brown plant hopper, Green leaf hopper, Whorl maggot	280+50	1000	500	58
Lambda-Cyhalothrin 5% + Clothianidin 5% + Carbendazim 37% SC					
Chilli	Cercospora leaf spot (*Cercospora capsica*), Chilli thrips (*Scirtothrips dorsalis*)	(25+25+185) to (31.25+31.25+231.25)	500 - 625	375	3
Lambda cyhalothrin 10% + Lufenuron 12% SC					
Chilli	Chilli fruit borer (*Helicoverpa armigera*),Tobacco caterpillar - *Spodoptera litura*), Chilli Thrips (*Scirtothrips dorsalis*)	25+30	250	500	33
Tetraniliprole 16% + Lambda-cyhalothrin 10% OD					
Paddy	Rice stem borer (Scirpophaga incertulas), Leaf folder (Cnaphalocrocis medinalis),	40+ 25.0	250	500	65
Lufenuron 20% + Chlorantraniliprole 10% + Pyriproxyfen 5% w/w WG					
Chilli	Thrips (*Scirtothrips dorsalis*) Mites (*Polyphagotarsonemus latus*) and Fruit borers (*Helicoverpa armigera* and *Spodoptera litura*)	22.50+ 11.25+ 5.63	112.50	500	19
Lufenuron 4% + Emamectin Benzoate 1.5% EC					

Maize	Fall armyworm (*Spodoptera litura*)	(30+11.25) to (46+17.25)	750-1150	400-500	35
Soybean	Spodoptera spp. (*Spodoptera litura*)	(25+9.37)	625	375	37
METHOXYFENOZIDE 20% + CHLORANTRANILIPROLE 5% SC					
Groundnut	Tobacco caterpillar (Spodoptera litura), American bollworm (*Helicoverpa armigera*) and Groundnut leaf miner (Aproaerema modicella)	100 + 25	500	500	27
METHOXYFENOZIDE 5% + CHLORPYRIPHOS 25% SE					
Bengal gram	Pod borer (*Helicoverpa armigera*)	100 + 500	2000	500	15
METHOXYFENOZIDE 18% + EMAMECTIN BENZOATE 1.8% SC					
Chilli	Fruit borer (*Helicoverpa armigera*), Thrips (*Thrips tabaci*) and Mites (*Polyphagotarsonemus latus*)	90 + 9	500	500	3
Paddy	Stem borer (*Scirpophaga incertulas*) Leaf folder (*Cnaphalocrocis medinalis*)	90+9	500	500	34
Nitenpyram 15.0 % + Fipronil 20.0% w/w WP					
Cotton	Thrips (*Thrips tabaci*) and Leaf hoppers (*Amrasca biguttula biguttula*)	140	400	500	70
Novaluron 05.25 % + Emamectin Benzoate 0.9% SC					
Cabbage	Diamond Back Moth (*Plutella xylostella*), Tobacco Caterpillar (*Spodoptera litura*)	45.94 + 7.87	875	500	3
Chilli	Gram Pod Borer (*Helicoverpa armigera*), Tobacco Caterpillar (*Spodoptera litura*)	45.94 + 7.87	875	500	3
Red Gram	Gram Pod Borer (*Helicoverpa armigera*)	45.94 + 7.87	875	500	25

Rice	Stem Borer (*Scirpophaga incertulas*)	78.75 + 13.50	1500	500	32
Brinjal	Shoot & Fruit Borer	78.75 + 13.50	1500	500	5
Chickpea	Pod Borer	45.94 + 7.88	875	500	11
Maize	Fall armyworm (*Spodoptera frugiperda* & Stem borer (*Chilo partellus*)	78.75+13.50	1500	500	27
Novaluron 05.25 % + Indoxacarb 04.50 % SC					
Tomato	Fruit borer (*Helicoverpa armigera*) & Leaf eating caterpillar (*Spodoptera litura*)	43.31 + 37.13-45.94 + 39.38	825 – 875	500	05
Chickpea	Gram pod borer (*Helicoverpa armigera*)	43.31 + 37.13-45.94 + 39.38	825 – 875	500	9
Soybean	*Spodoptera* spp., *Helicoverpa armigera* and *Semilooper*	43.31 + 37.13-45.94 + 39.38	825 – 875	500	14
Pigeon pea (Red Gram/Arhar/Tur)	Pod borer complex (*Helicoverpa armigera & Melanogromyza abtuse)*	43.31 + 37.13-45.94 + 39.38	825 – 875	500	25
Chilli	Fruit borer complex (*Helicoverpa armigera, Spodoptera litura)*	43.31 + 37.13-45.94 + 39.38	825 – 875	500	7
Black gram	Black gram pod borer complex (*Etiella zinckenella*, *Spodoptera litura* and *Maruca vitrata*)	43.31 + 37.13-45.94 + 39.38	825 – 875	500	14
Rice (Paddy)	Rice leaf folder (*Cnaphalocrocis medinalis*)	22.97 + 19.69	437.5	500	40
Groundnut	*Helicoverpa armigera & Spodoptera litura*	45.94 + 39.38	875	500	34

NOVALURON 9.45% + LAMBDA-CYHALOTHRIN 1.9% ZC					
Red gram	Pod borer (*Helicoverpa armigera*), Leaf hopper (Empoasca kerri)	75+15	750	500	39
Tomato	Tomato fruit borer (*Helicoverpa armigera*), Tobacco caterpillar (Spodoptera litura)	75+15	750	500	7
Cabbage	Diamond back moth (Plutella xylostella), Tobacco caterpillar (Spodoptera litura)	75+15	750	500	7
Nitenpyram 20 % + Pymetrozine 60% DF for formulation indigenous manufacture					
Paddy	Brown plant hopper (*Nilaparvata lugens*) and White backed plant hopper (*Sogatella furcifera*)	30 + 90	150	500	40
Phenthoate 45% + Cypermethrin 6% EC					
Paddy	Yellow Stem Borer, Leaf Folder and Brown Plant Hopper	450+60	1000	500	At the end of the Harvest
Profenofos 40 % + Cypermethrin 04 % EC					
Cotton	Bollworm complex	440 – 660	1000 – 1500	500 – 1000	14
Sugarcane	Early shoot borer	440	1000	500	138
Profenofos 40 % + Cypermethrin 04 % EW					
Rice	Yellow stem borer (*Scirpophaga incertulas*) and Leaf folder (*Cnaphalocrocis medinalis*)	440	1000	500	25
Phenthoate 35% + Emamectin benzoate 3.5% w/w EC					
Chilli	Mites, Thrips and fruit borer	262.5g+26.25g	750	500	5

Profenofos 50% + Fenpropathrin 5% EC					
Cotton	Pink Bollworm (*Pectinophora gossipiella*), Thrips (*Thrips tabaci*), Aphids (*Aphis gossypii*), Jassids (*Amrasca biguttula*) and Whitefly (*Bemisia tabaci*)	825 (750 + 75)	1500	500	22
Profenofos 40 % + Fenpyroximate 02.50 % w/w EC					
Chilli	Thrips, Mites, Fruit borer	0.4 + 0.025	1000	500	07
Brinjal	Whitefly, Mites (*Tetranychus urticae*), Fruit & Shoot Borer	0.4 + 0.025	1000	500	07
Profenofos 40% (w/w) + Lambda Cyhalothrin 1.5% (w/w) EC					
Cotton	Aphids, thrips, Leafhopper, Whitefly, Bollworm	400 + 15	1000	500-1000	15
Propargite 50 % + Bifenthrin 5 % w/w SE					
Okra	Mite , White fly & Jassids	594 + 59.4 – 621 + 62.1	1100 – 1150	500	05
Tomato	Mite , White fly & Jassids	594 + 59.4 – 621 + 62.1	1100 – 1150	500	05
Propargite 42 % + Hexythiazox 2 % EC					
Tea	Red spider Mites	525 + 25	1250	400 – 500	07
Chilli	Yellow Mite	525 + 25	1250	500	7
Apple	European Red Mite	4.2 + 0.2 (g/10 L water)	10 (ml/10 L water)	10-15 L/tree depending on tree canopy	15
Brinjal	Mite	525+25	1250	500	7
Pymetrozine 29.2 % + Dinotefuran 11.7 % w/w WDG					
Rice	Brown plant hopper (Nilaparvata lugens)	146 + 58.5	500	500	22
Pymetrozine 14.8% + Fipronil 7.5% + Dinotefuran 4.8% w/w SC					
Paddy	Stem Borer, Brown Plant Hopper, White	129.5+65.62 + 42	875	500	25

	Backed Plant Hopper, Green Leaf Hopper				
Pymetrozine 30.0% + Tebuconazole 37.0% WG					
Paddy	Brown plant hopper (Nilaparvata lugens), and diseases viz., Sheath blight, Brown spot, Blast, and Glume discoloration (Dirty panicle)	150 +185	500	500	33
Pyrifluquinazon 6% + Buprofezin 30% SC					
Cotton	Whiteflies and Jassids	60 + 300	1000	500	14
Pymetrozine 30% + Dinotefuran 10% + Pyraclostrobin 20% WG					
Paddy	Brown plant hopper (Nilaparvata lugens), white backed plant hopper (*Sogatella furcifera*), and blast disease	112.5 + 37.5 + 75	375	500	22
Pymetrozine 25.0% + Thiamethoxam 17.5% + Hexaconazole 12.5% WG					
Paddy	**Pests:** Brown Plant Hopper, White Backed Plant Hopper, Green Leaf Hopper, Yellow Stem Borer, Leaf Folder **Diseases:** Sheath Blight, Leaf Blast	100 + 70 + 50	400	500	19
Pymetrozine 11.3% w/w + Triflumezopyrim 1.9% w/w SC					
Rice	Brown plant hopper, White backed plant hopper, Green leafhopper	120 + 20	1000	500	31
Pyraclostrobin 10% + Fipronil 5% w/v SC					
Rice	Yellow stem borer, Leaf folder, Brown Plant Hopper (BPH), Sheath Blight, Leaf blast	50+100	1000	500	53
Chilli	Thrips, Chilli fruit Borer, Powdery Mildew, Anthracnose/ Fruit Rot	50+100	1000	500	7
Pyraclostrobin 3.5% + Thiram 15.0% + Clothianidin 22.5% FS					
Ground nut	Aphids, Jassids, termites and white grub	2.45 + 10.5 + 15.75	70 ml / 10 Kg of seed	Not applicable for seed treatment	

(Seed Treatment)					
Pyriproxyfen 05 % + Fenpropathrin 15 % EC					
Cotton	Bollworms	25 + 75- 37.5 + 112.5	500-750	500-750	14
	Whitefly	60 + 60	600	500	19
Brinjal	Whitefly, Shoot & fruit borer	25 + 75 - 37.5 + 112.5	500 - 750	500 – 750	07
Okra (Bhindi)	Whitefly, Fruit borer	25 + 75- 37.5 + 112.5	500 – 750	500 – 750	07
Chilli	Whitefly, Fruit borer	25 + 75- 37.5 + 112.5	500 – 750	500 – 750	07
Pyriproxyfen 05 % + Diafenthiuron 25 % SE					
Cotton	Whitefly (*Bemisia tabaci*), Thrips (*Thrips tabaci*), Jassid (*Amrasca biguttula biguttula*), Aphid (*Aphisgossypi*)	50 + 250- 62.50 + 312.5	1000 – 1250	500	35
Chilli	Whitefly (*Bemisia tabaci*), Thrips (*Thrips tabaci*) and Mites (*Polyphagotarsonemus latus*)	37.5 + 187.5 – 50 + 250	750 - 1000	500	10
Pyriproxyfen 10 % + Bifenthrin10 % w/w EC					
Cotton	Whitefly	60 + 60	600	500	19
Chilli	Yellow mites, Thrips, whitefly, fruit borer	125+125	1250	500	15
Brinjal	Whitefly, Shoot & Fruit Borer	125+125	1250	500	15
Soybean	Whitefly	100+100	1000	500	35
Green gram	Whitefly	100+100	1000	500	25
Pyriproxyfen 8% + Dinotefuran 5% + Diafenthiuron 18% SC					
Brinjal	Whitefly, Jassid, Thrips and Aphids	48+30+108 to 66+41.25+148.5	600-825	500	8-10
Cotton	Whitefly, Jassids, Thrips and Aphids	48 + 30 + 108	600	500	48

Chilli	Whitefly, Jassids, Thrips and Aphids	66 + 41.25 + 148.5	825	500	5
Spinetoram 5.66% + Methoxyfenozide 28.3% w/w SC					
Rice	Stem Borer, Leaf Folder	135-144	375-400	500	56
Cabbage	Diamond Back Moth, Leaf Eating Caterpillar	135-144	375-400	375-500	10
Cauliflower	Diamond Back Moth, Leaf Eating Caterpillar	135-144	375-400	375-500	10
Blackgram	Pod Borer (*H. armigera*), Spotted Pod Borer (*M. vitrata*)	135-144	375-400	375-500	48
Spiromesifen 15% + Hexythiazox 2.2 % + Abamectin 1.15 % w/ SC					
Chilli	Yellow mites (*Polyphagotarsonemus latus*) and Black thrips (*Thrips parvispinus*)	93.75 + 13.75 + 7.1875	625	500	16
Spirotetramat 11.01 % + Imidacloprid 11.01 % w/w SC					
Okra (Bhindi)	Red spider mites	60 + 60	500	500	03
Brinjal	Whitefly, Red spider mites	60 + 60	500	500	05
Mango	Mealy bug	0.018%	0.075%	Spray fluid as required depending upon size of tree.	15
Cotton	Mealy bug	75+75	625	500	22
Grapes	Thrips, Mites and Mealy Bugs	90 + 90	750	1000	60
Spiropidon 30% w/w + Acetamiprid 24% w/w WG					
Tomato	Thrips (*Thrips tabaci*), Aphids (*Aphis gossypii*), Whiteflies (*Bemisia tabaci*)	108 (Spiropidion 60 + Acetamiprid 48) – 135 (Spiropidion 75 + Acetamiprid 60)	200 - 250	500	5
Cotton	Mealy bugs (*Phenococcus solenopsis*, Thrips (*Thips tabaci*), Jassids (*Amrasca devastans*),	108 (Spiropidion 60 + Acetamiprid 48) – 135 (Spiropidion 75 + Acetamiprid 60)	200 - 250	500	85

	Aphids (*Aphis gossypii*), Whiteflies (*Bemisia tabaci*)				
SULFOXAFLOR 7.5% + BUPROFEZIN 15% SC					
Paddy	Brown Plant Hopper (BPH)	60 + 120 – 75 + 150	800 - 1000	500	26
Tetraniliprole 0.4 % w/w + Fipronil 0.6 % w/w GR					
Rice	Stem borer (*Scirpophaga incertulas*) and leaf folder (*Cnaphalocrocis medinalis*)	40 + 60 – 50 + 75	10 - 12.5	NA	65
Sugarcane	Early shoot borer (*Chilo infuscatellus*) and White grub (*Holotrichia serrata*)	50 + 75 – 60 + 90	12.5 – 15	NA	314
Tetraniliprole 10.81 % w/w + Spirotetramat 21.62 % w/w SC					
Okra	Fruit & Shoot borer, Aphids, Whitefly and Mites	45 +90	375	500	3
Cabbage	Diamond back moth, Tobacco caterpillar and Aphids	45 +90	375	500	7
Chilli	Fruit Borer, Thrips, Whitefly and Mites	45 +90	375	500	5
Brinjal	Shoot and Fruit borer, Whitefly and Mites	45 +90	375	500	5
Tomato	Fruit Borer, Leaf miner, Thrips and Whitefly	45 +90	375	500	5
Tetraniliprole 10.08 % w/w + Thiacloprid 30.25% w/w SC					
Rice	Stem Borer and Leaf Folder	37.5 + 112.5	312.5	500	43
Thiamethoxam 12.60 % + Lambda-cyhalothrin 09.50 % ZC					
Cotton	Jassids, Aphids, Thrips, Bollworms	44	200	500	26
Maize	Aphid, Shoot fly, Stem borer	27.50	125	500	42
Groundnut	Leaf hopper, Leaf eating caterpillar	27.5	150	500	28
Soybean	Stem fly, Semilooper, Girdle beetle	27.50	125	500	48
Chilli	Thrips, Fruit borer	33	150	500	03

Tea	Tea mosquito bug, Thrips, Semilooper	33	150	400	01
Tomato	Thrips, Whiteflies & Fruit borer	27.5	125	500	05
Citrus	Psyllids, Thrips and Leaf miner)	33.15	150	500	10
Thiamethoxam 0.4% + Bifenthrin 0.8% GR					
Groundnut	White grub & Termite	48+96	12	-	105
Thiamethoxam 00.90 % + Fipronil 00.20 % w/w GR					
Ground nut	White grub, Termite	108 + 24 - 135 + 30	12.15	106	48
Sugarcane	White grub (*Holotrichia consanguineo*) and Termite (*adontotermes obesus*)	135+30	15	-	296
Thiamethoxam 1.25% + Fipronil 1.25% + Chlorantraniliprole 0.60% w/w GR					
Paddy (Rice)	Yellow stem borer (*Scirpophaga incertulas*) Leaf folder (*Cnaphalocrocis medinalis*)	56.25 + 56.25 + 27	4500	-	77
Thiocyclam Hydrogen Oxalate 3.0% + Clothianidin 1.2% GR					
Paddy	Yellow Stem Borer (*Scirpophaga incertulas*), Leaf Folder (*Cnaphalocrocis medinalis*), Brown Plant Hopper (*Nilaparvata lugens*), and White Backed plant Hopper (*Sogatella furcifera*).	300 + 120 – 375 + 150	10000-12500	-	56
TOLFENPYRAD 15% + BIFENTHRIN 7.5% SE					
Cotton	Leaf hoppers, Aphids, Thrips, Pink bollworm	112.50 + 56.25	750	500	54
Transplanted Paddy	Brown plant hopper (Nilaparvata lugens) Green leaf folder (Nephotettix nigropictus) Stem borer (Scirpophaga incertulus) Leaf folder	112.50 + 56.25	750	500	35

	(Cnaphalocrosis medinalis)				
Tolfenpyrad 18.75% + Emamectin Benzoate 0.94% SC					
Cauliflower	Diamond black moth (*Plutella xylostella*) Tobacco caterpillar (*Spodoptera litura*)	140 + 7	700	500	7
Brinjal	Shoot & Fruit borer (*Leucinodes orbonalis*) Jassids (*Amrasca biguttula biguttula*)	140 + 7	700	500	3
Tolfenpyrad 30 % + Pyriproxyfen 10 % + Acetamiprid 4% w/w EC					
Chilli	Aphids, Black Thrips, Whitefly and Jassids	150 + 50 + 20	500	500	7
Triflumezopyrim 4.79% + Spinetoram 8.62% w/w SC					
Paddy	Brown Plant Hopper (*Nilaparvata lugens*), White Backed Plant Hopper (*Sogatella furcifera*), Stem borer (*Scirpophaga incertulas*)	70 (25+45)	500	500	21
Triflumezopyrim 3.74 % + Benzpyrimoxan 14.02 % w/w SC					
Rice	Brown plant hopper (*Nilaparvata lugens*), white backed plant hopper (*Sogatella furcifera*) and green leaf hopper (*Nepotettix virescens*)	95 (25 + 70)		500	21
Triflumezopyrim 10% w/w + Spinetoram 12% w/w WG					
Paddy	Brown Plant Hopper (*Nilaparvata lugens*), White Back Plant Hopper (*Sogatella furcifera*) Leaf Folder (*Cnaphalocrosis medinalis*)	55 (25 + 30)	250	500	21
PUBLIC HEALTH USE					
Pest	**Habitat**	**a.i. (mg/m^2)**	**Formulation (gm)**		**Dilution (Ltr.)**
Alphacypermethrin 05 % WP					
Adult Mosquito	-	25 (2 cycles application torepeat	Dilute 250 gm of Alphacypermethrin5% WP		250

		after 3month)	in 10 litres of water tocover 500 sq m area.		
	-	40 (single cycle application)	Dilute 250 gm of Alphacypermethrin 5% WP in 10 litres of water tocover 500 sq m area.		400
Alphacypermethrin Impregnated long lasting nets 00.667 % w/w (200 mg/m^2) (For Import only)					
Ready to use Impregnated Bed Net		To control mosquitoes under Public Health			
Bifenthrin 10.00%WP					
Adult Mosquito	-	25 (2 rounds of spraying 3 months apart	125	Dilute 125 gm of Bifenthrin 10% WP in 10liters of water to cover 500 m² areas.	-
Chlorpyriphos Methyl 40 % EC					
-	Used to control of adult vector mosquitoes				
Cyfluthrin 10 % WP					
Under Public Health Programme (Adult Mosquitoes)	-	25 (2 cycles application to be Repeated after 3 months.	250	Dilute 250 gm of Cyfluthrin 10% WP in 10 litres of water to cover 500 m2 areas.	
		40 (single cycles application)	400	Dilute 400 gm of Cyfluthrin 10% WP in 10 litres of water to cover 500 m2 areas.	
DDT 50 % WP					
Adult mosquitoes	-	1-2gm	-	-	-
Deltamethrin 00.15 % + Piperonyl 00.55 % EC					
Adult mosquitoes	-	Mosquitoes control underPublic Health	-	-	-
Deltamethrin 01.25 % w/w or 01.00 % w/v EC					
Insect	**Method of Application**	**Dosage/ha.**			

		a.i. (gm)	Formulation (ml)	Dilution in diesel Oil (Litre)
Adult Mosquitoes	Thermal fogging	0.50	50	10
	Ultra low volume application	0.50	50	0.50
Deltamethrin 02.50 % WP				
Adult Mosquitoes	For public health purpose only	625-1250 mg/50 m^2	25-50 g/50 m^2	1.5-2.5 Ltr./50 m^2
Deltamethrin impregnated Bed Net 55 mg/m^2 (For Import only)				
Ready to use insecticide Impregnated Bed net			Mosquitoes control under Public Health	
Diflubenzuron 02 % GR				
Name of the insect pest	**Habitat**	**Dosage/ha (Kg.)**	**-**	**Waiting period**
Mosquito larvae	Water bodies (Cess pits, Drains, Disused wells and Pools)	1.25-3.0	-	-
Fenitrothion 40 % WP				
Common name of pest	**a.i. (gm)**	**Formulation**	**-**	**Dilution in water (litres)**
Mosquitoes & files	400	1000	-	80
Lambda Cyhalothrin 9.7% w/w (10% W/V) CS				

Purpose and Target Pest	Spray Deposit Rate (mg a.i./ sq.m meter)	Gram a.i dose		Dose per 500 sq.m. meter surface area	
		Spray Deposit Rate (g.a.i/sq. meter)	g. a.i per 500 sq. mtrs	Formulation Dose (ml)	Water Volume (litre)
For public health use for controlling mosquitoes transmit malaria *(Anopheles culicifacies)*	25	0.025	12.5	1.25	10

Name of Insect Pest	Dosage			
	Infestation	Requirement mg a.i./sq.m	Formulation (ml)/ liter water	Spray solution (ml)/ sq m

Mosquitoes (*Aedes aegypti, Anopheles stephensi, Culex quinquefasciatus*); Housefly (*Musca domestica*), Cockroaches (*Periplaneta americana, Blattella germanica*) and Bed bugs (*Cimex hemipterus*)	Moderate	20	4	50
	High	25	5	50

Lambda-cyhalothrin 10 % WP

Pest	**Use**	**Dosage 500 m^3 floor area**		**Dilution in water (Litre)**
		a.i. (gm)	**Formulation (gm)**	
Mosquitoes	For public health only	7.50 - 15	75 - 150	10
Mosquito, housefly, cockroach	For household use	10	100	10

Malathion 25 % WP

Crop	**Common name of the pest**	**Dosage/m^2**			**Waiting Period (days)**
		a.i. (gm)	**Formulation (gm)**	**Dilution in water (Liter)**	
-	Adult mosquitoes	02/m^2	08/m^2	100	Repeat after 6-8 weeks

Novaluron 10 % EC

Place of Application	**Insect**	**Dosages**		**Waiting Period**
		a.i. (gm)	**Formulation (ml)**	
Clean surface water	*Anopheles stephensi, Aedes aegypti*	30	0.03 ml/m^2	Every 12 weeks
Polluted surface water	*Culex quinquefasciatus, Anopheles subpictus*	60	0.06 ml/m^2	Every 6th week

Pyriproxyfen 00.50 % GR

Breeding habitats	**Dosage/ha**		**Interval between application**
	a.i. (gm)	**Formulation (Kg.)**	
Clean water/Domestic containers	10 (0.01ppm)	2	08 weeks
Polluted/ Peri-domestic breeding habitat	20 (0.02ppm)	4	08 weeks

Pirimiphos methyl 50 % EC

Location	**Name of the pest**	**Dosage**	**-**	**Waiting period**

Mosquito breeding surface	Mosquito larvae	25 ml/ha	-	-

Spinosad 20.62% w/w EC

Regime of application	Target Pests	Dosage a.i.	Formulation applied in g/100 L, ml/100 L or ml/m^2 per unit area	Frequency of application
Polluted water stagnant, drains cement lined "U" shaped drains, slow running / stagnant drains and waste water drains	*Culex quinquefasciatus*	25 mg/m^2	0.121 ml / m^2	Weekly
Clean water containers, plastic barrels, cement tanks, barrels, earthen pots, plastic overhead tanks, and plastic containers	*Aedes aegypti* *Anopheles stephensi*	0.274 mg/L	0.133 ml /100 L	Weekly

SPINOSAD 7.48% w/w TABLET

Regime of application	Target Pests	Dosage a.i.	Formulation applied in g/100 L, ml/100 L or ml/m^2 per unit area	Frequency of application
Domestic clean / potable waters, Clean water containers, plastic barrels, cement tanks, barrels, earthen pots, plastic overhead tanks and plastic containers	*Aedes aegypti* *Anopheles stephensi*	1 tab /200 L or 1/2 tab /100 L	1 tab /200 L or 1/2 tab /100 L	6 Weeks

Spinosad 2.50% w/w GR

Polluted water stagnant, drains cement lined "U" shaped drains, slow running / stagnant drains and waste water drains	*Culex quinquefasciatus*	150 mg /m^2, 1.5 mg/L	6 g /m^2 or 60 Kg/Ha	2 weeks
Clean water containers, plastic barrels, cement tanks, barrels, earthen pots, plastic overhead tanks, and plastic containers	*Aedes aegypti* *Anopheles stephensi*	0.074mg/L	0.296 g /100 L	4 weeks

Temephos 1% granules			
Mosquito	Highly polluted (Drains, Cesspits)	200-500	20-50 Kg/Ha
	Moderately Polluted (Tidal waters, Marshes, Swamps, Etc)	100-200	10-20 Kg/Ha
	Clean (Standing water, Shallow Ponds, Lakes, Woodland pools)	50-100	5-10 Kg/Ha
Cyclops Sp	Ponds, Step wells	0.5-1.0	5 – 10 g

Temephos 50 % EC

Regime of application	**Common name of pest**	**Dosage/ha**		**Waiting period (days)**
		a.i. (g)	**Formulation (ml)**	
Mosquito larval treatment area, ponds, swamps, drainage, ditches,canals and other, Breeding areas.	Mosquitoes larvae	37.5 - 125	75 - 250	200

HOUSEHOLD INSECTICIDES

Alphacypermethrin 0.1 % w/w (RTU)

Common name of pest	**Dose/m^2 (a.i./mg)**	**Formulation (ml)**
Cockroaches, Adult mosquitoes, Adult houseflies	25 - 50	25 - 50

Alphacypermethrin 00.50 % Chalk	
Ready to use household insecticides	To control cockroaches
Allethrin 00.50 % Coil	
Ready to use household insecticides	Used to control of house hold flying insectlike houseflies and mosquitoes

Allethrin 00.50 % Mosquito Coil	
Ready to use household insecticides	To control of adult mosquitoes
Allethrin 00.20 % Coil Adult Mosquitoes	
Ready to use household insecticide	To control of mosquito
Allethrin 00.50 % Coil Adult Mosquitoes	
Ready to use household insecticide	To control of mosquito
Allethrin 04 % Mat Adult Mosquitoes	
Ready to use household insecticide	To control of mosquito
Allethrin 05 % Aerosol	
Ready to use household insecticide	To control of mosquito
Allethrin 03.60 % LV	
Ready to use household insecticide	To control of mosquito
Bifenthrin 00.05 % Mosquito coil (8 hours Min.)	
Ready to use household insecticide	Used to control adult mosquitoes

Bifenthrin 2.5% + Acetamiprid 2.1% w/w ME

Common name of pest	**Dosage**		**Dilution in water (L)**
	a.i. in gm	**Formulation (ml)**	
American Cockroach (*Periplaneta americana*) , German Cockroach (*Blattella germanica*), Housefly (*Musa domestica*), Bed bug (*Cimex hemipterus*) in house hold (houses, commercial establishments, etc.,)	0.115 g a.i per liter of water	2.5 (Dilute 2.5 ml of formulation with 1000 ml water; Spray @ 50 ml / m2 or 1000 ml/ 20 m2 surface area.)	1 liter

Clothianidin 50% WDG

Common name of pest	**Dosage**		**Dilution in water**
	a.i. in (mg/m^2)	**Formulation (mg/m^2)**	
Cockroach (*Periplanata americana, Blatella germanica*)	60	120	Dillute 120 mg of Clothianidin 50% WDG in 75 ml of water per m^2 area.
House fly (*Musca domestica*)			
Bedbugs (*Cimex spp*)			

Cyfluthrin 10 % WP

Common name of pest	Dosage		Use
	a.i. in mg/m^2	Formulation (gm/m^2)	
Adult mosquitoes, Cockroaches, Houseflies & Mosquitoes in house	25	0.250 for each spray	100 gm of Cyfluthrin 10% WP to be diluted in 8 liters of potable water 40 gm ofCyfluthrin 10% WP tobe diluted in 10% litres water.
	20	0.200 for each spray	100 gm of Cyfluthrin 10% WP to be diluted in 8 liters of potable water 40 gm of Cyfluthrin 10%WP to be diluted in 10% litres water.
Cyfluthrin 10 % WP			
For house hold use Cockroach HouseflyMosquitoes	25-40	250 - 400	Dilute 250-400 gm of Cyfluthrin 10% WP in10 litres of water to cover 500 m^2 areas.
Chlorpyriphos 02 % w/w EC			
Ready to use household insecticides		Used for protecting wood from the attack of termites & borers.	
Chlorpyriphos Methyl 40 % EC			
Used to control adult mosquitoes			
Cyphenothrin 5 % EC			

Pest	Dosage (Per square meters)		
	a.i (mg)	Formulation (ml)	Dilution in water
Cockroaches	5.0	1.0	Dissolve 500ml of formulated material in 10 ltr water to cover500 sq m area.
Housefly, Adult mosquitoes	0.2	0.004	Dissolve 4 ml of formulated material in 20 ltr water to cover 1000 sq m area.
Thermal fogging Indoor			
Anopheles stephensi, Culex quinquefasciatus and *Aedes aegypti*	0.5	0.01	Dissolve 5 ml of formulation in kerosene/water to cover 500 sq m area.
Outdoor			

Anopheles stephensi, Culex quinquefasciatus and *Aedes aegypti*	3.5	70	Dissolve 70 ml of formulation in kerosene/water to cover 1 hectare area.

Cyphenothrin 07.20 % VP w/w (For use by pest control operator only)	
American Cockroaches & German Cockroaches	To control of American Cockroaches &German Cockroaches (In house)
Cypermethrin 03 % Smoke Generator	
Ready to use household insecticide.	To control Cockroaches in house, hotels & warehouse.
Cypermethrin 01.00% Dust	
Ready to use household insecticide.	To control Cockroaches in house.
Cypermethrin 01 % Chalk	
Ready to use household insecticide.	To control Cockroaches in house.

Cyfluthrin 05 % EW				
Ready to use	Cockroaches, House flies, mosquitoes, in-house. Bed net impregnation	8.0 ml	1.0	50 ml diluted solution/m^2

Cyfluthrin 00.025 % + Transfluthrin 00.04 % Aerosol	
Ready to use	Used for controlling /repelling Mosquitoes, Houseflies & cockroaches in homes.
Transfluthrin 1 % w/w + Cypermethrin 0.2 % w/w Spray	
Ready to use house hold	Mosquitoes (*Culex quinquefasciatus, Aedes aegypti*), Houseflies (*Musca domestica*), Cockroaches (*Periplaneta americana, Blatella germenica)* and Ants(Red ants)
Deltamethrin 1.0% SB (Anti- Insect Enamel) (Insecticidal Paint formulation)	
Recommended for the control of Cockroaches, Houseflies, Mosquitoes and Ants in houses.	1 litre of insect control paint is sufficient for an area of 90 to 110 sq feet
Deltamethrin 02.50 % Flow	

Name of insectpest	**Type of use**	**Dosage /m^2 area of bed net**		**-**	**-**
		a.i.	**Formulation**		

Adult Mosquitoes	For impregnation of polyester , nylon and cotton bed net	25 mg	1 ml	-	-
Deltamethrin 02.50 % WP					
Name of insect pest	**Habitat**	**Dosage /m^2 area of bed net**			-
		a.i.	**Formulation**	**Dilution in water (Liter)**	
Lesser grain borerRice moth, Saw toothed grain beetle, Red flour beetle, Khapra beetle, Almond moth	Grain and seeds in stacks	30 mg/m^2	1.2 g/m^2	1 Liter for 30 m^2	-
Rice weevil	Grain and seeds in stacks	30 mg/m^2	1.2 g/m^2	1 Liter for 30 m^2	-
	Walls, Ceilings & Floor	30 mg/m^2	1.2 g/m^2	1 Liter for 30 m^2	-
Diflubenzuron 02 % Tablets					
Name of pest	**Habitat**	**Dosage**		**Dilution in water**	
Mosquitoes larvae	Unused Coolers	0.5-1.0 ppm		0.5-1.0 tablet in 40 liter water	
Diflubenzuron 0.1% W/W Termite Bait to be used by pest control operators only					
Termites	-	-		Bait mixed with water in 1:2 ratio	
Diflubenzuron 20 % + Deltamethrin 02% SC					
Name of the insectpest	**Habitat**	**Dosage/ha (kg.)**		**Waiting period**	
House fly maggot	Poultry Manure & kitchen garbage	1.50-2.00 ml/liter water (5 litreof water /10 m^2		-	
Diflubenzuron 25 % WP					

Name of pest	Habitat	Dosage	Dilution in water
Mosquitoes larvae	Clean surface water	25 - 50 g a.i./ha	-
	Polluted surface water	50 - 100 g a.i./ha	-
	Sewage pits, Soak pits, Latrines, Septic tanks	01.0 mg a.i./liter	-
House fly maggots control	In poultry manure garbage, filth & dumping areas	5.0 gm/10 m^2	05 liters water/10m^2

Dinotefuran 0.5% RB Gel	
Ready to use house hold	Used for controlling American cockroach (*Periplaneta americana*) and German Cockroach (*Blattella germanica*)

Deltamethrin 00.05 % + Allethrin 00.04 % w/w EC		
Common name of house hold insect	**Dosage/ha**	
	g a.i.	**Formulation (ml)**
Cockroaches, House flies, Mosquitoes	12.5 - 25.0	25 - 50

Deltamethrin 02.50 % + D-trans allethrin 02 % w/w EC		
Insects	**Dosage/m^2**	
	a.i. (mg)	**Quantity of solution(ml)**
Cockroach, Houseflies, Mosquitoes	12.5 - 25.0 + 10 - 20	25 - 50

Deltamethrin 00.02 % + Allethrin 00.13 % w/w	
Ready to use	To control cockroaches, mosquitoes andflies

Deltamethrin 00.50 % w/w Chalk	
Ready to use household insecticide	To control Cockroaches, ants and bedbugs

D-Trans Allethrin 00.10 % + Permethrin 00.03 % + Imiprothrin 00.02 % Aerosol w/w (all InsectKiller Aerosol)	
Ready to use	To control cockroaches, mosquitoes andhouse flies

Deltamethrin 01 % RTU

Ready to use household insecticide	To control Cockroaches in house. One litre of insect control of paints sufficientfor an area of 22 sq. meters. Two coats of insect control paint are recommendedgiving 18 hours of drying between the coats.
D-Trans Allethrin 02% Mosquito Mat	
Ready to use household insecticide.	To control Adult Mosquitoes in house.
D-Trans Allethrin 00.10 % w/w Mosquito Coil	
Ready to use household insecticide.	To control and repel of Adult Mosquitoes in the house.
D-Allethrin 21.97 % w/w Mosquito Mat.	
Used to control Adult Mosquitoes	Open Area like Park, Garden and FarmHouses etc. only.

Benzoate 00.10 % w/w Gel

Name of Insect/Pest	Dose (g a.i.)	Formulation Dose	Application Usages
American Cockroach (*Periplaneta americana*)	0.001 g a.i./m^2	1.0 gm of Gel Bait/m^2 (2-5 spots)	Place "Ready to Use Gel Bait" (RB) for use as spot or cracks and crevices treatment in residential institutional, commercial andindustrial areas e.g. application at or nearharborage or aggregation areas, such ascorners, areas where cockroaches forage or crack and crevices, holes, hidden surfaces, any other places where cockroaches are typically known to hide etc. for the control of cockroaches.
German Cockroach (*Blattella germanica*)	0.001 g a.i./m^2	1.0 gm of Gel Bait/m^2 (1-2 spots)	Place "Ready to Use Gel Bait" (RB) foruse as spot or cracks and crevices treatment in residential institutional, commercial andindustrial areas e.g. application at or near harborage or aggregation areas, such as corners, areaswhere cockroaches forage or crack andcrevices, holes, hidden surfaces, any otherplaces where cockroaches are typically known to hide etc. for the control of cockroaches.

Fenitrothion 20 % OL

Name of Pest	Dose (g a.i.)	Formulation (ml)	Instruction for use
Bedbug (*Cimex* spp.)	2	10	Take 10 ml of BILFOL 20 and dilute in 200 ml ofkerosene. Apply spotspray thoroughly in all bed bug infested areas like charpoy furniture etc. taking care that the spray is the directed into cracks and crevices where bedbugs are hiding. 200ml of spray wash will approx cover 10 m^2 it can also be applied with a brush where ever bedbugsoccur.

Fipronil 00.03 % & 0.5 % Gel	
Ready to use household insecticide	Used to control of German & AmericanCockroaches.

Fipronil 00.05 % GEL		
House hold	**Common name of the pest**	**Dosage/m 2**
House hold	American Cockroach (*Periplanata mericana*), German cockroach (*Blattella germanica*)	0.03 g (in a bait gun), 3-4 spot/m^2
House hold	German cockroach (*Blattella germanica*)	*(100 mg spot= approx 5 mm diameter)* Low Density – 1 spot /M^2 High Density – 2 spots / M^2
	American Cockroach (*Periplanata americana*)	Low Density – 2 spots /M^2 High Density – 3 spots / M^2

Imiprothrin 00.10 % + Cyphenothrin 00.13 % w/w	
Ready to use	Used for controlling cockroaches inhomes.
Imiprothrin 00.70 % + Cypermethrin 00.20 % w/w Aerosol	
Ready to use household insecticides	Used against Cockroaches.
Imiprothrin 00.05 % + Cypermethrin 01 % CL	
Ready to use	Used for controlling cockroaches inhouses.
Imidacloprid 00.03 % w/w Gel	

Species	Recommended Dose
Pharaoh ant (*Monomorium pharaonis*), Small black ant (*Monomorium indicum*), Crazy ant (*Paratrechina longicomis*), Ghost ant (*Tapinoma melanocephalum*)	Low infestation level (one spot of 200 mg/m^2 of infested area). Moderate to highinfestation level (one spot of 300 mg/m^2 of infested area).

Scoring of ant activity will be done based on the following:

Low activity=1-50 ants passing from a given point in the time period of one minute.

Medium activity=51-200 ants passing from a given point in the time period of one minute.

High activity= 201 ants passing from a given point in the time period of one minute.

Imidacloprid 02.15 % w/w Gel

Ready to use household insecticide	Used to control of German & American Cockroaches

Imidacloprid 21 % + Beta-cyfluthrin 10.50 % w/w SC

Name of Insect pests	Places	Dosage
American Cockroaches, German Cockroaches	Private Houses, Factories, Offices, Market places, Restaurants, Hotels, Shops, Ships, Hospital etc.	Dilute 04 ml of Imidacloprid 21.0% w/w + Beta-cyfluthrin 10.5% w/w SC with 01L of water. Apply 50 ml of this solution to spray per square meter area or apply 01 L of this solution to cover 20 square meter area.
Bed Bug	Hospitals, Houses, Commercial establishments, Hotels, Dormitories, Old age Homes, Hostels, etc.	
Housefly	Houses, Factories, Offices,Market places, Restaurants, Hotels, Shops, Ships, Hospital, Poultry, Diaries, Cattle sheds etc	
Adult Mosquitoes (Aedes sp., Anopheles sp. and Culex sp.)	Houses, factories, offices, market places, restaurants, hotels, shops, ships, hospitals etc.	Dilute 3 to 4 ml of Imidacloprid 21% w/w + Betacyfluthrin 10.5% w/w SC with 1 liter of water. Apply 50 ml of this solution to spray per sq.m. area. Or apply 1 liter of this solution to cover 20 sq.m. area

Lambda-cyhalothrin 00.50 % Chalk

Ready to use household insecticides	Used to control Cockroaches.

Lambda-cyhalothrin 02.43 % CS

Purpose and target pest	Dosage/m2 of netting		
	a.i. (mg)	Concentration of spray fluid	Quantity of spray fluid (ml)
Impregnation of bed nets	10.0	0.05%	1000 (depending on the type of the

to prevent attack from mosquitoes			net)
Lambda-cyhalothrin 02.43 % CS			
Common Name of pest	**Dosage**		
Adult mosquitoes, Adult house flies, Cockroaches	20 - 30 mg/m^2	10-15 ml/litres of water to cover 50 m^2area	
Lambda-cyhalothrin 02.43 % CS			
Target insect	**Dosage**		
	Mg a.i./m^2	**Method of application**	
Non-porous surfaces – Mosquitoes, Houseflies & Cockroaches	12.50	Mix 20 ml of product in 1 liter of water& spray the solution uniformly @ 25 ml/m^2 on non porous & @ 50 ml/m^2 onporous surfaces.	
Porous surfaces – Mosquitoes House flies& Cockroaches	25	Mix 20 ml of product in 1 liter of water& spray the solution uniformly @ 25 ml/m^2 on non porous & @ 50 ml/m^2 onporous surfaces.	
Lambda-cyhalothrin 02.43 % CS			
Name of pest	**Dosage/m^2**		
	a.i. (mg)	**Formulation (ml)**	**Dilution in water**
Cockroaches	50	1	Dissolve 500 ml of formulated material in 10 litre water to cover 500 square meter area.
Housefly, Adult mosquitoes	0.2	0.004	Dissolve 4 ml of formulated material in 20 litre water to cover1000 square meter area.
Indoor			
Anopheles stephensi, *Culex* q*uinquefasciatus*, *Aedes aegypti*	0.5	0.01	Dissolve 5 ml of formulated material in kerosene to cover 500 square meter area.
Outdoor			
Anopheles stephensi, Culex quinquefasciatus, Aedes aegypti	3.5	70	Dissolve 70 ml Formulation in kerosene to cover 1 hectare area.

Target Pest	Active Ingredient Dose (g a.i.)	Formulation Dose (ml)	Method of application (water volume)	Application Usage
Houseflies (*Musca domestica*) Mosquito (*Anopheles* spp.) American Cockroaches (*Periplaneta americana*) German Cockroaches (*Blattella germanica*)	0.375 – 0.5 g a. i. per Litre water (15 – 20 mg. a. i. per square meter)	15 – 20 per litre water	**For Low Pest Infestation (Maintenance Rate)** Mix 15 ml of the product in one litre of water and spray the solution uniformly @ 40 ml per square meter. **For High Rate Infestation (Cleanout Rate)** Mix 20 ml of the product in one litre of water and spray the solution uniformly @ 40 ml per square meter.	For use as indoor oroutdoor as a surface crack and crevice or spot spraytreatment in residential, institutional, commercial and industrial areas/establishment setc.
Anopheles stephensi, *Culex quinquefasciatus*, *Aedes aegypti*	0.5	0.01	Dissolve 5 ml of formulated material inkerosene to cover 500 square meter area.	
Outdoor				
Anopheles stephensi, Culex quinquefasciatus, Aedes aegypti	3.5	70	Dissolve 70 ml Formulation in keroseneto cover 1 hectare area.	
Malathion 02 % House Hold Spray				
Ready to use		To control of Bed, Bugs, Flies, Ants, Gnats, Mosquitoes, Moths and Cockroaches in houses.		
Metofluthrin 00.005 % (Mosquito Coil)-Min. 07 Hrs. Burning time				
Ready to use household insecticide.		To control of mosquitoes in houses.		
Metofluthrin 00.005 % (Mosquito Coil)-Min.12 Hrs.				
Ready to use household insecticide.		To control of mosquitoes in houses.		
METOFLUTHRIN 0.14% w/w AEROSOL				
This is ready to use aerosol insecticide		Mosquito species (Anopheles stephensi, Aedes aegypti, and Culex quinquefasciatus) and housefly Musca domestica		
Metofluthrin 0.32 % Liquid Vaporizer				
House hold		for Mosquitoes		

Metofluthrin 0.32 % Liquid Vaporizer (with 1% perfume)				
Household		For mosquitoes (*Anopheles stephensi, Aedes aegypti & Culex quinquefasciatus*)		
Novaluron 10 % EC				
Clean surface water	*Anopheles stephensi* *Aedes aegypti*	30	0.03 ml/m^2	-
Polluted water	*Culex quinquefasciatus* *Anopheles subpictus*	60	0.06 ml/m^2	-
Permethrin 02.00% (Olyset @ Net) w/w for Import only				
Ready to use household insecticides	For control of mosquitoes both indoors andoutdoors. After unpacking and before usingthe new bed net, keep it in and open place for 12 hrs away from the sunlight.			
Permethrin 2% w/w Aerosol (To be used in Aircraft only)				
Ready to use insecticides	For use inside aircraft prior to embarkation of passengers to control and prevent the spread of mosquitoes.			
Propetamphos 01 % Spray				
Ready to use household insecticide	To control of Cockroaches, Bed bugs, Flies,fleas, Mosquitoes & Silverfish.			
Propoxur 00.75 % + Cyfluthrin 00.025 % Aerosol				
Ready to use household insecticide	Cockroaches, Mosquitoes & Houseflies			
Propoxur 20 % EC				
Common name of pest	**Dose (g a.i.)**	**Formulation (ml)**	**Dilution in water (litres)**	
Flying insect-Mosquitoes, Files, Cockroaches , Bed bugs, Flash, Ticks crickets , Woodlice , Mite, Silver fish, Spider, Ants etc.	200	1000	40	
Pirimiphos-methyl 01 % Spray				
Location	**Pest**	**Dosage**	**Exposure period (min.hrs.)**	

Spot spray inhouses	Cockroaches , bed bugs, flea etc.	100 ml/1 m^2	01
Space spray inhouses	Mosquitoes, houseflies	50 ml/100 m^3	01
Pyrethrin 00.05 % + Malathion 01 %			
Insects		Used to control of Cockroaches, Mosquitoes and Flies.	
Propoxur 02 % Bait			
Ready to use household insecticides		Used to control of Cockroaches andFlies.	
Pyrethrin 00.20 % Spray			
Ready to use household insecticide		To control of Cockroaches, Houseflies,Mosquito and bugs	
Propoxur 01 % Spray			
Ready to use household insecticide		Used to control of Cockroaches, Houseflies and Adult Mosquitoes	
Prallethrin 01 % w/w Red Mosquitoes Mat			
Ready to use household insecticide.		Used to control of adult mosquitoes	
Prallethrin 00.04 % Coils (Min.11Hrs.)			
Ready to use household insecticide		Used to control mosquitoes in Houses	
Prallethrin 00.04 % Coils (Min.6 Hrs.)			
Ready to use household insecticide		Used to control mosquitoes in Houses	
Prallethrin 00.80 % w/w Red Mosquitoes Mat			
Ready to use household insecticide.		Used to control of Mosquitoes.	
Prallethrin 00.50 % w/w Mosquitoes Coil			
Ready to use household insecticide.		Used to control of adult mosquitoes.	
Prallethrin 01.20 % Mat			
Ready to use household insecticide.		Used to control of adult mosquitoes.	
Prallethrin 00.04 % w/w Mosquito Coil			
Ready to use household insecticide.		Used to control of adult mosquitoes.	
Prallethrin 19 % w/w VP			
Ready to use household insecticide.		Used to control of adult mosquitoes.	

Prallethrin 02.40 % w/w Liquid Vaporizer			
Ready to use household insecticide.		Used to control of Mosquitoes.	
Renofluthrin 0.025% w/w Mosquito Coil		Used to control adult mosquitoes of *Aedes aegypti*, *Anopheles stephensi* & *Culex quinquefasciatus.*	
Renofluthrin 1.0% W/W Incense Stick		For the control of mosquito species *Anopheles stephensi, Aedes aegypti, and Culex quinquqfasciatus* under household conditions.	
Renofluthrin 1.5% W/W Liquid vaporizer		For the control of mosquito species *Anopheles stephensi, Aedes aegypti, and Culex quinquqfasciatus* under household conditions.	
S-Bioallethrin 02.40 % Mosquitoes Mat			
Ready to use household insecticide.		Used to control of adult mosquitoes.	
Thiamethoxam 00.01 % w/w Gel Bait			
Common Name of the Insect/Pest	**Dose (g a.i.)**	**FormulationDose**	**Application/Usage**
Black Carpenter Ants (*Camponotus* spp.)	0.0001 g.a.i. per spot (2-4 spotsper square meter)	1.0 gm of gelbait per spot(2-4 spots persquare meter)	Locate the ant trails or location where ants are most active. Place" Readyto Use Gel Bait" (RB) for controlling ants for use as spot or cracks and crevices treatment in residential, Institutional, commercial and industrialareas e.g. application at ornear harborage oraggregation areas, such as corners areas where ants forage or crack and crevices, holes, hiddensurfaces any other places where ants are typically known to hide.
Transfluthrin 0.08 % w/w Aerosol			
Ready to use household insecticide		Used to control in household Mosquitoes(*Aedes aegypti, Culex quinquefasciatus*) and Housefly (*Musca domestica*).	
Transfluthrin 0.15 % w/w Mosquito Coil			
Ready to use household insecticide		To control / repel mosquitoes in houses	
Transfluthrin 0.75% Mosquito Agarbatti			
Ready to use household insecticide		Used to control of Adult Mosquitoes.	
Transfluthrin 00.88 % & 01.60 % Liquid Vaporizer			

Ready to use household insecticide.	Used to control of Adult Mosquitoes and House fly.
Transfluthrin 01.60 % Liquid Vaporizer (For 30 Nights (25 ml)	
Ready to use household insecticide.	Used to control of Adult Mosquitoes.
Transfluthrin 20 % w/w MV Gel	
Ready to use household insecticide.	Used to control of Mosquitoes in thehouse.
Transfluthrin 00.03 % w/w Mosquito Coil	
Ready to use household insecticide	Used for controlling/repelling of Mosquitoes in the house
Transfluthrin 01 % EU (Smoke generator)	
Use / recommendation	It is used for controlling/repellingadult mosquitoes in the houses (Effective for 6 hrs.)
Transfluthrin 01.20 % Liquid Vaporizer (For 60 Nights (45 ml) & 90 nights (67 ml.)	
Ready to use household insecticide	Used to control of adult mosquitoes
Transfluthrin 12 % AE	
Ready to use household insecticide.	Used to control/repel mosquitoes (Culex quinquefasciatus, Aedes aegypti) and houseflies (Musca domestica) in the houses (effective for 12 hours)
Zinc Phosphide 01 % bait (Household Product)	
To be ready to use household insecticide	To control Rats

Recommended chemicals by FAO for Locust Control

Sr. No.	Chemical	Dose (gram activeingredient per ha.)	
		Hoppers	Adults
1	Chlorpyriphos 20 % & 50 % EC	240	240
2	Deltamethrin 2.8 % EC & 1.25 % ULV	12.5	12.5
3	Diflubenzuron 25 % WP	60	NA
4	Fipronil 5 % SC & 2.92 % EC	6.25	6.25
5	Lambdacyhalothrin 5 % EC & 10% WP	20	20
6	Malathion 50 % EC & 25 % WP &96 %ULV	925	925
7	Fenitrothion is also recommended for the control of locust but only inscheduled desert area and public health but banned in agriculture.(refer copy of Gazette of India, S.O.706 (E) dated 03rdMay, 2007)		
8	Powder formulations are approved (RC-413) for control of desertlocust in Scheduled Desert Area: - Fenvalerate 0.4 % DP Malathion 5 % DP Quinalphos 1.5 % DP		