



- **National EMS Quality Alliance (NEMSQA) Performance Measures** (for additional information, see www.nemsqa.org)
 - *Pediatrics—03: Documentation of Estimated Weight in Kilograms*
 - *Trauma—01: Pain Assessment of Injured Patients*

References

1. American Burn Association. Advanced Burn Life Support (ABLS) Handbook; 2011
2. Chung K, Salinas J, Renz E, et al. Simple derivation of the initial fluid rate for the resuscitation of severely burned adult combat casualties: in Silico validation of the rule of ten. *J Trauma*. 2010;69 Suppl 1: S49–54
3. Dries DJ, Endorf FW. Inhalation injury: epidemiology, pathology, treatment strategies. *Scand J Trauma Resusc Emerg Med*. 2013; 21:31
4. Endorf FW, Gamelli RL. Inhalation injury, pulmonary perturbations, and fluid resuscitation. *J Burn Care Res*. 2007; 28(1):80-83
5. Fluid Rate charts (based on Parkland formula) and TBSA diagrams courtesy of the University of Utah Burn Center; 2014. As presented in [Appendix VI. Burn and Burn Fluid Charts](#)
6. Hettiaratchy S, Papini R. Initial management of a major burn: II— assessment and resuscitation. *BMJ*. 2004;329(7457):101–103. doi:10.1136/bmj.329.7457.101
7. McIntire AM, Harris SA, Whitten JA, et al. Outcomes Following the Use of Nebulized Heparin for Inhalation Injury (HIHI Study). *J Burn Care Res* 2017; 38:45-52
8. Miller AC. Influence of nebulized unfractionated heparin and N-acetylcysteine in acute lung injury after smoke inhalation injury. *J Burn Care Res*. 2009; 30:249–256

Revision Date

March 11, 2022