


Graphic 1. Prehospital External Hemorrhage Control Protocol



* Use of tourniquet for extremity hemorrhage is strongly recommended if sustained direct pressure is ineffective or impractical; Use a commercially produced, windlass, pneumatic, or ratcheting device, which has been demonstrated to occlude arterial flow and avoid narrow, elastic, or bungee-type devices; Utilize improvised tourniquets only if no commercial device is available; Do not release a properly applied tourniquet until the patient reaches definitive care

Apply a topical hemostatic agent, in combination with direct pressure, for wounds in anatomic areas where tourniquets cannot be applied and sustained direct pressure alone is ineffective or impractical; Only apply topical hemostatic agents in a gauze format that support wound packing; Only utilize topical hemostatic agents which have been determined to be effective and safe in a standardized laboratory injury model

Source: *Bulger et al. 2014*

References

1. Bedri H, Ayoub H, Engelbart J, Lilienthal M, Galet C, Skeete D. Tourniquet application for bleeding control in a rural trauma system: outcomes and implications for prehospital providers. *Prehosp Emerg Care*. 2021; DOI: 10.1080/10903127.2020.1868635
2. Bulger E et al. An evidence-based prehospital guideline for external hemorrhage control: American College of Surgeons Committee on Trauma. *Prehosp Emerg Care*. 2014;18(2):163–73
3. Charlton NP, et al. Pediatric Tourniquet Types: First Aid New TF SR. CoSTR.ILCOR.org [Internet]. Available from: <https://costr.ilcor.org/document/pediatric-tourniquet-types-first-aid-new-tf-sr>. Accessed March 11, 2022
4. Cornelissen M, Brandwijk A, Schoonmade L, Giannakopoulos G, van Oostendorp S, Geeraedts L. The safety and efficacy of improvised tourniquet in life-threatening hemorrhage: A systematic review. *European Journal of Trauma and Emergency Surgery*. 2020; 46:531–538
5. Cunningham A, Auerbach M, Cicero M, Jafri M. Tourniquet usage in prehospital care and resuscitation of pediatric trauma patients – Pediatric Trauma Society position statement. *J Trauma Acute Care Surg*. 2018;85(4):665–667
6. Doyle G, Taillac P. Tourniquets: a review of current use with proposals for expanded prehospital use. *Prehosp Emerg Care*. 2008;12(2):241–56
7. Kelly JR, Levy MJ, Reyes J, Anders J. Effectiveness of the combat application tourniquet for