



VI. Neurologic Status Assessment

Neurologic status assessment involves establishing a baseline and then trending any change in patient neurologic status. Glasgow Coma Score (GCS) is frequently used, but there are often errors in applying and calculating this score. With this in consideration, Glasgow Coma Score may not be more valid than a simpler field approach. Either AVPU (Alert, Verbal, Painful, Unresponsive) or only the motor component of the GCS may more effectively serve in this capacity.

Glasgow Coma Score

	Points	Pediatric	Adult
Eyes	1	No eye opening	
	2	Eye opening to pain	
	3	Eye opening to verbal	
	4	Eyes open spontaneously	
Verbal	1	No vocalization	No verbal response
	2	Inconsolable, agitated	Incomprehensible sounds
	3	Inconsistently consolable, moaning	Inappropriate words
	4	Cries but consolable, inappropriate interactions	Confused
	5	Smiles, oriented to sounds, follows objects, interacts	Oriented
Motor	1	No motor response	
	2	Extension to pain	
	3	Flexion to pain	
	4	Withdraws from pain	
	5	Localizes pain	
	6	Obeys commands	

Table X: AVPU

- A:** The patient is alert
V: The patient responds to verbal stimulus
P: The patient responds to painful stimulus
U: The patient is completely unresponsive