



- Pain score if applicable

Performance Measures

- If patient in pain, pain score change
- If patient is nauseated, symptom relief
- If patient is dehydrated, symptom relief or vital sign change

References

1. Coyne PJ, Viswanathan R, Smith TJ. Nebulized fentanyl citrate improves patients' perception of breathing, respiratory rate, and oxygen saturation in dyspnea. *J Pain Symptom Manage.* 2002;23(2):157–60
2. Farahmand S, Shiralizadeh S, Talebian MT, et al. Nebulized fentanyl vs intravenous morphine for ED patients with acute limb pain: a randomized clinical trial. *Am J Emerg Med.* 2014;32(9):1011–5
3. Portenoy RK, Mehta Z, Ahmed E. Cancer pain management with opioids: Optimizing analgesia. Post TW, ed. UpToDate. Waltham, MA: UpToDate Inc. <https://www.uptodate.com/contents/cancer-pain-management-with-opioids-optimizing-analgesia..> Accessed March 11, 2022
4. Prehospital Evidence Based Practice Program (PEP) [Internet]. Halifax, Nova Scotia: Dalhousie University - Division of Emergency Medical Services <https://emspep.cdha.nshealth.ca>. Accessed March 11, 2022
5. Shirk MB, Donahue KR, Shirvai J. Unlabeled uses of nebulized medications. *Am J Health Syst Pharm.* 2006;63(18):1704–16

Revision Date

March 11, 2022