



15. Lamhaut L, Dagron C. Comparison of intravenous and intraosseous access by pre-hospital medical emergency personnel with and without CBRN protective equipment. *Resuscitation*. 2010;81(1):65–8
16. Mahmoudian T, Zadeh MM. Comparison of intranasal midazolam with intravenous diazepam for treating acute seizures in children. *Epilepsy Behav*. 2004;5(2):253–5
17. McIntyre J, Robertson S, Norris E, et al. Safety and efficacy of buccal midazolam versus rectal diazepam for emergency treatment of seizures in children: a randomized controlled trial. *Lancet*. 2005;366(9481):205–10
18. McMullan J, Sasson C, Pancioli A, Silbergleit R. Midazolam versus diazepam for the treatment of status epilepticus in children and young adults: a meta-analysis. *Acad Emerg Med*. 2010;17(6):575–82
19. Mittal P, Manohar R, Rawat AK. Comparative study of intranasal midazolam and intravenous diazepam sedation for procedures and seizures. *Indian J Pediatr*. 2006;73(11):975–8
20. Mpimbaza A, Ndeezi G, Staedke S, Rosenthal PJ, Byarugaba J. Comparison of buccal midazolam with rectal diazepam in the treatment of prolonged seizures in Ugandan children: a randomized clinical trial. *Pediatrics*. 2008;121(1):58–64
21. Muchohi SN, Kokwaro GO, Ogutu BR, et al. Pharmacokinetics and clinical efficacy of midazolam in children with severe malaria and convulsions. *Br J Clin Pharmacol*. 2008;66(4):529–38
22. Muchohi SN, Obiero K, Newton CR, et al. Pharmacokinetics and clinical efficacy of lorazepam in children with severe malaria and convulsions. *Br J Clin Pharmacol*. 2008;65(1):12–21
23. Rainbow J, Browne GJ, Lam LT. Controlling seizures in the prehospital setting: diazepam or midazolam? *J Paediatr Child Health*. 2002;38(6):582–6
24. Schwartz D, Amir L, Dichter R, et al. The use of a powered device for intraosseous drug and fluid administration in a national EMS: a 4-year experience. *J Trauma*. 2008;64(3):650–5
25. Shah I, Deshmukh CT. Intramuscular midazolam vs. intravenous diazepam for acute seizures. *Indian J Pediatr*. 2005;72(8):667–70
26. Shah MI, Macias CG, Dayan PS, et al. An evidence-based guideline for pediatric prehospital seizure management using GRADE methodology. *Prehosp Emerg Care*. 2014;18 Suppl 1:15–24
27. Silbergleit R, Durkalski V, Lowenstein D, et al. Intramuscular versus intravenous therapy for prehospital status epilepticus. *N Engl J Med*. 2012;366(7):591–600
28. Silbergleit R, Durkalski V, Lowenstein D, et al; NETT Investigators. Intramuscular versus intravenous therapy for prehospital status epilepticus. *N Engl J Med*. 2012;366(7):591–600
29. Silbergleit R et al. RAMPART (rapid anticonvulsant medication prior to arrival trial): a double-blind randomized clinical trial of the efficacy of IM midazolam versus IV lorazepam in the pre-hospital treatment of status epilepticus by paramedics. *Epilepsia*. 2011;52 Suppl 8:45–7
30. Sporer KA, Johnson NJ. Detailed analysis of prehospital interventions in medical priority dispatch system determinants. *West J Emerg Med*. 2011;12(1):19–29
31. Sreenath TG, Gupta P, Sharma KK, Krishnamurthy S. Lorazepam versus diazepam-phenytoin combination in the treatment of convulsive status epilepticus in children: a randomized controlled trial. *Eur J Paediatr Neurol*. 2010;14(2):162–8
32. Talukdar B, Chakrabarty B. Efficacy of buccal midazolam compared to intravenous diazepam in controlling convulsions in children: a randomized control trial. *Brain Dev*. 2009;31(10):744–9
33. Vilke GM, Sharieff GQ, Marino A, Gerhart AE, Chan TC. Midazolam for the treatment of out-of-hospital pediatric seizures. *Prehosp Emerg Care*. 2002;6(2):215–7
34. Wuerz RC, Swope GW, Holliman J, Vazquez-de Miguel G. Online medical direction: a prospective study. *Prehosp Disaster Med*. 1995;10(3):51–4