

- community physicians is associated with improved outcome. *Pediatrics*. 2013;112(4):793–9
16. Hartholt KA, van Lieshout EM, Thies WC, Patka P, Schipper IB. Intraosseous devices: a randomized controlled trial comparing three intraosseous devices. *Prehosp Emerg Care*. 2010;14(1):6–13
 17. Howell MD, David AM. Management of sepsis and septic shock. *JAMA*. 2017;317(8):847–8
 18. Hunter CL, Silvestri S, Dean M, Falk JL, Papa L. End-tidal carbon dioxide is associated with mortality and lactate in patients with suspected sepsis. *Am J Emerg Med* (2013) 31, 64–71
 19. Jansen TC, van Bommel J, Mulder PG, Rommes JH, Schieveld SJ, Bakker, J. The prognostic value of blood lactate levels relative to that of vital signs in the pre-hospital setting: a pilot study. *Crit Care*. 2008;12(6): R160
 20. Kumar A, Roberts D, Wood KE, et al. Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med*. 2006;34(6):1589–96
 21. Lampard JG, Lang E. Vasopressors for hypotensive shock. *Ann Emerg Med*. 2013;31(3):351–2
 22. Larsen GY, Mecham N, Greenberg R. An emergency department septic shock protocol and care guideline for children initiated at triage. *Pediatrics*. 2011;127(6): e1585–92
 23. Levy B, Bastien O, Karim B, et al. Experts' recommendations for the management of adult patients with cardiogenic shock. *Ann Intensive Care*. 2015;5(1):17
 24. Lillis KA, Jaffe DM. Prehospital intravenous access in children. *Ann Emerg Med*. 1992;21(12):1430–4
 25. Martin C, Papazian L, Perrin G, Saux P, Gouin F. Norepinephrine or dopamine for the treatment of hyperdynamic septic shock? *Chest*. 1993;103(6):1826–31
 26. Martin C, Viviani X, Leone M, Thirion X. Effect of norepinephrine on the outcome of septic shock. *Crit Care Med*. 2000;28(8):2758–65
 27. Mikkelsen ME, Miltiades AN, Gaieski DF, et al. Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock. *Crit Care Med*. 2009;37(5):1670–7
 28. Morimatsu H, Singh K, Uchino S, Bellomo R, Hart G. Early and exclusive use of norepinephrine in septic shock. *Resuscitation*. 2004;62(2):249–54
 29. Nawrocki PS, Poremba M, Lawner BJ. Push dose epinephrine use in the management of hypotension during critical care transport. *Prehosp Emerg Care*. 2020;24(2):188–95. [//doi.org/10.1080/10903127.2019.1588443](https://doi.org/10.1080/10903127.2019.1588443).
 30. Oliveira CF, Nogueira de Sá FR, Oliveira DS, et al. Time- and fluid-sensitive resuscitation for hemodynamic support of children in septic shock: barriers to the implementation of the American College of Critical Care Medicine/Pediatric Advanced Life Support Guidelines in a pediatric intensive care unit in a developing world. *Pediatr Emerg Care*. 2008;24(12):810–5
 31. Patel GP, Grahe JS, Sperry M, et al. Efficacy and safety of dopamine versus norepinephrine in the management of septic shock. *Shock*. 2010;33(4):375–80
 32. Rhodes A, Evans LE, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med*. Epub 2017 Jan 18
 33. Santhanam I, Sangareddi S, Venkataraman S, Kisson N, Thiruvengadamudayan V, Kasthuri RK. A prospective randomized controlled study of two fluid regimens in the initial management of septic shock in the emergency department. *Pediatr Emerg Care*. 2008;24(10):647–55
 34. Sebat F, Johnson D, Musthafa AA, et al. A multidisciplinary community hospital program for early and rapid resuscitation of shock in nontrauma patients. *Chest*. 2005;127(5):1729–43
 35. Seymour CW, Band RA, Cooke CR, et al. Out-of-hospital characteristics and care of patients with severe sepsis: a cohort study. *J Crit Care*. 2010;25(4):553–62