



8. Haut ER, Kalish BT, Cotton BA, et al. Prehospital intravenous fluid administration is associated with higher mortality in trauma patients: a National Trauma Data Bank analysis. *Ann Surg.* 2011;253(2):371–7
9. Jagoda AS, Bazarian JJ, Bruns JJ Jr, et al; American College of Emergency Physicians; Centers for Disease Control and Prevention. Clinical policy: neuroimaging and decision making in adult mild traumatic brain injury in the acute setting. *Ann Emerg Med.* 2008;52(6):714–48
10. Kleinman ME, Chameides L, Schexnayder SM, et al. Part 14: pediatric advanced life support: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation.* 2010;122: S876–908
11. Reed D. *Adult Trauma Clinical Practice Guidelines: Initial Management of Closed Head Injury in Adults: 2<sup>nd</sup> Edition*. New South Wales Institute of Trauma and Injury Management; 2011
12. Roberts I, Schierhout G. Hyperventilation therapy for acute traumatic brain injury. *Cochrane Database Syst Rev.* 1997;(4):CD000566
13. Spaite DW, Bobrow BJ, Keim SM, et al: Association of Statewide Implementation of the Prehospital Traumatic Brain Injury Treatment Guidelines With Patient Survival Following Traumatic Brain Injury: The Excellence In Prehospital Injury Care (EPIC) Study. *JAMA Surg.* 2019;154(7): e191152. doi:10.1001/jamasurg.2019.1152. NIH Manuscript System ID: NIHMSID: 1663161; PubMed PMID: 31066879; PMCID: PMC6506902
14. Stocchetti N, Maas AIR, Chieriegato A, van der Plas AA. Hyperventilation in head injury a review. *Chest.* 2005;127(5):1812–27
15. Zebrack M, Dandoy C, Hansen K, Scaife E, Mann NC, Bratton SL. Early resuscitation of children with moderate-to-severe traumatic brain injury. *Pediatrics.* 2009;124(1):56–64

#### **Revision Date**

March 11, 2022