



- and preliminary investigation for ratio scale properties. *Pain* 1990;41:139–150
3. Brewster GS, Herbert ME, Hoffman JR. Medical myth: analgesia should not be given to patients with acute abdominal pain because it obscures the diagnosis. *West J Med.* 2000;172(3):209–10
  4. De Nadal M, Munar F, Poca MA, Sahuquillo J, Garnacho A, Rosselló J. Cerebral hemodynamic effects of morphine and fentanyl in patients with severe head injury: absence of correlation to cerebral autoregulation. *Anesthesia.* 2000; 92:1–11
  5. Hicks CL, von Baeyer CL, Spafford P, van Korlaar I, Goodenough B. The Faces Pain Scale – Revised: Toward a common metric in pediatric pain measurement. *Pain.* 2001; 93:173–83
  6. Jennings PA, Cameron P, Bernard S. Ketamine as an analgesic in the pre-hospital setting: a systematic review. *Acta Anaesthesiol Scand.* 2011;55(6):638–43
  7. Lindbeck, George et al. ([Evidence-Based Guidelines for Prehospital Pain Management: Literature and Methods \(nasemso.org\)](https://www.nasemso.org/Evidence-Based-Guidelines-for-Prehospital-Pain-Management-Literature-and-Methods)). Accessed March 11, 2022
  8. LoVecchio F, Oster N, Sturmann K, Nelson LS, Flashner S, Finger R. The use of analgesics in patients with acute abdominal pain. *J Emerg Med.* 1997;15(6):775–9
  9. Manterola C, Astudillo P, Losada H, Pineda V, Sanhueza A, Vial M. Analgesia in patients with acute abdominal pain. *Cochrane Database Syst Rev.* 2007 Jul 18;(3) CD005660
  10. Merkel S, et al. The FLACC: A behavioral scale for scoring postoperative pain in young children., *Pediatr Nurse.* 1997;23(3):293–7
  11. Pace S, Burke TF. Intravenous morphine for early pain relief in patients with acute abdominal pain. *Acad Emerg Med.* 1996;3(12):1086–92
  12. *Prehospital use of Ketamine in Battlefield Analgesia 2012–13.* Falls Church, VA: Defense Health Agency; March 8, 2012. Correspondence to Assistant Secretary of Defense (Health Affairs)
  13. Porter K. Ketamine in prehospital care. *Emerg Med J* 2004; 21:351–4
  14. Powell, Jonathan R., et al. Evidence-Based Guidelines for Prehospital Pain Management: Literature and Methods. *Prehospital Emergency Care.*
  15. Ranji SR, Goldman LE, Simel DL, Shojania KG. Do opiates affect the clinical evaluation of patients with acute abdominal pain? *JAMA.* 2006;296(14):1764–74
  16. Svenson JE, Abernathy MK. Ketamine for prehospital use: new look at an old drug. *Am J Emerg Med.* 2007; 25:977–80
  17. Vermuelen B, Morabia A, Unger PF, et al. Acute appendicitis: influence of early pain relief on the accuracy of clinical and US findings in the decision to operate – a randomized trial. *Radiology.* 1999;210(3):639–43
  18. Wiel E, Zitouni D, Assez N, et al. Continuous infusion of ketamine for out-of-hospital isolated orthopedic injuries secondary to trauma: a randomized controlled trial. *Prehosp Emerg Care.* 2015;19(1):10–16
  19. Wood PR. Ketamine: prehospital and in-hospital use. *Trauma.* 2003;5(2):137–40

#### **Revision Date**

March 11, 2022