



References

1. Abramo TJ, Wiebe RA, Scott SM, Primm PA, McIntyre D, Mydlyer T. Noninvasive capnometry in a pediatric population with respiratory emergencies. *Pediatr Emerg Care*. 1996;12(4):252–4
2. Al-Ansari K, Sakran M, Davidson BL, El Sayyed R, Mahjoub H, Ibrahim K. Nebulized 5% or 3% hypertonic or 0.9% saline for treating acute bronchiolitis in infants. *J Pediatr*. 2010;157(4):630–4
3. Cambonie G, Milési C, Jaber S, et al. Nasal continuous positive airway pressure decreases respiratory muscles overload in young infants with severe acute viral bronchiolitis. *Intensive Care Med*. 2008;34(10):1865–72
4. Chavasse R, Seddon P, Bara A, McKean M. Short acting beta2-agonists for recurrent wheeze in children under two years old. *Cochrane Database Syst Rev*. 2002;(3):CD002873
5. Chowdhury MM, McKenzie SA, Pearson CC, et al. Heliox therapy in bronchiolitis: phase III multicenter double-blind randomized controlled trial. *Pediatrics*. 2013;131(4):661–9
6. Corneli HM, Zorc JJ, Mahajan P, et al. A multicenter, randomized, controlled trial of dexamethasone for bronchiolitis. *N Engl J Med*. 2007;357(4):331–9
7. Denver Metro Airway Study Group. A prospective multicenter evaluation of prehospital airway management performance in a large metropolitan region. *Prehosp Emerg Care*. 2009;13(3):304–10
8. Ehrlich PF, Seidman PS, Atallah O, Haque A, Helmkamp J. Endotracheal intubations in rural pediatric trauma patients. *J Pediatr Surg*. 2004;39(9):1376–80
9. Everard ML, Bara A, Kurian M, Elliot TM, Ducharme F. Anticholinergic drugs for wheeze in children under the age of two years. *Cochrane Database Syst Rev*. 2002;(1):CD001279
10. Freedman SB, Haladyn JK, Floh A, Kirsh JA, Taylor G, Thull-Freedman J. Pediatric myocarditis: emergency department clinical findings and diagnostic evaluation. *Pediatrics*. 2007;120(6):1278–85
11. Gausche-Hill M, Lewis RJ, Stratton SJ, Haynes BE, Gunter CS, Goodrich SM, et al. Effect of out-of-hospital pediatric endotracheal intubation on survival and neurological outcome. *JAMA*. 2000;283(6):783–90
12. Grewal S, Ali S, McConnell DW, Vandermeer B, Klassen TP. A randomized trial of nebulized 3% hypertonic saline with epinephrine in the treatment of acute bronchiolitis in the emergency department. *Arch Pediatr Adolesc Med*. 2009;163(11):1007–12
13. Hartling L, Russell KF, Patel H, Klassen TP, Liang Y. Epinephrine for bronchiolitis. *Cochrane Database Syst Rev*. 2004;(1):CD003123
14. Ho J, Casey B. Time saved with use of emergency warning lights and sirens during response to requests for emergency medical aid in an urban environment. *Ann Emerg Med*. 1998;32(5):585–8
15. Ho J, Lindquist M. Time saved with the use of emergency warning lights and siren while responding to requests for emergency medical aid in a rural environment. *Prehosp Emerg Care*. 2001;5(2):159–62
16. Hunt RC, Brown LH, Cabinum ES, Whitley TW, Prasad NH, Owens JCF, et al. Is ambulance transport time with lights and siren faster than that without? *Ann Emerg Med*. 1995;25(4):507–11
17. Javouhey E, Barats A, Richard N, Stamm D, Floret D. Non-invasive ventilation as primary ventilatory support for infants with severe bronchiolitis. *Intensive Care Med*. 2008;34(9):1608–14
18. Kuzma K, Sporer KA, Michael GE, Youngblood GM. When are prehospital intravenous catheters used for treatment? *J Emerg Med*. 2009;36(4):357–62