



- Neurologic status (GCS, nystagmus, pupil size, focal neurologic deficit, or signs of stroke)
- Blood glucose level

#### **Performance Measures**

- Frequency of performing glucometry
- Time to administration of anticonvulsant medication
- Rate of respiratory failure
- Rate of seizure recurrence

#### **References**

1. Alldredge BK, Gelb AM, Isaacs SM, et al. A comparison of lorazepam, diazepam, and placebo for the treatment of out-of-hospital status epilepticus. *N Engl J Med*. 2001;345(9):631–7
2. Alldredge BK, Wall DB, Ferriero DM. Effect of prehospital treatment on the outcome of status epilepticus in children. *Pediatr Neurol*. 1995;12(3):213–6
3. Appleton R, Sweeney A, Choonara I, Robson J, Molyneux E. Lorazepam versus diazepam in the acute treatment of epileptic seizures and status epilepticus. *Dev Med Child Neurol*. 1995;37(8):682–8
4. Arya R, Gulati S, Kabra M, Sahu JK, Kalra V. Intranasal versus intravenous lorazepam for control of acute seizures in children: a randomized open-label study. *Epilepsia*. 2011;52(4):788–93
5. Bhattacharyya M, Kalra V, Gulati S. Intranasal midazolam vs rectal diazepam in acute childhood seizures. *Pediatr Neurol*. 2006;34(5):355–9
6. Cain E, Ackroyd-Stolarz S, Alexiadis P, Murray D. Prehospital hypoglycemia: the safety of not transporting treated patients. *Prehosp Emerg Care*. 2003;7(4):458–65
7. Chamberlain JM, Altieri MA, Futterman C, Young GM, Ochsenchlager DW, Waisman Y. A prospective, randomized study comparing intramuscular midazolam with intravenous diazepam for the treatment of seizures in children. *Pediatr Emerg Care*. 1997;13(2):92–4
8. Chin RF, Neville BG, Peckham C, Wade A, Bedford H, Scott RC. Treatment of community-onset, childhood convulsive status epilepticus: a prospective, population-based study. *Lancet Neurol*. 2008;7(8):696–703
9. Fisgin T, Gurer Y, Tezic T, et al. Effects of intranasal midazolam and rectal diazepam on acute convulsions in children: prospective randomized study. *J Child Neurol*. 2002;17(2):123–6
10. Frascione RJ, Jensen J, Wewerka SS, Salzman JG. Use of the pediatric EZ-IO needle by emergency medical services providers. *Pediatr Emerg Care*. 2009;25(5):329–32
11. Galustyan SG, Walsh-Kelly CM, Szewczuga D, Bergholte J, Hennes H. The short-term outcome of seizure management by prehospital personnel: a comparison of two protocols. *Pediatr Emerg Care*. 2003;19(4):221–5
12. Holliman CJ, Wuerz RC, Vazquez-de Miguel G, Meador SA. Comparison of interventions in prehospital care by standing orders versus interventions ordered by direct (online) medical command. *Prehosp Disaster Med*. 1994;9(4):202–9
13. Holsti M, Dudley N, Schunk J, et al. Intranasal midazolam vs rectal diazepam for the home treatment of acute seizures in pediatric patients with epilepsy. *Arch Pediatr Adolesc Med*. 2010;164(8):747–53
14. Lahat E, Goldman M, Barr J, Bistrizter T, Berkovitch M. Comparison of intranasal midazolam with intravenous diazepam for treating febrile seizures in children: prospective randomized study. *Br Med J*. 2000;321(7253):83–6