



- West J Emerg Med. 2019 Jul;20(4):688-689]. *West J Emerg Med.* 2019;20(2):409-418. doi:10.5811/westjem.2019.1.41344
8. Halstead SM, Deakyne SJ, Bajaj L, Enzenauer R, Roosevelt GE. The effect of ketamine on intraocular pressure in pediatric patients during procedural sedation. *Acad Emerg Med.* 2012;19(10):1145–50.
 9. Ho JD, Smith SW, Nystrom PC, et al. Successful management of excited delirium syndrome with prehospital ketamine: two case examples. *Prehosp Emerg Care.* 2013;17(2): 274–9.
 10. Isbister GK, Calver LA, Page CB, Stokes B, Bryant JL, Downes MA. Randomized controlled trial of intramuscular droperidol versus midazolam for violence and acute behavioral disturbance: the DORM study. *Ann Emerg Med.* 2010;56(4):392-401 e1.
 11. Kupas DF, Wydro GC. Patient restraint in emergency medical services systems. *Prehosp Emerg Care.* 2002;6(3):340–5.
 12. Sonnier L, Barzman D. Pharmacologic management of acutely agitated pediatric patients. *Paediatr Drugs.* 2011 1;13(1):1–10.
 13. Swift RH, Harrigan EP, Cappelleri JC, Kramer D, Chandler LP. Validation of the behavioural activity rating scale (BARS): a novel measure of activity in agitated patients. *J Psychiatr Res.* 2002;36(2):87–95.
 14. Tsze DS, Steele DW, Machan JT, Akhlaghi F, Linakis JG. Intranasal ketamine for procedural sedation in pediatric laceration repair: a preliminary report. *Pediatr Emerg Care.* 2012;28(8):767–70
 15. *White Paper Report on Excited Delirium Syndrome.* ACEP Excited Delirium Task Force, American College of Emergency Physicians; September 10, 2009.

Revision Date

March 11, 2022