



- Neurological status with vitals: AVPU, GCS
- Exams: Neurological and Mental Status Assessment pre- and post-treatment

### **Performance Measures**

- No oxygen desaturation *less than* 90%
- No hypotension:
  - **Adults:** *less than* 110 mmHg
  - **Pediatrics:**
    - Age *less than* 1 month: *less than* 60 mmHg
    - Age 1–12 months: *less than* 70 mmHg
    - Age 1–10 years: *less than* 70 + 2x age in years
- Assess the patient's blood pressure prior to the administration of any medication that may cause hypotension.
- EtCO<sub>2</sub> target 40 mmHg (range 35–45 mmHg). Meticulous prevention of hypocapnia in all patients
- Triage to the appropriate level hospital within the local trauma system
- **National EMS Quality Alliance (NEMSQA) Performance Measures** (for additional information, see [www.nemsqa.org](http://www.nemsqa.org))
  - *Pediatrics—03: Documentation of Estimated Weight in Kilograms*
  - *Trauma—01: Pain Assessment of Injured Patients*

### **References**

1. Ambrosi PB, Valença MM, Azevedo-Filho H. Prognostic factors in civilian gunshot wounds to the head: a series of 110 surgical patients and brief literature review. *Neurosurg Rev.* 2012;35(3):429–35; discussion 435–6
2. Badjatia N, Carney N, Crocco TJ, et al; Brain Trauma Foundation; BTF Center for Guidelines Management. Guidelines for prehospital management of traumatic brain injury 2<sup>nd</sup> edition. *Prehosp Emerg Care.* 2008;12 Suppl 1: S1–52
3. Berlot G, La Fata C, Bacer B, et al. Influence of prehospital treatment on the outcome of patients with severe blunt traumatic brain injury: a single-centre study. *Eur J Emerg Med.* 2009;16(6):312–17
4. Davis DP, Koprowicz KM, Newgard CD, et al. The relationship between out-of-hospital airway management and outcome among trauma patients with Glasgow Coma Scale scores of 8 or less. *Prehosp Emerg Care.* 2011;15(2):184–92
5. Dumont TM, Visoni AJ, Rughani AI, Tranmer BI, Crookes B. Inappropriate prehospital ventilation in severe traumatic brain injury increases in-hospital mortality. *J Neurotrauma.* 2010 Jul;27(7):233–41
6. Franschman G, Peerdeman SM, Andriessen TM, et al; Amsterdam Lifeline: Analysis of Results and Methods--Traumatic Brain Injury (ALARM-TBI) Investigators. Effect of secondary prehospital risk factors on outcome in severe traumatic brain injury in the context of fast access to trauma care. *J Trauma.* 2011;71(4):826–32
7. Gaither JB, Spaite DW, Bobrow BJ, et al: Impact of Implementing the Prehospital Traumatic Brain Injury Treatment Guidelines: The Excellence In Prehospital Injury Care for Children (EPIC4Kids) Study. *Ann Emerg Med.* 2021;77(2):139–153. DOI: 10.1016/j.annemergmed.2020.09.435. NIH Manuscript System ID: NIHMSID:1654418; PubMed PMID:33187749