

Optimality

Alexander S. Kulikov

Steklov Mathematical Institute at St. Petersburg, Russian Academy of Sciences
and
University of California, San Diego

Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$

Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard

Subset without x and $2x$

Maximizing Profit



A factory produces milk chocolate (\$10 per box) and dark chocolate (\$30 per box). The daily demands are 500 and 200 boxes for milk and dark chocolate, respectively. The factory produces 600 boxes of chocolate per day.

What is the optimum production plan?

Consulting

- A consulting company claims that the maximum profit per day is \$10 000

Consulting

- A consulting company claims that the maximum profit per day is \$10 000
- How can they convince the factory that this profit is indeed **optimal** (i.e., maximum)?

Consulting

- A consulting company claims that the maximum profit per day is \$10 000
- How can they convince the factory that this profit is indeed **optimal** (i.e., maximum)?
- They need to show two things:

Consulting

- A consulting company claims that the maximum profit per day is \$10 000
- How can they convince the factory that this profit is indeed **optimal** (i.e., maximum)?
- They need to show two things:
 1. **Existential statement**: there **exists** a plan achieving profit \$10 000

Consulting

- A consulting company claims that the maximum profit per day is \$10 000
- How can they convince the factory that this profit is indeed **optimal** (i.e., maximum)?
- They need to show two things:
 1. **Existential statement**: there **exists** a plan achieving profit \$10 000
 2. **Universal statement**: **all** plans give profit at most \$10 000

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day
- It is indeed a valid plan:

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day
- It is indeed a valid plan:
 - no more than 500 boxes of milk chocolate

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day
- It is indeed a valid plan:
 - no more than 500 boxes of milk chocolate
 - no more than 200 boxes of dark chocolate

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day
- It is indeed a valid plan:
 - no more than 500 boxes of milk chocolate
 - no more than 200 boxes of dark chocolate
 - no more than 600 boxes

Existential Part: Production Plan

- Plan: produce 400 boxes of milk chocolate and 200 boxes of dark chocolate per day
- It is indeed a valid plan:
 - no more than 500 boxes of milk chocolate
 - no more than 200 boxes of dark chocolate
 - no more than 600 boxes
- Profit: $400 \times 10 + 200 \times 30 = 10\,000$

Universal Part: All Plans are Not Better

- Let M and D be the number of boxes of milk and dark chocolate, respectively, produced per day

Universal Part: All Plans are Not Better

- Let M and D be the number of boxes of milk and dark chocolate, respectively, produced per day
- $M \leq 500, D \leq 200, M + D \leq 600$

Universal Part: All Plans are Not Better

- Let M and D be the number of boxes of milk and dark chocolate, respectively, produced per day
- $M \leq 500$, $D \leq 200$, $M + D \leq 600$
- Want to show that $10M + 30D \leq 10\,000$

Universal Part: All Plans are Not Better

- Let M and D be the number of boxes of milk and dark chocolate, respectively, produced per day
- $M \leq 500$, $D \leq 200$, $M + D \leq 600$
- Want to show that $10M + 30D \leq 10\,000$
- Sum up the inequalities

$$10M + 10D \leq 6\,000$$

$$20D \leq 4\,000$$

Summary

- A proof of the fact that some value α is optimal usually consists of two parts:
 1. Existential statement: there exists a solution achieving the value α
 2. Universal statement: all solutions achieve the value not greater than α
- In this lesson, we'll see several proofs of optimality, following the same pattern

Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$

Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard

Subset without x and $2x$

Problem

What is the maximum number of two-digit integers $(10, 11, \dots, 99)$ that one can select, if it is not allowed to select simultaneously x and y such that $x + y = 100$?

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa
- Similarly, 11 and 89 cannot be taken simultaneously

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa
- Similarly, 11 and 89 cannot be taken simultaneously
- 40 pairs: $(10, 90), (11, 89), \dots, (49, 51)$

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa
- Similarly, 11 and 89 cannot be taken simultaneously
- 40 pairs: $(10, 90), (11, 89), \dots, (49, 51)$
- 10 numbers without pairs:
 $50, 91, 92, \dots, 99$

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa
- Similarly, 11 and 89 cannot be taken simultaneously
- 40 pairs: $(10, 90), (11, 89), \dots, (49, 51)$
- 10 numbers without pairs:
 $50, 91, 92, \dots, 99$
- Definitely no more than $40 + 10 = 50$ numbers

Solving the Problem

- If one takes 10, then one cannot take 90, and vice versa
- Similarly, 11 and 89 cannot be taken simultaneously
- 40 pairs: $(10, 90), (11, 89), \dots, (49, 51)$
- 10 numbers without pairs:
 $50, 91, 92, \dots, 99$
- Definitely no more than $40 + 10 = 50$ numbers
- Optimal solution: $50, 51, \dots, 99$

Once Again, Formally

Theorem

The maximum number of two-digit numbers
s.t. no two of them sum up to 100 is 50.

Once Again, Formally

Theorem

The maximum number of two-digit numbers s.t. no two of them sum up to 100 is 50.

Proof

- Solution of size 50: 50, 51, ..., 99 (the sum of any two is greater than 100)
- Any solution should exclude at least one number from each of the 40 pairs. Hence, at most $90 - 40 = 50$ numbers □

Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$


Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard


Subset without x and $2x$

Chess Rook



A chess **rook** moves vertically and horizontally

Chess Rook



A chess **rook** moves
vertically and horizon-
tally

Maximum Number of Rooks

Problem

What is the maximum number of rooks on a chessboard s.t. no two attack each other?

The Pigeonhole Principle

If n pigeons are placed into m boxes and $m < n$, then there is a box containing more than one pigeon



<https://commons.wikimedia.org/w/index.php?curid=4658682>

Solving the Problem

- There should be at most one rook in each row


Solving the Problem

- There should be at most one rook in each row
- Hence, the total number of rooks is at most 8

Solving the Problem


- There should be at most one rook in each row
- Hence, the total number of rooks is at most 8
- In other words, if the number of rooks is greater than the number of rows, then, by the pigeon hole principle, there is a row containing at least two rooks (and these two rooks attack each other)

Solution




- Placing 8 rooks is not difficult

Solution




- Placing 8 rooks is not difficult

Solution




- Placing 8 rooks is not difficult
- The maximum number of rooks is 8

Solution




- Placing 8 rooks is not difficult
- The maximum number of rooks is 8
- There are many optimum solutions

Solution



- Placing 8 rooks is not difficult
- The maximum number of rooks is 8
- There are many optimum solutions

Solution



- Placing 8 rooks is not difficult
- The maximum number of rooks is 8
- There are many optimum solutions

Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$


Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard


Subset without x and $2x$

Chess Knight



A chess **knight** moves
in an L-shape

Chess Knight




A chess **knight** moves
in an L-shape

Maximum Number of Knights

Problem


What is the maximum number of knights on a chessboard s.t. no two attack each other?

Speculating




- A knight attacks only cells of opposite colors

Speculating




- A knight attacks only cells of opposite colors
- There is a solution with 32 knights: place knights on all white cells

Speculating




- A knight attacks only cells of opposite colors
- There is a solution with 32 knights: place knights on all white cells

Speculating Further




- why can't we place 64 knights?

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further




- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further



- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Speculating Further



- why can't we place 64 knights?
- at most one knight can be placed to these two cells
- if we could partition the cells into 32 disjoint pairs, we would be done!
- let's try...

Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$


Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard


Subset without x and $2x$

Chess Bishop



A chess **bishop** moves diagonally

Chess Bishop




A chess **bishop** moves diagonally

Maximum Number of Bishops


Problem

What is the maximum number of bishops on a chessboard s.t. no two attack each other?

Solving the Problem




Solving the Problem




- partition the board into 15 diagonals

Solving the Problem




- partition the board into 15 diagonals
- each diagonal contains at most one bishop, so at most 15 bishops

Solving the Problem



- partition the board into 15 diagonals
- each diagonal contains at most one bishop, so at most 15 bishops
- but! the diagonals 1 and 15 cannot both contain a bishop, so at most 14

14 Bishops



Outline

Warm-up: Producing Chocolate

Subset without x and $100 - x$

Rooks on a Chessboard

Knights on a Chessboard

Bishops on a Chessboard

Subset without x and $2x$

Problem

What is the maximum number of integers among $1, 2, \dots, 50$ that one can select, if it is not allowed to select simultaneously x and $2x$?

Speculating

- 1 and 2 cannot be taken simultaneously

Speculating

- 1 and 2 cannot be taken simultaneously
- 2 and 4 cannot be taken simultaneously

Speculating

- 1 and 2 cannot be taken simultaneously
- 2 and 4 cannot be taken simultaneously
- More generally, no two neighbors from the following chain can be taken simultaneously: $1 - 2 - 4 - 8 - 16 - 32$

Speculating

- 1 and 2 cannot be taken simultaneously
- 2 and 4 cannot be taken simultaneously
- More generally, no two neighbors from the following chain can be taken simultaneously: 1 – 2 – 4 – 8 – 16 – 32
- Another chain: 3 – 6 – 12 – 24 – 48

Speculating

- 1 and 2 cannot be taken simultaneously
- 2 and 4 cannot be taken simultaneously
- More generally, no two neighbors from the following **chain** can be taken simultaneously: $1 - 2 - 4 - 8 - 16 - 32$
- Another chain: $3 - 6 - 12 - 24 - 48$
- The integers $1, 2, \dots, 50$ can be **partitioned** into such chains, each chain starting with an odd number

Partitioning into Chains

1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	...	49
2	6	10	14	18	22	26	30	34	38	42	46	50				
4	12	20	28	36	44											
8	24	40														
16	48															
32																

- Chains are independent

Partitioning into Chains

1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	...	49
2	6	10	14	18	22	26	30	34	38	42	46	50				
4	12	20	28	36	44											
8	24	40														
16	48															
32																

- Chains are **independent**
- To maximize the number of integers taken in each chain, take every second number in the chain, starting with the first one

Partitioning into Chains

1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	...	49
2	6	10	14	18	22	26	30	34	38	42	46	50				
4	12	20	28	36	44											
8	24	40														
16	48															
32																

- Chains are **independent**
- To maximize the number of integers taken in each chain, take every second number in the chain, starting with the first one
- Optimum size: 33