



NABH ACCREDITED



CARE • COMPASSION • CURE

MGM**NEW BOMBAY HOSPITAL, VASHI**

Plot No. 35, Sector 3, Vashi, Navi Mumbai - 400703. Tel.: (022) 5066 6777



JCI ACCREDITED

| | | |
|-------------|--|---|
| CNS | Emotional State | Anxious (A), Calm (C), Angry (AN), Co-operative (CO), Fearful (F), Restless (R), Withdrawn (W), sedated (S) |
| | LOC | Alert (A), Lethargic (L), Confused (C), Semiconscious (S), Comatose (CO) |
| | Upper Limbs | Preserved (P), Paresis (PA), Plegia (PL), Paraesthesia (PR) |
| | Lower Limbs | Preserved (P), Paresis (PA), Plegia (PL), Paraesthesia (PR) |
| Respiratory | Respiratory Pattern | Normal (N), Dyspnea (D), Tachypnea (T), Bradypnea (B), Orthopnea (O), Kussmaul (K), Cheyne Strokes (CS), Use of Accessory muscle (A) |
| | Oxygenation | Spontaneous (S), Nasal Prongs (N), Mask (M), Endotracheal Tube (ET) Trachestomy (T), Non-Invasive ventilation (N/V), Thermovent (TH) |
| | Pulmonary Auscultation | Air Entry Bilaterally Equal (AEBE) Reduced Right (RR), Reduced Left (RL) |
| | Adventitious Sounds | Ronchi (R), Rales (RA), Crepts (C), |
| | Coughing | Absent (A), Productive (P), Non – Productive (NP) |
| CVS | Pulse | Regular (R), Irregular (I), Palpable (P), Impalpable (IM), Felt by Doppler (D) |
| | Skin colour | Normal (N), Pale (P), Cyanosis (C), Icteric (I) |
| | Capillary refill time | Brisk \leq 2 seconds (B), Sluggish \geq 2 seconds (S) |
| | Oedema | Anasarca (A), Non Pitting (NP) Pitting (P), (1+ 2mm mid, 2+ 4mm moderate, 3+ 6mm moderate fever, 4+ 8mm severe) |
| GI | Mouth | Clean (C), Sordes (S), Halitosis (H) |
| | Appetite | Normal (N), Increased (I), Reduced (R) |
| | Route Of Administration | Oral (O), Nasogastric (N), Parenteral (P), PEG feeds (PEG), Jejunostomy feeds (JT) |
| | Presence Of | Nausea (N), Vomiting (V), Bloating (B) |
| | Abdomen | Flat (F), Distended (D), Tender (T) |
| | Peristalsis | Present (P), Absent (A), Increased (I), Reduced (R) |
| | Elimination | Bowel opened (BO), Constipation (C), Diarrhea (D), Bowel not opened (BNO) |
| | Stool Characteristic | Hard (H), Semisolid (SS), Watery (W), Melena (M), Mucoid (MU), Others frequency (specify) |
| Renal | Urinary Elimination | Retention (R), Incontinence (I), Dysuria (D), Oliguria (O), Polyuria (P), Adequate (A) |
| | Method of Voiding | Self voiding (S), Foleys Catheter (F), Condom Catheter (C) |
| | Urine | Clean (C), Cloudy (CL), Hematuria (H), Sediments (S) |
| Skin | Skin | Intact (I), Cynosis (C), Jaundice (J), Pallor (P), Itching (IT), Petechiae (PE), Ecchymosis (E), Break (B), Blister (BL), Redness (R), Infection (IN) |
| S.J | Oozing | None (N), Sanguinous (SG), Sero Sanguinous (SS), Serous (S), Pus (P), Gaping-Specify size in cm (G), Dressing present (D) |
| Others | Eyes | Clean (C), Discharge (D), Icterus (I) |
| | Sight | Normal (N), Altered (A) |
| | Hearing | Normal (N), Altered (A) |
| | Speech | Aphonia (A), Dysphasia (D), Aphasia (AP), Other- Specify (O) |
| | Isolation precaution | ($\sqrt{\times}$) |
| | Barrier Nursing | ($\sqrt{\times}$) |
| Invasive | Lines | Peripheral (P), Central (C), Arterial (A), Femoral (F) |
| | Catheter | HD catheter (HD), IABP catheter (IA), Urinary Cath (U) |
| | Tubes | Endotracheal tube (E), Tracheostomy (T), Feeding tube (F) |
| | Drains | Drain (D) |
| | Modified Morse Fall risk Scale (Adult) | Low Risk (0 – 24) Medium Risk (25 – 45) High Risk (\geq 45) |
| | Humpty Dumpty Fall risk Scale (Paediatric) | Low Risk (7 – 11) High Risk (12 – 23) |
| | Braden Scale Score | No Risk (19 – 23), Low Risk (15 – 18), Moderate Risk (13 – 14), High Risk (10 – 12), Very High Risk (6 – 9) |
| | Braden Q Scale Score | Low Risk (22 – 28), Moderate Risk (17 – 21), High Risk (\leq 16) |