



## DVT RISK REASSESSMENT

SR No	PARAMETERS	Date	0/1/0	1/2/0	3/0	4/1/0	5/1/0
		Time	9am	9-	9.	9--	9
		Score	1	2	3	4	5
1	Active cancer (on-going treatment or diagnosed within 6 months or palliative care)	0/1	0	0	0	0	0
2	Bedridden recently >3 days or major surgery within four weeks.	0/1	1	2	1	1	1
3	Calf swelling >3 cm compared with asymptomatic side, measured at 10 cm below tibial tubercle (Assess for both legs)	0/1	0	0	0	0	0
4	Collateral (non varicose) superficial veins present (Assess for both legs)	0/1	0	0	0	0	0
5	Entire leg swelling (Assess for both legs)	0/1	0	0	0	0	0
6	Localized tenderness along the deep venous system (Assess for both legs)	0/1	0	0	0	0	0
7	Pitting edema, greater in the symptomatic leg (Assess for both legs)	0/1	0	0	0	0	0
8	Paralysis, paresis, or recent plaster immobilization of the lower extremity (Assess for both legs)	0/1	0	0	2	2	2
9	Previously documented DVT (Assess for both legs)	0/1	0	0	0	0	0
10	Alternative diagnosis to DVT as likely or more likely (Assess for both legs)/Co-morbidity like ESLD / Renal disease, Renal Failure, CCF Cellulitis (Commonly mistaken as DVT), Dependent (stasis) oedema, Lymphatic obstruction, Septic arthritis, cirrhosis, Nephrotic syndrome, Calf muscle tear or strain, Hematoma ( collection of blood) in the muscle, Sprain or rupture of a leg tendon, Fracture	0/-2	0	0	0	0	0
FINAL SCORE			1	2	2	2	
Interpretation of score			mod	mod	mod	mod	mod?
DVT Prophylaxis activated			✓	✓	✓	✓	✓
Initials & Emp. ID			b1670	9-	Ku	BK	
DVT Prophylaxis: For Low risk patient (-2 to 0) Ambulation Moderate Risk : ( 1 to 2) Pharmacological prophylaxis / Mechanical Prophylaxis High risk patients : (3 to 7) Pharmacological prophylaxis along with mechanical Prophylaxis		Mechanical Prophylaxis - Intermittent Pneumatic compressions (IPC) Pharmacological Prophylaxis - Low-Molecular weight heparin Unfractionated Heparin Oral anticoagulant					