



NABH ACCREDITED

**MGM****NEW BOMBAY HOSPITAL, VASHI**

Plot No. 35, Sector 3, Vashi, Navi Mumbai - 400703. Tel.: (022) 5066 6777



JCI ACCREDITED

CNS	Emotional State	Anxious (A), Calm (C), Angry (AN), Co-operative (CO), Fearful (F), Restless (R), Withdrawn (W), sedated (S)
	LOC	Alert (A), Lethargic (L), Confused (C), Semiconscious (S), Comatose (CO)
	Upper Limbs	Preserved (P), Paresis (PA), Plegia (PL), Paraesthesia (PR)
	Lower Limbs	Preserved (P), Paresis (PA), Plegia (PL), Paraesthesia (PR)
Respiratory	Respiratory Pattern	Normal (N), Dyspnea (D), Tachypnea (T), Bradypnea (B), Orthopnea (O), Kussmaul (K), Cheyne Strokes (CS), Use of Accessory muscle (A)
	Oxygenation	Spontaneous (S), Nasal Prongs (N), Mask (M), Endotracheal Tube (ET), Tracheostomy (T), Non-Invasive ventilation (N/V), Thermovent (TH)
	Pulmonary Auscultation	Air Entry Bilaterally Equal (AEBE) Reduced Right (RR), Reduced Left (RL)
	Adventitious Sounds	Ronchi (R), Rales (RA), Crepts (C)
	Coughing	Absent (A), Productive (P), Non-Productive (NP)
CVS	Pulse	Regular (R), Irregular (I), Palpable (P), Impalpable (IM), Felt by Doppler (D)
	Skin colour	Normal (N), Pale (P), Cyanosis (C), Icteric (I)
	Capillary refill time	Brisk ≤ 2 seconds (B), Sluggish ≥ 2 seconds (S)
	Oedema	Anasarca (A), Non Pitting (NP), Pitting (P), (1+ 2mm mid, 2+ 4mm moderate, 3+ 6mm moderate fever, 4+ 8mm severe)
GI	Mouth	Clean (C), Sordes (S), Halitosis (H)
	Appetite	Normal (N), Increased (I), Reduced (R)
	Route Of Administration	Oral (O), Nasogastric (N), Parenteral (P), PEG feeds (PEG), Jejunostomy feeds (JT)
	Presence Of	Nausea (N), Vomiting (V), Bloating (B)
	Abdomen	Flat (F), Distended (D), Tender (T)
	Peristalsis	Present (P), Absent (A), Increased (I), Reduced (R)
	Elimination	Bowel opened (BO), Constipation (C), Diarrhea (D), Bowel not opened (BNO)
	Stool Characteristic	Hard (H), Semisolid (SS), Watery (W), Melena (M), Mucoid (MU), Others frequency (specify)
Renal	Urinary Elimination	Retention (R), Incontinence (I), Dysuria (D), Oliguria (O), Polyuria (P), Adequate (A)
	Method of Voiding	Self voiding (S), Foleys Catheter (F), Condom Catheter (C)
	Urine	Clean (C), Cloudy (CL), Hematuria (H), Sediments (S)
Skin	Skin	Intact (I), Cynosis (C), Jaundice (J), Pallor (P), Itching (IT), Petechiae (PE), Ecchymosis (E), Break (B), Blister (BL), Redness (R), Infection (IN)
S.J	Oozing	None (N), Sanguinous (SG), Sero Sanguinous (SS), Serous (S), Pus (P), Gaping-Specify size in cm (G), Dressing present (D)
Others	Eyes	Clean (C), Discharge (D), Icterus (I)
	Sight	Normal (N), Altered (A)
	Hearing	Normal (N), Altered (A)
	Speech	Aphonia (A), Dysphasia (D), Aphasia (AP), Other- Specify (O)
	Isolation precaution	($\sqrt{\times}$)
	Barrier Nursing	($\sqrt{\times}$)
Invasive	Lines	Peripheral (P), Central (C), Arterial (A), Femoral (F)
	Catheter	HD catheter (HD), IABP catheter (IA), Urinary Cath (U)
	Tubes	Endotracheal tube (E), Tracheostomy (T), Feeding tube (F)
	Drains	Drain (D)
	Modified Morse Fall risk Scale (Adult)	Low Risk (0 - 24) Medium Risk (25 - 45) High Risk (≥ 45)
	Humpty Dumpty Fall risk Scale (Paediatric)	Low Risk (7 - 11) High Risk (12 - 23)
	Braden Scale Score	No Risk (19 - 23), Low Risk (15 - 18), Moderate Risk (13 - 14), High Risk (10 - 12), Very High Risk (6 - 9)
	Braden Q Scale Score	Low Risk (22 - 28), Moderate Risk (17 - 21), High Risk (≤ 16)