



**MGM**  
**NEW BOMBAY HOSPITAL, VASHI**



MR. KIRAN BHAGWAN TAWARE  
UHID : MGM240017131 IPD : MGMIP2406720  
Age/Sex : 56 Years / M DOA : 05/10/2024  
BED : SICU-010 WARD : SICU  
DR : DR. PRASHANT ATHALE

## FALL RISK ASSESSMENT PHYSIOTHERAPY

### FALL RISK ASSESSMENT ( MODIFIED MORSE SCALE ) :

Variables	Numeric Value		CNS / CVS Medication	No Yes	
History of Falling	No	0	Gait		0
	Yes	25	Normal / bed rest / wheel chair		0
Secondary Diagnosis / Elimination Problem	No	0	Weak		10
	Yes	15	Impaired		20
Ambulatory Aid			Mental Status		
None / bed rest / nurse assist		0	Oriented to own stability		0
Crutches / Cane / Walker		15	Overestimated or forgets limitations		15
Furniture		30	Total Score		35

### FALL RISK ASSESSMENT ( THE HUMPTY DUMPTY SCALE )

Parameter	Criteria	Score	
Age	Less than 3 years old	4	Total score : _____ 7-11 : Low risk 12 or above : High risk
	3 to less than 7 years old	3	
	7 to less than 13 years old	2	
	13 years and older	1	
Gender	Male	2	Interventions for low risk ● Orient the patient / attendant to surroundings ● Bed is in low position and Bed locked ● Orient the patient / attendant on use of call bell ● Keep all side rails up ● Patient instructed not to get up without assistance ● Assistant with patient while using washroom ● Place the yellow colour ID band on bed
	Female	1	
Diagnosis	Neurological Diagnosis	4	
	Alterations in oxygenation (respiratory diagnosis, dehydration, anaemia, anorexia, syncope / dizziness, etc)	3	
	Psychiatric behavioural disorders	2	Interventions for High risk ● Family companion instructed to be present at bedside ● Need for 24 hrs. Supervision / Assistance ● Initiate the use of bedside commode if appropriate ● Place the red colour ID band on bed
	Others diagnosis	1	
Cognitive Impairment	Not aware of limitation	3	
	Forgets limitations	2	
	Oriented	1	
Environmental factors	History of falls or infant - toddler placed in bed	4	
	Patient uses assertive devices or infant toddler in crib or furniture / lighting	3	
	Patient placed in bed	2	
	Outpatient Area	1	
Response to surgery / sedation and anaesthesia	Within 24 Area	3	
	Within 48 hours	2	
	More than 48 hours / none	1	
Medication Usage	Multiple usage of : sedatives (excluding ICU patients sedated and paralyzed) Hypnotic, Barbiturates, Phenothiazine, Antidepressants, Laxatives diuretics, Narcotics	3	
	One of the medicines listed above	2	
	Other medications / none	1	